

## Subjective evaluation of physical activity

1. Kaur M, Ribeiro DC, Lamb P, et al. Low knee-related quality of life and persistent physical asymmetries in participants up to 10 years post-ACL reconstruction-A cross-sectional study. *Phys Ther Sport* 2021;48:35-42.
2. Sonesson S, Österberg A, Gauffin H, et al. Low correlation between functional performance and patient reported outcome measures in individuals with non-surgically treated ACL injury. *Phys Ther Sport* 2021;47:185-92.
3. Heijne A, Ang BO, Werner S. Predictive factors for 12-month outcome after anterior cruciate ligament reconstruction. *Scand J Med Sci Sports* 2009;19:842-9.
4. Tengman E, Brax Olofsson L, Nilsson KG, et al. Anterior cruciate ligament injury after more than 20 years: I. Physical activity level and knee function. *Scand J Med Sci Sports* 2014;24:e491-500.
5. MOON Knee Group. Ten-year outcomes and risk factors after anterior cruciate ligament reconstruction: A MOON longitudinal prospective cohort study. *Am J Sports Med* 2018;46:815-25.
6. Spindler KP, Huston LJ, Wright RW, et al. The prognosis and predictors of sports function and activity at minimum 6 years after anterior cruciate ligament reconstruction: A population cohort study. *Am J Sports Med* 2011;39:348-59.
7. Cox CL, Huston LJ, Dunn WR, et al. Are articular cartilage lesions and meniscus tears predictive of IKDC, KOOS, and Marx activity level outcomes after anterior cruciate ligament reconstruction? A 6-year multicenter cohort study. *Am J Sports Med* 2014;42:1058-67.
8. Kostogiannis I, Ageberg E, Neuman P, et al. Activity level and subjective knee function 15 years after anterior cruciate ligament injury: A prospective, longitudinal study of nonreconstructed patients. *Am J Sports Med* 2007;35:1135-43.
9. Lee DY, Karim SA, Chang HC. Return to sports after anterior cruciate ligament reconstruction - A review of patients with minimum 5-year follow-up. *Ann Acad Med Singap* 2008;37:273-8.
10. Fältström A, Hägglund M, Kvist J. Patient-reported knee function, quality of life, and activity level after bilateral anterior cruciate ligament injuries. *Am J Sports Med* 2013;41:2805-13.
11. Kessler MA, Behrend H, Henz S, et al. Function, osteoarthritis and activity after ACL-rupture: 11 years follow-up results of conservative versus reconstructive treatment. *Knee Surg Sports Traumatol Arthrosc* 2008;16:442-8.
12. Månsson O, Sernert N, Rostgard-Christensen L, et al. Long-term clinical and radiographic results after delayed anterior cruciate ligament reconstruction in adolescents. *Am J Sports Med* 2015;43:138-45.
13. Minzlaff P, Heidt T, Feucht MJ, et al. Patient satisfaction with health is substantially improved following ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2018;26:582-8.

14. Möller E, Weidenhielm L, Werner S. Outcome and knee-related quality of life after anterior cruciate ligament reconstruction: A long-term follow-up. *Knee Surg Sports Traumatol Arthrosc* 2009;17:786-94.
15. Klasan A, Putnis SE, Grasso S, et al. Tegner level is predictive for successful return to sport 2 years after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2021;29:3010-6.
16. Gignac MA, Cao X, Ramanathan S, et al. Perceived personal importance of exercise and fears of re-injury: A longitudinal study of psychological factors related to activity after anterior cruciate ligament reconstruction. *BMC Sports Sci Med Rehabil* 2015;7:4.

### **Lower extremity injuries, including hip or ankle**

1. Hubbard-Turner T, Turner MJ. Physical activity levels in college students with chronic ankle instability. *J Athl Train* 2015;50:742-7.
2. Pan F, Byrne KS, Ramakrishnan R, et al. Association between musculoskeletal pain at multiple sites and objectively measured physical activity and work capacity: Results from UK Biobank study. *J Sci Med Sport* 2019;22:444-9.
3. Kretzschmar M, Lin W, Nardo L, et al. Association of physical activity measured by accelerometer, knee joint abnormalities, and cartilage T2 measurements obtained from 3T magnetic resonance imaging: Data from the osteoarthritis initiative. *Arthritis Care Res (Hoboken)* 2015;67:1272-80.
4. Thanoo N, Gilbert AL, Trainor S, et al. The relationship between polypharmacy and physical activity in those with or at risk of knee osteoarthritis. *J Am Geriatr Soc* 2020;68:2015-20.
5. Brisson NM, Gatti AA, Maly MR. Association of pain and steps per day in persons with mild-to-moderate, symptomatic knee osteoarthritis: A mixed-effects models analysis of multiple measurements over three years. *Arthritis Care Res (Hoboken)* 2020;72:114-21.
6. Sliepen M, Mauricio E, Lipperts M, et al. Objective assessment of physical activity and sedentary behaviour in knee osteoarthritis patients - beyond daily steps and total sedentary time. *BMC Musculoskelet Disord* 2018;19:64.
7. Whitaker KM, Pettee Gabriel K, Laddu D, et al. Bidirectional associations of accelerometer measured sedentary behavior and physical activity with knee pain, stiffness, and physical function: The CARDIA study. *Prev Med Rep* 2021;22:101348.
8. Ceroni D, Martin X, Lamah L, et al. Recovery of physical activity levels in adolescents after lower limb fractures: A longitudinal, accelerometry-based activity monitor study. *BMC Musculoskelet Disord* 2012;13:131.
9. Whittaker JL, Toomey CM, Nettel-Aguirre A, et al. Health-related outcomes after a youth sport-related knee injury. *Med Sci Sports Exerc* 2019;51:255-63.

10. Glaviano NR, Baellow A, Saliba S. Physical activity levels in individuals with and without patellofemoral pain. *Phys Ther Sport* 2017;27:12-6.

### **Did not estimate relevant outcomes**

1. Bailey CA, Bardana DD, Costigan PA. Using an accelerometer and the step-up-and-over test to evaluate the knee function of patients with anterior cruciate ligament reconstruction. *Clin Biomech (Bristol, Avon)* 2016;39:32-7.

2. Havens KL, Cohen SC, Pratt KA, et al. Accelerations from wearable accelerometers reflect knee loading during running after anterior cruciate ligament reconstruction. *Clin Biomech (Bristol, Avon)* 2018;58:57-61.

3. Tomite T, Saito H, Kijima H, et al. Evaluation of anteroposterior accelerometric change after bi-cruciate stabilized total knee arthroplasty and posterior stabilized total knee arthroplasty. *Knee* 2021;32:121-30.

4. Bryant AL, Newton RU, Steele J. Successful feed-forward strategies following ACL injury and reconstruction. *J Electromyogr Kinesiol* 2009;19:988-97.

5. Ezzat AM, Brussoni M, Mâsse LC, et al. New or recurrent knee injury, physical activity, and osteoarthritis beliefs in a cohort of female athletes 2 to 3 years after ACL reconstruction and matched healthy peers. *Sports Health* 2022;14:842-8.

6. Fischer F, Blank C, Dünwald T, et al. Isokinetic extension strength is associated with single-leg vertical jump height. *Orthop J Sports Med* 2017;5:2325967117736766.

7. Dunn WR, Spindler KP; MOON Consortium. Predictors of activity level 2 years after anterior cruciate ligament reconstruction (ACLR): A multicenter orthopaedic outcomes network (MOON) ACLR cohort study. *Am J Sports Med* 2010;38:2040-50.

8. De Carlo MS, Sell KE. The effects of the number and frequency of physical therapy treatments on selected outcomes of treatment in patients with anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 1997;26:332-9.

9. Lisee C, Davis-Wilson HC, Evans-Pickett A, et al. Linking gait biomechanics and daily steps after ACL reconstruction. *Med Sci Sports Exerc* 2022;54:709-16.

10. Pamukoff DN, Pietrosimone B, Ryan ED, et al. Whole-body vibration improves early rate of torque development in individuals with anterior cruciate ligament reconstruction. *J Strength Cond Res* 2017;31:2992-3000.

11. Sole G, Lamb P, Pataky T, et al. Immediate and 6-week effects of wearing a knee sleeve following anterior cruciate ligament reconstruction: A cross-over laboratory and randomised clinical trial. *BMC Musculoskelet Disord* 2021;22:655.

## Not an original study

1. Abrams GD, Harris JD, Gupta AK, et al. Functional performance testing after anterior cruciate ligament reconstruction: A systematic review. *Orthop J Sports Med* 2014;2:2325967113518305.
2. Kuenze C, Collins K, Pfeiffer KA, et al. Assessing physical activity after ACL injury: Moving beyond return to sport. *Sports Health* 2022;14:197-204.
3. Brown C, Marinko L, LaValley MP, et al. Quadriceps strength after anterior cruciate ligament reconstruction compared with uninjured matched controls: A systematic review and meta-analysis. *Orthop J Sports Med* 2021;9:2325967121991534.
4. Zalta J. Massage therapy protocol for post-anterior cruciate ligament reconstruction patellofemoral pain syndrome: a case report. *Int J Ther Massage Bodywork* 2008;1:11-21.
5. Darware M, Naqvi WM. A case report on physiotherapy rehabilitation accelerating the recovery of older patient with anterior cruciate ligament reconstruction. *Med Sci* 2020;24:1803-8.
6. de Oliveira FCL, Roy JS, Pappas E. ACL injury, physical activity, and overweight/obesity: a vicious cycle? *Knee Surg Sports Traumatol Arthrosc* 2020;28:667-9.
7. Everhart JS, Yalcin S, Spindler KP. Twenty-year outcomes after anterior cruciate ligament reconstruction: A systematic review of prospectively collected data. *Am J Sports Med* 2022;50:2842-52.
8. Arnold JB, Walters JL, Ferrar KE. Does physical activity increase after total hip or knee arthroplasty for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther* 2016;46:431-42.
9. Wallis JA, Webster KE, Levinger P, et al. What proportion of people with hip and knee osteoarthritis meet physical activity guidelines? A systematic review and meta-analysis. *Osteoarthr Cartil* 2013;21:1648-59.
10. Ekegren CL, Beck B, Climie RE, et al. Physical activity and sedentary behavior subsequent to serious orthopedic injury: A systematic review. *Arch Phys Med Rehabil* 2018;99:164-77.
11. Butler CR, Allen K, DiStefano LJ, et al. Protracted cardiovascular impairments after anterior cruciate ligament injury: A critically appraised topic. *J Sport Rehabil* 2019;29:680-3.
12. Yang L, Wang P, McGill B. The relationship between experience of knee pain and physical activity participation: A scoping review of quantitative studies. *Int J Nurs Sci* 2023;10:258-67.
13. Lisee C, Lepley AS, Birchmeier T, et al. Quadriceps strength and volitional activation after anterior cruciate ligament reconstruction: A systematic review and meta-analysis. *Sports Health* 2019;11:163-79.

14. Sašek M, Kozinc Ž, Löfler S, et al. Objectively measured physical activity, sedentary behavior and functional performance before and after lower limb joint arthroplasty: A systematic review with meta-analysis. *J Clin Med* 2021;10:5885.
15. Barchek AR, Baez SE, Hoch MC, et al. The relationship between musculoskeletal injury and objectively measured physical activity levels: A critically appraised topic. *J Sport Rehabil* 2020;29:243-47.
16. Oliveira CB, Franco MR, Maher CG, et al. Physical activity interventions for increasing objectively measured physical activity levels in patients with chronic musculoskeletal pain: A systematic review. *Arthritis Care Res* 2016;68:1832-42.
17. Le CY, Hui C, Emery CA, et al. Every athlete has a unique trajectory: Knee-related quality of life in young athletes following anterior cruciate ligament reconstruction. 2019.
18. Lowenhar MC. Femoral cartilage resiliency in individuals with anterior cruciate ligament reconstruction and healthy controls (Doctoral dissertation, The University of North Carolina at Chapel Hill). 2021.
19. Barber-Westin SD, Noyes FR. Factors used to determine return to unrestricted sports activities after anterior cruciate ligament reconstruction. *Arthroscopy* 2011;27:1697-705.

### **Only abstracts available**

1. Hall M, Wang X, Fortin K, et al. Moderate to vigorous physical activity is associated with greater patellar cartilage volume over 2 years following anterior cruciate ligament reconstruction. *Osteoarthr Cartil* 2016; 24:S195-S196.
2. Soutakbar H, Scammell BE, Hodgson L, et al. Are there any interactions between physical activity, obesity and injury on the risk of incidence or progression of knee osteoarthritis? Data from OAI and MOST. *Osteoarthr Cartil* 2016;24:S195.
3. Cheung C, Wyman J, Resnick B. Is yoga effective for knee osteoarthritis in older women? *Osteoarthr Cartil* 2012;20:S280.