






# Self-reported reproductive health of retired elite women's footballers: a cross-sectional study

Sean Carmody <sup>1,2</sup> Steve den Hollander <sup>3</sup> Kirsty Elliott-Sale <sup>4</sup>  
Margo Lynn Mountjoy <sup>5,6,7</sup> Jane S Thornton <sup>8</sup> Andrew Massey,<sup>9</sup>  
Gino Kerkhoffs,<sup>1,2,10,11</sup> Vincent Gouttebarge <sup>1,2,3,10,11,12</sup>

**To cite:** Carmody S, den Hollander S, Elliott-Sale K, *et al.* Self-reported reproductive health of retired elite women's footballers: a cross-sectional study. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e002028. doi:10.1136/bmjsem-2024-002028

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjsem-2024-002028>).

Accepted 11 July 2024

## ABSTRACT

**Objectives** The primary objective of this study was to describe the self-reported reproductive health of retired elite women's footballers with specific reference to menstrual function, pregnancy and motherhood, contraceptive use and pelvic floor function.

**Methods** An electronic survey was disseminated to women's footballers (18 years or older) who had retired from elite football within 10 years of completing the survey.

**Results** 69 respondents completed the survey (mean age 35.8 years, mean age at retirement 30.3 years). One-third of participants self-reported experiencing at least one episode of amenorrhoea (>3 months without menstruation) for reasons other than hormonal contraceptive use or pregnancy. Three participants (mean age of 41 years, range 30–54) reported having reached menopause at the time of the study. 54 (78.3%) participants were not using any form of contraception. 17 (24.6%) of the participants are mothers (range 1–3 children). 51 of the participants (73.9%) had never been pregnant, and the majority of those who had been pregnant (86.7%) became pregnant in less than 2 years following the onset of desire for pregnancy. Four of the retired players gave birth during their playing career, and the mean time to return to competitive matches following delivery was 22 weeks. The mean Pelvic Floor Distress Inventory-20 score for participants was 72.

**Conclusion** These insights can be used to inform future efforts to promote positive reproductive health outcomes for current, former and future women's footballers. Research efforts should focus on improving the understanding of how to effectively support women's footballers in the perinatal period. Best practice guidelines on the use of menstrual cycle monitoring and pelvic health support would improve standards of care for women's footballers. Stakeholders should consider gender-specific postretirement care for women's footballers.

## INTRODUCTION

Elite women's footballers often compete during their peak reproductive years, in some cases compelling them to choose between motherhood and their football career. This professional and personal dilemma can be due to a combination of sport-related

### WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Elite women athletes often compete during their peak reproductive years and this has potential implications for their reproductive health.
- ⇒ There are high rates of amenorrhoea among elite women athletes, and this is likely to be associated with negative health outcomes.

### WHAT THIS STUDY ADDS

- ⇒ This study provides insight into the self-reported reproductive health outcomes of retired elite women's footballers with specific reference to menstrual function, pregnancy and motherhood, contraceptive use and pelvic floor function.

### HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ The information presented here can be used to inform future efforts to promote positive reproductive health outcomes for current, former and future women's footballers.

perinatal and postpartum considerations, a lack of widespread maternity policies or a lack of contractual security.<sup>1–5</sup> While there is an increasing research interest in the relationship between the menstrual cycle and its relationship with health and performance in women's footballers, the understanding of more detailed reproductive outcomes in elite women's footballers is limited, both during and after a player's career.<sup>6 7</sup>

A recent scoping review<sup>8</sup> highlighted that there are very few studies examining long-term health outcomes among retired elite women's footballers, which reflects the broader research landscape in women's sport (including football).<sup>9 10</sup> A small number of studies have investigated musculoskeletal<sup>11 12</sup> and mental health<sup>13–15</sup> conditions among retired women's professional footballers, but there are no studies examining female-specific health considerations in this population. These female-specific



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

### Correspondence to

Dr Sean Carmody;  
seanocearmaide@gmail.com

considerations include (but are not limited to) Relative Energy Deficiency in Sport (REDs), fertility and pregnancy, contraceptive use, pelvic floor dysfunction and menstrual cycle disorders.

Female athletes (with or without REDs) can experience menstrual dysfunction, with unknown impacts on long-term fertility or pregnancy outcomes.<sup>1 16–19</sup> A study among retired elite women rowers and rugby players (n=74) reported high rates of amenorrhoea, older age at childbirth and earlier onset of menopause suggesting a narrower 'reproductive window' for female athletes.<sup>20</sup> One study (n=34) identified positive pregnancy outcomes in a cohort of active elite athletes, with athletes getting pregnant easily, delivering healthy babies and returning to sport post partum.<sup>3</sup> There is some understanding of hormonal contraceptive use among active women's footballers, with one study showing that about a third of players at the highest level in England use hormonal contraception.<sup>21</sup> Further studies are needed to understand the reasons for hormonal contraceptive use in women's footballers, including the risks and benefits from a health and performance perspective. With respect to pelvic floor health, athletes appear to be at higher risk of pelvic floor dysfunction than the general population, and this can lead to psychological distress.<sup>22</sup> Athletes who have given birth are likely to be at greater risk of pelvic floor dysfunction.<sup>23</sup>

An improved understanding of female-specific health considerations among retired elite women's footballers would empower key stakeholders (eg, sporting organisations, governing bodies, athlete health and performance teams) to support athletes during and after their career.<sup>24</sup> This could be done by putting preventative measures in place where any potential concerns or risks have been identified. In particular, greater knowledge with respect to reproductive health outcomes (eg, menstrual function, fertility) among retired elite women's footballers may help to promote better provisions for players who would like to have children during or soon after their career. Therefore, the primary objective of this study was to describe the self-reported reproductive health of retired elite women's footballers with specific reference to menstrual function, pregnancy and motherhood, contraceptive use, and pelvic floor function. Secondary objectives of this study were to explore the likelihood of a history of a stress fracture among retired players whose menstrual period had stopped for 3 months or more (for reasons other than contraception or pregnancy), and to compare pelvic floor health outcome measures between retired players who had given birth against those who had not.

## METHODS

### Study design

An observational study based on a cross-sectional design through means of an electronic survey was conducted. In order to ensure a high quality of reporting, the Strengthening the Reporting of Observational Studies

in Epidemiology statement and Checklist for Reporting Results of Internet E-Surveys were adhered to.<sup>25–27</sup>

### Participants

Inclusion criteria were women, 18 years or older, retired from elite football within the 10 years prior to completing the survey and able to understand text written in English. Participants were excluded if they had retired more than 10 years ago, or if they had not played an elite level of football according to the study's inclusion criteria, which was defined as an individual who competed at the senior international level and/or in the highest or second highest national domestic league, which would be considered tier 4 or tier 5 according to standard participant classification criteria.<sup>28</sup> Remuneration for playing football was not considered a prerequisite for participation in the study due to the largely semiprofessional/amateur landscape of elite women's football until relatively recent advancements. A sample of retired elite women's footballers was recruited via convenience and snowball sampling. Participants were recruited through a variety of means including invitations from national governing bodies and players' unions (national and international), 'word of mouth' and social media recruitment.

### Patient and public involvement

One retired professional women's player was invited to complete the study and to provide feedback on its content and design prior to wider dissemination. The retired player gave feedback on survey design, wording of questions and proposed including an additional question.

### Outcome measures

#### Menstrual function

Seven questions were included to assess menstrual function. Age of menarche was assessed with a single question (eg, 'At about what age did you begin to menstruate?'). Participants were asked whether they have reached menopause ('Have you reached menopause?' YES/NO). Menopause was not defined for the purposes of the question. Mean menstrual cycle length and duration of menstruation were measured in a number of days. Participants were invited to describe the regularity of their cycle as either regular, irregular or very irregular according to their perception. Participants were asked about their experience of pain during their menstrual cycle, with options to specify whether the pain occurred before, during or after menstruation. Participants were asked to recall whether following menarche, they had ever experienced their periods stop for three or more consecutive months (for reasons other than hormonal contraceptive use or pregnancy) with YES/NO answer options.

#### Contraception use and hormonal replacement

Two questions assessed contraception and hormonal replacement (eg, hormone replacement therapy for menopause) use. Participants were asked about their current use of contraception use (including listed options

for barrier and hormonal methods), with 10 options available to choose from. Postmenopausal hormonal replacement use was assessed ('Are you currently using hormone replacement?') with YES/NO answer options.

### Pregnancy and motherhood

A total of 11 questions were included that evaluated pregnancy and motherhood. Motherhood and pregnancy were initially assessed with the questions ('Are you a mother?'; 'Have you been pregnant?') with YES/NO answer options. Follow-up questions included 'How many children do you have?', 'How did you give birth?' (Vaginal delivery; Caesarean delivery). Participants were asked to report the duration of time it took them to become pregnant following the onset of desire for pregnancy; 'How long on average did it take you to become pregnant following the onset of desire for pregnancy?' (Less than 1 year; 1–2 years; more than 2 years). The number of previous pregnancies and pregnancy losses were recorded. For participants who responded that they had previously given birth, they were asked whether they had given birth during their playing career. If they answered 'YES', they were asked 'How long did it take to return to training?' (number of weeks) and 'How long did it take to return to competitive matches?' (number of weeks).

### Pelvic floor dysfunction

The Pelvic Floor Distress Inventory (PFDI-20) was used to assess pelvic floor function among this cohort of retired elite women's footballers. The PFDI-20 is a 20-item questionnaire divided into three subscales: six items evaluating Pelvic Organ Prolapse Distress (POPDI-6), 8 items assessing Colorectal anal Distress (CRADI-8) and six items evaluating Urinary Distress (UDI-6).<sup>29</sup> Each question relates to the presence of an individual symptom. An example question includes 'do you usually experience frequent urination?'. If the symptom is present, the responder scores it on a 4-point scale to indicate the extent of distress; score 1 is not at all, and 4 is quite a bit. POPDI-6, CRADI-8 and UDI-6 are each scored similarly. First, subscale scores for each scale are obtained by taking the mean value of all items answered within each subscale. These subscale scores are determined by multiplying each scale's mean score by 25, which gives each subscale score a range of 0–100. Therefore, the total scale score is calculated by summing the subscale scores (a range of 0–300). A higher score is suggestive of more symptoms and higher levels of distress due to the symptoms.

### Procedures

An online questionnaire was developed in English using an electronic data management system (Castor EDC, www.castoredc.com). Eligible patients received an invitation by email between November 2021 and July 2023, followed by periodic reminders at random intervals. This questionnaire included the following descriptive

variables: age, height, body weight, duration of elite football career, level of play, nature and duration of retirement, level of education, and current employment status. Information about the study was sent to potential participants. All participants gave their electronic informed consent and filled in the questionnaire. Players participated voluntarily in the study and did not receive any financial remuneration for their participation. Participants were provided with a free copy of the book 'The Secret Balance of Champions' as a show of gratitude for completing the survey.<sup>30</sup> The questionnaire has been included in online supplemental material.

### Statistical analysis

All data were self-reported. The statistical software IBM SPSS V.28.0 for Apple Mac was used for data analysis. Descriptive analyses (mean, SD, frequency and range) were performed for all variables, using percentages for categorical variables and means and SD for continuous variables. Pearson correlation coefficient ( $r$ ) was used to assess the association of amenorrhoea and a history of stress fracture ( $r$  value less than 0.3 was considered small; 0.3–0.5 moderate; 0.5–0.7 large; 0.7–0.9 very large and >0.9 near perfect for predicting relationships).<sup>31</sup> The Mann-Whitney test (independent samples) was used to determine normal distribution and make comparisons between groups, respectively.<sup>32</sup> Associations with  $p \leq 0.05$  were considered to be statistically significant.

## RESULTS

### Demographics

69 respondents completed the survey in full and were included in the final analysis. The mean age of participants was 35.8 years (SD=5.7, range 24–54) while their mean age at retirement was 30.3 years. The mean number of seasons as an elite footballer was 10.6 seasons (SD=5.2, range 2–21 years). The mean duration of retirement at the time of survey was 5.5 years (SD=5.4). 64 (92.8%) participants reported playing in the highest national league, and 79% ( $n=53$ ) played senior international football. Eighteen different countries (across six continents) were represented in the survey, with England representing the highest number of participants ( $n=15$ , 21.7%). Over one-third ( $n=26$ , 37.7%) retired from elite football due to injury. Participant characteristics are presented in [table 1](#).

### Menstrual function

The mean age of menarche among respondents was 14 years (SD=2). The mean duration of menstrual cycle was 23.7 days, with 4 days of menstrual bleeding. 11 participants (15.9%) subjectively described their menstrual cycle as irregular, with 6 (8.7%) describing their menstrual cycle as very irregular. One-third ( $n=23$ ) of participants experienced at least one occasion of their period stopping for 3 months or more for reasons other than hormonal contraception or pregnancy. 16 of the participants (23.2%) had been diagnosed with a stress



**Table 1** Demographic information of survey respondents (n=69)

Current age (years)	35.8 mean, 5.7 SD, range 23–54
Height (cm)	167 mean, range 146–182
Weight (kg)	65.2, range 50–96
Age at retirement (years)	30.3 mean
Currently in employment:	
Yes	n=59, 85.5%
Number of years since retirement	5.5 (SD=5.4)
Number of years as a professional footballer	10.6 mean, range 2–22
Reason for retirement (more than one option available for selection):	
Injury	n=26, 37.7%
Personal choice	n=41, 59.4%
Financial reasons	n=11, 15.9%
Other	n=7, 10.1%
Highest level of education:	
No school completed	n=0, 0%
Nursery/elementary school	n=0, 0%
High school	n=9, 13%
Vocational/technical school	n=1, 1.4%
College, university or equivalent	n=59, 85.5%
Position (more than one option available for selection):	
Goalkeeper	n=8, 11.6%
Defender	n=34, 49.3%
Midfielder	n=23, 33.3%
Forward	n=11, 15.9%
Level of play:	
Highest national league	n=64, 92.8%
Second highest national league	n=3, 4.4%
Other	n=1, 1.4%
Played international football	n=53, 76.8%

fracture at some point during their football career. There was a weak correlation ( $r=0.1$ ) between a history of amenorrhoea for 3 months or more and having sustained a stress fracture. Three participants (mean age of 41 years, range 30–54) reported having reached menopause at the time of the study. The findings are summarised in [table 2](#).

### Contraception use and hormonal replacement

54 (78%) participants were not using any form of contraception at the time of the study. Three (4%) were using a form of contraceptive pill and 5 (7%) were using a contraceptive implant. No participants were using hormonal

**Table 2** Menstrual function of participants (n=69)

Age of menarche (years)	14 (SD=2), range 10–20
Mean duration of menstrual cycle (days)	23.7
Mean number of days of bleeding during menstrual cycle	4 days
Reached menopause:	
Yes	n=3, 4.4%
Perception of menstrual cycle regularity:	
Regular	n=46, 66.7%
Irregular	n=11, 15.9%
Very irregular	n=6, 8.7%
Did not say	n=6, 8.7%
Experience pain during menstrual cycle:	
No	n=17, 24.6%
Yes, before a bleeding period	n=16, 23.2%
Yes, during a bleeding period	n=29, 42%
Yes, after a bleeding period	n=0
Did not say	n=7, 10.1%
Period stopped for 3 months or more (for reasons other than pregnancy or contraception):	
Yes	n=23, 33.3%

replacement therapy. The findings are summarised in [table 3](#).

### Pregnancy and motherhood

17 (26%) of the participants are mothers (range=1–3 children) (note motherhood does not imply previous pregnancy). 51 of the participants (77%) had never been pregnant. For those who had been pregnant (n=15), nine became pregnant in less than 1 year following the onset of desire for pregnancy, four became pregnant within 1–2 years and two became pregnant over 2 years after the onset of desire for pregnancy. Eight out of 15 participants who had been pregnant became pregnant

**Table 3** Contraceptive and hormonal replacement use among participants (n=69)

Contraception:	
None	n=54, 78.2%
Contraceptive pill	n=3, 4.4%
Condom	n=2, 2.9%
Contraceptive implant	n=5, 7.3%
Other	n=4, 5.8%
Did not say	n=2, 2.9%
Hormone replacement therapy	n=0

**Table 4** History of pregnancy and motherhood amongst participants (n=69)

Number of participants who are mothers	17, 24.6%, range 1–3 children
Have you ever been pregnant:	
Yes	n=15, 21.7%
No	n=51, 73.9%
Did not say	n=3, 4.4%
Pregnancy onset desire (n=15 participants):	
Became pregnant in less than 1 year	n=9, 60%
Became pregnant in 1–2 years	n=4, 26.7%
Became pregnant in 2+ years	n = 2, 13.3%
Number of pregnancies (n=15 participants):	
Total number of pregnancies among participants	27, range 1–5 per participant
Mode of conception (n=15 participants):	
Sexual intercourse	n=8, 53.3%
Sperm donation	n=5, 33.3%
Insemination with partner's sperm	n=1, 6.7%
Would rather not say	n=1, 6.7%
Mode of delivery (n=17 term deliveries)	
Vaginal delivery	n=13
Caesarean delivery	n=4
Pregnancy loss	n=8, 14 total (range 1–5 pregnancy losses)
Number of participants who gave birth during their playing career	4
Number of weeks to return to training	range 6–52, mean 19 weeks
Number of weeks return to competitive matches	range 6–52, mean 22 weeks

through sexual intercourse, 5 through sperm donation, 1 through insemination with partner's sperm and 1 participant selected the option 'would rather not say'. Eight participants reported having at least one pregnancy loss, with one participant having experienced five pregnancy losses. Four of the retired women's players gave birth during their playing career, and the mean time to return to competitive matches following delivery was 22 weeks. The findings are summarised in [table 4](#).

### Pelvic floor dysfunction

The mean PFDI-20 score for participants was 72 (lower scores indicate better pelvic floor function) (SD 35.2; range 8–151). There was no significant difference in PFDI-20 scores between participants who had delivered a baby through vaginal delivery (n=13) compared with those who had not (n=53) (Mann-Whitney test:  $U=327.00$ ,  $Z=-0.28$ ).

### DISCUSSION

This study describes the self-reported reproductive health of an elite cohort of women's footballers (79% of participants previously played senior international football) with specific reference to menstrual function, pregnancy and motherhood, contraceptive use and pelvic floor function. 33% of retired elite women's footballers had experienced amenorrhoea at some point during their lifetime, with

less than a quarter of the cohort being mothers. Of those who had become pregnant, 60% became pregnant within 1 year of trying. There did not appear to be any self-reported evidence of significant pelvic floor dysfunction in this population.

### Pregnancy and fertility considerations among retired women's footballers

With increasing numbers of women playing elite football<sup>33</sup> and potentially more competing after pregnancy, a better understanding of how to support these players is necessary. There is a growing interest in understanding pregnancy outcomes among active and retired elite female athletes,<sup>1</sup> with emerging research suggesting that appropriately modified elite sport participation during pregnancy is not associated with adverse events during pregnancy, labour or delivery.<sup>3 18</sup> With reference to elite football, outcomes of interest may include (but not be limited to): recognising barriers to pregnancy for elite footballers, understanding their experiences of pregnancy, assessing the extent to which the provision of care that is sensitive to their sporting ambitions, best practice guidelines for how they should continue to train/return to training post partum and information on time to return to full training and match play. The mean return to match play for players post partum in our study was 22 weeks. Although we did not assess risk of injury

following return, there is evidence to suggest risk may be elevated during this period (eg, due to rapid changes in training loads).<sup>3</sup> Evidence-based guidelines are required to safely support the return to sport of elite footballers in the postpartum period.<sup>34</sup>

Elite female athletes have described how there is a common societal and sporting narrative (including pressure from coaches) that suggests they must choose between being an athlete or becoming pregnant.<sup>35</sup> Peak performance and fertility windows may overlap, complicating decisions for athletes. A study of retired elite rowers and rugby players reported higher rates of amenorrhoea, older age at childbirth and a trend towards earlier symptoms or onset of menopause indicating a potentially narrower 'reproductive window' for female athletes.<sup>20</sup> In our study, 33% of participants reported amenorrhoea and three participants had reached menopause (range 30–54 years), although the reasons (eg, premature ovarian failure) for early menopause in two participants aged 30 and 39 are unclear. This experience of amenorrhoea is lower than previous data recorded among retired elite athletes (45%)<sup>20</sup> but higher than the general population.<sup>36</sup> Less than a quarter of retired elite players in our study had been pregnant, which is relatively low considering the demographic of the population. Another study in a retired, mostly subelite, population reported that nearly half the participants had become pregnant.<sup>15</sup> Understanding the reasons for low pregnancy rates among retired elite women's footballers compared with the subelite population, is worth considering further.

### Pelvic floor health

There has been a positive promotion of pelvic floor health among professional women's footballers.<sup>37</sup> These initiatives have the potential to improve outcomes for women's footballers and to increase awareness of important health issues. Contact sports, such as rugby, have shown high rates (43%) of stress urinary incontinence (SUI) in active elite players (n=396) and players who had given birth were most at risk (106/172 of active elite players with SUI had given birth).<sup>38</sup> The mean PFDI-20 score for participants was 72 (SD=35.27) in our study which is lower than reference groups used in validation studies indicating better pelvic floor function, although this should be interpreted with caution due to the limited validity of interpreting the PFDI-20 in the general population.<sup>39 40</sup> Although pelvic function was satisfactory among the retired elite women football players in this study, steps should be taken to identify and support those who may be most at risk (eg, during the perinatal period).<sup>41</sup> Using screening tools validated in athlete populations (eg, PFD-SENTINEL) may provide the most clinically relevant information, although this had not been developed at the time of the survey design.<sup>42</sup>

### Clinical implications

This study addresses the limited evidence with respect to the reproductive health of retired elite women's

footballers.<sup>8</sup> The data collected here and in the study by Ling *et al*<sup>15</sup> provide useful information on women players which can be used to provide gender-specific support postretirement, such as has occurred with retired men's players.<sup>43</sup> This may include, but not be limited to, providing support with respect to fertility, pregnancy and pelvic floor health. These initiatives should complement efforts and policies to support women's footballers becoming pregnant during their career, should they wish to do so.

The number of former players who had experienced menstrual irregularity (n=17, 24.6%) and/or amenorrhoea (n=23, 33.3%) mirrors findings from Ling *et al*, where a number (n=115/560, 21.7%) of retired football players reported their menstrual periods had stopped for  $\geq 3$  consecutive months.<sup>15</sup> These findings lend support to the importance of menstrual cycle monitoring in professional football to support women's health and performance (eg, to screen for early signs of REDs).<sup>44</sup>

### Future research directions

This study highlights the need to understand reproductive health outcomes among retired women's footballers in greater detail—to empower practitioners and governing bodies to better support women's footballers during and after their career. Research efforts should focus on evaluating how to effectively support women's footballers in pregnancy, through to postpartum return to play. Additionally, understanding the best practice approach with respect to menstrual cycle monitoring in professional football, and how to advise those with irregular cycles is likely to benefit players across women's football. These approaches may lead to, among other benefits, an improved understanding of the impact of REDs on professional footballers both during and after their careers (eg, injury, fertility). For future survey design, it may be worth including 'I don't know' as an option, and while menopause was not defined for the purposes of the study (as a level of understanding among participants was anticipated), future studies could consider adding a question that increases the accuracy of participants' responses; for example, 'Have you been 'diagnosed' with menopause by a medical doctor?'. In addition to assessing for the presence of menstrual pain and regularity, future studies should consider assessing for the presence of other menstrual cycle symptoms as this may prove important to understanding the reproductive health of retired women players.

### Strengths and limitations

This study describes self-reported reproductive health outcomes in an elite women football population and paves the way for future research in this area. The findings may not necessarily apply to current professional women's footballers considering the changing nature of the sport and the recent improvement in conditions. The major limitation of our study is the number of respondents that was relatively low, with only 69 respondents

completing the survey in a 20-month period. The self-reported nature of the survey and potential for selection and recall bias are also important considerations which may further limit the generalisability of our findings. The use of PFDI-20 in this study is limited by not establishing the presence of pelvic floor dysfunction prior to dissemination.

## CONCLUSION

This study describes the self-reported reproductive health of an elite cohort of women's footballers with specific reference to menstrual function, pregnancy and motherhood, contraceptive use and pelvic floor function. A number (n=23, 33.3%) of retired elite women's footballers reported experiencing amenorrhoea, with less than a quarter of the cohort being mothers. For those who had become pregnant, the majority became pregnant within 2 years of trying. There did not appear to be any evidence of significant pelvic floor dysfunction in this population. Future research efforts should focus on improving the understanding of how to effectively support women's footballers in pregnancy, through to postpartum return to play. Best practice guidelines on the use of menstrual cycle monitoring and pelvic health support would also likely improve standards of care for women's footballers. Finally, stakeholders should consider gender-specific postretirement care for women's footballers, as has been developed for men's footballers.

## Author affiliations

- <sup>1</sup>Amsterdam UMC location University of Amsterdam, Department of Orthopedic Surgery and Sports Medicine, Meibergdreef 9, Amsterdam, Netherlands
- <sup>2</sup>Amsterdam Collaboration for Health & Safety in Sports (ACHSS), IOC Research Center, Amsterdam, Netherlands
- <sup>3</sup>Football Players Worldwide (FIFPRO), Hoofddorp, Netherlands
- <sup>4</sup>Department of Sport and Exercise Sciences, Manchester Metropolitan University, Manchester, UK
- <sup>5</sup>Family Medicine, McMaster University Michael G DeGroot School of Medicine, Waterloo, Ontario, Canada
- <sup>6</sup>Medical and Scientific Department - Games Group, International Olympic Committee, Lausanne, Switzerland
- <sup>7</sup>FIFA Scientific Advisory Committee, Zurich, Switzerland
- <sup>8</sup>Public Health and Family Medicine, University of Western Ontario Schulich School of Medicine and Dentistry, London, Ontario, Canada
- <sup>9</sup>Medical Department, Fédération Internationale de Football Association (FIFA), Zurich, Switzerland
- <sup>10</sup>Amsterdam Movement Sciences, Ageing & Vitality, Musculoskeletal Health, Sports, Amsterdam, Netherlands
- <sup>11</sup>Academic Center for Evidence based Sports medicine (ACES), Amsterdam, Netherlands
- <sup>12</sup>Section Sports Medicine, University of Pretoria, Pretoria, South Africa

X Steve den Hollander @steve\_dh1, Kirsty Elliott-Sale @ElliottSale, Margo Lynn Mountjoy @margo.mountjoy, Jane S Thornton @JaneSThornton and Vincent Gouttebarga @VGouttebarga

**Contributors** SC and VG designed the study and were responsible for data collection. SC performed the data analysis with input from VG and SdH. SC drafted the initial manuscript, with KE-S, MLM, SdH, JST, GK, AM and VG contributing to subsequent iterations. SC is the guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not applicable.

**Ethics approval** Ethical approval for the study was provided by the Medical Ethics Review Committee of the Amsterdam University Medical Centers (W21\_135#21.150; Amsterdam, The Netherlands). The study was conducted in accordance with the principles set out in the Declaration of Helsinki (2013).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

- Sean Carmody <http://orcid.org/0000-0001-8683-5532>  
 Steve den Hollander <http://orcid.org/0000-0002-6064-038X>  
 Kirsty Elliott-Sale <http://orcid.org/0000-0003-1122-5099>  
 Margo Lynn Mountjoy <http://orcid.org/0000-0001-8604-2014>  
 Jane S Thornton <http://orcid.org/0000-0002-3519-7101>  
 Vincent Gouttebarga <http://orcid.org/0000-0002-0126-4177>

## REFERENCES

- 1 Bø K, Artal R, Barakat R, *et al.* Exercise and pregnancy in recreational and elite athletes: 2016 evidence summary from the IOC expert group meeting, lausanne. part 1-exercise in women planning pregnancy and those who are pregnant. *Br J Sports Med* 2016;50:571–89.
- 2 Bø K, Artal R, Barakat R, *et al.* Exercise and pregnancy in recreational and elite athletes: 2016/17 evidence summary from the IOC expert group meeting, lausanne. part 3-exercise in the postpartum period. *Br J Sports Med* 2017;51:1516–25.
- 3 Sundgot-Borgen J, Sundgot-Borgen C, Myklebust G, *et al.* Elite athletes get pregnant, have healthy babies and return to sport early postpartum. *BMJ Open Sport Exerc Med* 2019;5:e000652.
- 4 Culvin A, Bowes A. The incompatibility of motherhood and professional women's football in England. *Front Sports Act Living* 2021;3:1–13.
- 5 Forsyth JJ, Sams L, Blackett AD, *et al.* Menstrual cycle, hormonal contraception and pregnancy in women's football: perceptions of players, coaches and managers. *Sport Soc* 2023;26:1280–95.
- 6 Martin D, Timmins K, Cowie C, *et al.* Injury incidence across the menstrual cycle in international footballers. *Front Sports Act Living* 2021;3:616999.
- 7 Bruinvels G, Burden RJ, McGregor AJ, *et al.* Sport, exercise and the menstrual cycle: where is the research? *Br J Sports Med* 2017;51:487–8.
- 8 Carmody S, Anemaat K, Massey A, *et al.* Health conditions among retired professional footballers: a scoping review. *BMJ Open Sport Exerc Med* 2022;8:e001196.
- 9 Costello JT, Bieuzen F, Bleakley CM. Where are all the female participants in sports and exercise medicine research? *Eur J Sport Sci* 2014;14:847–51.
- 10 Kryger K, Wang A, Mehta R, *et al.* Research on women's football: a scoping review. *Sci Med Football* 2021.
- 11 Prien A, Boudabous S, Junge A, *et al.* Every second retired elite female football player has MRI evidence of knee osteoarthritis before age 50years: a cross-sectional study of clinical and MRI outcomes. *Knee Surg Sports Traumatol Arthrosc* 2020;28:353–62.
- 12 Prien A, Prinz B, Dvořák J, *et al.* Health problems in former elite female football players: prevalence and risk factors. *Scand J Med Sci Sports* 2017;27:1404–10.



- 13 Prinz B, Dvořák J, Junge A. Symptoms and risk factors of depression during and after the football career of elite female players. *BMJ Open Sport Exerc Med* 2016;2:e000124.
- 14 Prien A, Feddermann-Demont N, Verhagen E, et al. Neurocognitive performance and mental health of retired female football players compared to non-contact sport athletes. *BMJ Open Sport Exerc Med* 2020;6:e000952.
- 15 Ling DI, Hannafin JA, Prather H, et al. The women's soccer health study: from head to toe. *Sports Med* 2023;53:2001–10.
- 16 Mountjoy M, Ackerman KE, Bailey DM, et al. 2023 International Olympic Committee's (IOC) consensus statement on relative energy deficiency in sport (reds). *Br J Sports Med* 2023;57:1073–97.
- 17 Taim BC, Ó Catháin C, Renard M, et al. The prevalence of menstrual cycle disorders and menstrual cycle-related symptoms in female athletes: a systematic literature review. *Sports Med* 2023;53:1963–84.
- 18 Wowdzia JB, McHugh T-L, Thornton J, et al. Elite athletes and pregnancy outcomes: a systematic review and meta-analysis. *Med Sci Sports Exerc* 2021;53:534–42.
- 19 Bo K, Artal R, Barakat R, et al. Exercise and pregnancy in recreational and elite athletes: 2016/17 evidence summary from the IOC expert group meeting, lausanne. part 4-recommendations for future research. *Br J Sports Med* 2017;51:1724–6.
- 20 Thornton J, Rosen C, Davenport M, et al. Beyond the medals: a cross-sectional study exploring retired elite female athletes' health. *BMJ Open Sport Exerc Med* 2023;9:e001479.
- 21 Parker LJ, Elliott-Sale KJ, Hannon MP, et al. An audit of hormonal contraceptive use in women's super league soccer players; implications on symptomology. *Sci Med Footb* 2022;6:153–8.
- 22 Almeida MBA, Barra AA, Saltiel F, et al. Urinary incontinence and other pelvic floor dysfunctions in female athletes in Brazil: a cross-sectional study. *Scand J Med Sci Sports* 2016;26:1109–16.
- 23 Jácome C, Oliveira D, Marques A, et al. Prevalence and impact of urinary incontinence among female athletes. *Int J Gynaecol Obstet* 2011;114:60–3.
- 24 Carmody S, Jones C, Malhotra A, et al. Put out to pasture: what is our duty of care to the retiring professional footballer? promoting the concept of the "exit health examination" (EHE). *Br J Sports Med* 2019;53:788–9.
- 25 Elm E von, Altman DG, Egger M, et al. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335:18064739:806–8.
- 26 Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (cherries). *J Med Internet Res* 2004;6:e34.
- 27 WMA. Wma Declaration of Helsinki – ethical principles for medical research involving human subjects, 2013. Available: <https://www.wma.net/policies-post/wmadeclaration-of-helsinki-ethical-principles-for-medical-research-involving-humansubjects/>
- 28 McKay AKA, Stellingwerff T, Smith ES, et al. Defining training and performance caliber: a participant classification framework. *Int J Sports Physiol Perform* 2022;17:317–31.
- 29 Barber MD, Walters MD, Bump RC. Short forms of two condition-specific quality-of-life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ-7). *Am J Obstet Gynecol* 2005;193:103–13.
- 30 Goutteberge V, Kerkhoffs G, Schutter M, eds. *The Secret Balance of Champions: Health Challenges in Elite Sport*. London: Future Science Group, 2020.
- 31 Hopkins WG. *A scale of magnitude for effect statistics*. 2009.
- 32 Woodward M. *Epidemiology: Study Design and Data Analysis*. Boca Raton: CRC Press, 2014.
- 33 UEFA Women's football across the National associations 2016/17, 2017. Available: [https://www.uefa.com/MultimediaFiles/Download/OfficialDocument/uefaorg/Women%27sfootball/02/43/13/56/2431356\\_DOWNLOAD.pdf](https://www.uefa.com/MultimediaFiles/Download/OfficialDocument/uefaorg/Women%27sfootball/02/43/13/56/2431356_DOWNLOAD.pdf) [Accessed 8 Mar 2024].
- 34 Kimber ML, Meyer S, McHugh T-L, et al. Health outcomes after pregnancy in elite athletes: a systematic review and meta-analysis. *Med Sci Sports Exerc* 2021;53:1739–47.
- 35 Davenport MH, Nesdoly A, Ray L, et al. Pushing for change: a qualitative study of the experiences of elite athletes during pregnancy. *Br J Sports Med* 2022;56:452–7.
- 36 He Y, Zheng D, Shang W, et al. Prevalence of oligomenorrhea among women of childbearing age in china: a large community-based study. *Womens Health (Lond Engl)* 2020;16:174550652092861.
- 37 The FA. Clubs to get support from The Well HQ on female athlete health. Available: <https://womensleagues.thefa.com/bwsl-bwc-the-well-hq-projects-female-athlete-health/> [Accessed 8 Mar 2024].
- 38 McCarthy-Ryan M, Perkins J, Donnelly GM, et al. Stress urinary incontinence prevalence and risk factors in female rugby players: a common health problem across four nations. *BMJ Open Sport Exerc Med* 2024;10:e001832.
- 39 Utomo E, Blok BF, Steensma AB, et al. Validation of the pelvic floor distress inventory (PFDI-20) and pelvic floor impact questionnaire (PFIQ-7) in a dutch population. *Int Urogynecol J* 2014;25:531–44.
- 40 Gafni-Kane A, Goldberg RP, Sand PK, et al. Enhanced interpretability of the PFDI-20 with establishment of reference scores among women in the general population. *Neurourol Urodyn* 2012;31:1252–7.
- 41 Zuchelo LTS, Bezerra IMP, Da Silva ATM, et al. Questionnaires to evaluate pelvic floor dysfunction in the postpartum period: a systematic review. *Int J Womens Health* 2018;10:409–24.
- 42 Giagio S, Salvioli S, Innocenti T, et al. PFD-sentinel: development of a screening tool for pelvic floor dysfunction in female athletes through an international delphi consensus. *Br J Sports Med* 2023;57:899–905.
- 43 Goutteberge V, Goedhart E, Kerkhoffs G. Empowering the health of retired professional footballers: the systematic development of an after career consultation and its feasibility. *BMJ Open Sport Exerc Med* 2018;4:e000466.
- 44 McGawley K, Sargent D, Noordhof D, et al. Improving menstrual health literacy in sport. *J Sci Med Sport* 2023;26:351–7.



16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

# Surveys of Health monitoring of female retired professional footballers - version 33.61



Printed on 16-04-2024 17:39:49 by Sean Carmody

## Survey '1. Personal characteristics, football career and medical history'

### 1. Personal characteristics, football career and medical history - Electronic informed consent

Number	Question	Answers
1.1	I have been fully informed about this study. Any questions I have asked about the study have been answered to my satisfaction. I had enough time to decide about my participation to the study.	<input type="checkbox"/> Yes
1.2	I have agreed to participate voluntarily in the study. I understand that I have the right to withdraw from the study at any time without disclosing any specific reasons.	<input type="checkbox"/> Yes
1.3	I consent to the collection and confidential use of my data in order to answer the research questions of this project.	<input type="checkbox"/> Yes
1.4	I consent to my data being used and stored confidentially. I understand that my data will not be accessible to anyone not directly involved in this study.	<input type="checkbox"/> Yes
1.5	I agree to participate in this study.	<input type="checkbox"/> Yes

### 1. Personal characteristics, football career and medical history - Personal characteristics

Number	Question	Answers
2.1	How old are you?	<input type="text"/> years
2.2	How tall are you (cm)?	<input type="text"/> cm
2.3	How much do you weigh (kg)?	<input type="text"/> kg

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

2.4 What is the highest level of education you have completed?

No school completed

Nursery/Elementary school

High school

Vocational/technical school

College, university or equivalent

2.5 Are you currently in employment?

Yes

No

2.5.1 **If 'Are you currently in employment?' is equal to 'Yes' answer this question:**  
What is your profession?

2.6 How many years ago did you retire from professional football?

years

2.7 What was your reason for retirement?

Injury

Personal Choice

Financial reasons

Other

2.7.1 **If 'What was your reason for retirement?' is equal to 'Other' answer this question:**  
What was your reason for retirement?

2.8 For how many seasons were you a professional footballer?

years

2.9 What position did you primarily play?

Goalkeeper

Defender

Midfield

Forward

2.10 What was the highest level of football that you played at primarily?

Highest national level/league

Second highest national level/league

Other

2.11 Did you play international football?

Yes

No

2.11.1 **If 'Did you play international football?' is equal to 'Yes' answer this question:**  
How many years ago did you retire from international football?

 years

## 1. Personal characteristics, football career and medical

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

## history - Medical history

Number	Question	Answers
3.1	Since retiring from professional football, have you been admitted to hospital?	<input type="radio"/> Yes <input type="radio"/> No
3.1.1	<b>If 'Since retiring from professional football, have you been admitted to hospital?' is equal to 'Yes' answer this question:</b> Why?	<input type="text"/>
3.2	Have you been diagnosed with a cardiovascular disease (e.g., heart attack, heart failure, angina, stroke, abnormal heart rhythms)?	<input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> I don't know
3.2.1	<b>If 'Have you been diagnosed with a cardiovascular disease (e.g., heart attack, heart failure, angina, stroke, abnormal heart rhythms)?' is equal to 'Yes' answer this question:</b> Which cardiovascular disease(s)?	<input type="text"/>
3.3	Have you been diagnosed with a mental health disorder (e.g., depression, anxiety, insomnia)?	<input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> I don't know
3.3.1	<b>If 'Have you been diagnosed with a mental health disorder (e.g., depression, anxiety, insomnia)?' is equal to 'Yes' answer this question:</b> Which mental health disorder(s)?	<input type="text"/>
3.4	Have you been diagnosed with a neurological disease (e.g., epilepsy, Alzheimer disease, Parkinson's disease, dementia)?	<input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> I don't know
3.4.1	<b>If 'Have you been diagnosed with a neurological disease (e.g., epilepsy, Alzheimer disease, Parkinson's disease, dementia)?' is equal to 'Yes' answer this question:</b> Which neurological disease(s)?	<input type="text"/>
3.5	Do you smoke?	<input type="radio"/> Yes <input type="radio"/> No
3.5.1	<b>If 'Do you smoke?' is equal to 'Yes' answer this question:</b> How many cigarettes a day do you smoke?	<input type="text"/>



16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

3.6 Do you use pain medication (prescription-only)?

Daily

Weekly

Monthly

Yearly or less

Do not use

3.7 Do you use pain medication (over-the-counter)?

Daily

Weekly

Monthly

Yearly or less

Do not use

3.8 Do you use sleeping tablets (prescription)?

Daily

Weekly

Monthly

Yearly or less

Do not use

3.9 Do you use sleeping tablets (over-the-counter)?

Daily

Weekly

Monthly

Yearly or less

Do not use

3.10 Do you use any medication? (please specify drug name, dose and frequency)

## 1. Personal characteristics, football career and medical history - Female Health Part 1

Number	Question	Answers
4.1	At about what age did you begin to menstruate?	<input type="text"/> years
4.2	Have you reached menopause?	<input type="radio"/> Yes <input type="radio"/> No
4.2.1	<b><i>If 'Have you reached menopause?' is equal to 'No' answer this question:</i></b> What is the average duration of your menstrual cycle? (i.e., average number of days between the first days of your menstrual cycles)	<input type="text"/> days

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

4.2.2	<b>If 'Have you reached menopause?' is equal to 'No' answer this question:</b> How many days do you typically bleed for during your period?	<input type="text"/> days
4.2.3	<b>If 'Have you reached menopause?' is equal to 'No' answer this question:</b> How would you describe your menstrual cycle?	<input type="checkbox"/> Regular <input type="checkbox"/> Irregular <input type="checkbox"/> Very irregular
4.2.4	<b>If 'Have you reached menopause?' is equal to 'No' answer this question:</b> Do you generally experience pain during your menstrual cycle?	<input type="radio"/> No <input type="radio"/> Yes, before a bleeding period <input type="radio"/> Yes, during a bleeding period; <input type="radio"/> Yes, after a bleeding period
4.3	Since your first menstrual cycle, did your menstrual bleeding period stop for three or more consecutive months (for reasons other than hormonal contraception)?	<input type="radio"/> Yes <input type="radio"/> No
4.4	Are you currently using contraceptives?	<input type="checkbox"/> No <input type="checkbox"/> Yes, cap <input type="checkbox"/> Yes, pill <input type="checkbox"/> Yes, condom <input type="checkbox"/> Yes, contraceptive implant <input type="checkbox"/> Yes, contraceptive injection <input type="checkbox"/> Yes, contraceptive patch <input type="checkbox"/> Yes, diaphragm <input type="checkbox"/> Yes, female condom <input type="checkbox"/> Yes, other
4.5	Are you currently using hormone replacement?	<input type="radio"/> Yes <input type="radio"/> No
4.6	Are you a mother?	<input type="radio"/> Yes <input type="radio"/> No
4.6.1	<b>If 'Are you a mother?' is equal to 'Yes' answer this question:</b> How many children do you have?	<input type="text"/>
4.7	Have you been pregnant?	<input type="radio"/> Yes <input type="radio"/> No
4.7.1	<b>If 'Have you been pregnant?' is equal to 'No' answer this question:</b> Have you tried / are you trying to become pregnant?	<input type="radio"/> Yes <input type="radio"/> No

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

---

4.7.2 **If 'Have you been pregnant?' is equal to 'Yes' answer this question:**  Less than one year  
How long on average did it take you to become pregnant following the onset of desire for pregnancy?  1-2 years  More than 2 years

---

4.7.3 **If 'Have you been pregnant?' is equal to 'Yes' answer this question:**  0  
How many times have you been pregnant?  1  2  3  4  5  6  7  8  9  10 or more

---

4.7.4 **If 'Have you been pregnant?' is equal to 'Yes' answer this question:**  Intercourse  
How did you get pregnant?  Insemination with partner's sperm  Insemination with sperm donation  I would rather not say

---

4.7.5 **If 'Have you been pregnant?' is equal to 'Yes' answer this question:**  Vaginal delivery  
How did you give birth?  Caesarean delivery  Adoption  Currently pregnant

---

4.8 Have you ever had a miscarriage?  Yes  No

---

4.8.1 **If 'Have you ever had a miscarriage?' is equal to 'Yes' answer this question:**  0  
How many miscarriages have you had?  1  2  3  4  5  6  7  8  9  10 or more

---

4.7.6 **If 'Have you been pregnant?' is equal to 'Yes' answer this question:**  Yes  
Did you ever give birth (vaginal or Caesarean) during your playing career?  No

---



16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

4.7.6.1 ***If 'Did you ever give birth (vaginal or Caesarean) during your playing career?' is equal to 'Yes' answer this question:***

How long (in weeks) did it take before you returned to training (average in case of multiple deliveries)?

4.7.6.2 ***If 'Did you ever give birth (vaginal or Caesarean) during your playing career?' is equal to 'Yes' answer this question:***

How long (in weeks) did it take before you returned to competitive matches (average in case of multiple deliveries)?

## 1. Personal characteristics, football career and medical history - Female Health Part 2

Number	Question	Answers
5.1	I think that my stomach is too big.	<input type="radio"/> Always <input type="radio"/> Usually <input type="radio"/> Often <input type="radio"/> Sometimes <input type="radio"/> Rarely <input type="radio"/> Never
5.2	I think that my thighs are too large.	<input type="radio"/> Always <input type="radio"/> Usually <input type="radio"/> Often <input type="radio"/> Sometimes <input type="radio"/> Rarely <input type="radio"/> Never
5.3	I think that my stomach is just the right size.	<input type="radio"/> Always <input type="radio"/> Usually <input type="radio"/> Often <input type="radio"/> Sometimes <input type="radio"/> Rarely <input type="radio"/> Never
5.4	I feel satisfied with the shape of my body.	<input type="radio"/> Always <input type="radio"/> Usually <input type="radio"/> Often <input type="radio"/> Sometimes <input type="radio"/> Rarely <input type="radio"/> Never

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

---

5.5 I like the shape of my buttocks.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.6 I think that my hips are too big.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.7 I think that my thighs are just the right size.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.8 I think that my buttocks are too large.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.9 I think that my hips are just the right size.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.10 I eat sweets and carbohydrates without feeling nervous.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

---

5.11 I think about dieting.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.12 I feel extremely guilty after overeating.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.13 I am terrified of gaining weight.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.14 I exaggerate or magnify the importance of weight.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.15 I am preoccupied with the desire to be thinner.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---

5.16 If I gain a pound, I worry that I will keep gaining.

Always  
 Usually  
 Often  
 Sometimes  
 Rarely  
 Never

---



16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

---

5.17 Do you usually experience pressure in the lower abdomen?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.18 Do you usually experience heaviness or dullness in the pelvic area?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.19 Do you usually have a bulge or something falling out that you can see or feel in your vaginal area?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.20 Do you ever have to push on the vagina or around the rectum to have or complete a bowel movement?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.21 Do you usually experience a feeling of incomplete bladder emptying?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.22 Do you ever have to push up on a bulge in the vaginal area with your fingers to start or complete urination?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.23 Do you ever feel you need to strain too hard to have a bowel movement?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

---

5.24 Do you ever feel you have not completely emptied your bowels at the end of a bowel movement?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.25 Do you usually lose stool beyond your control if your stool is well formed?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.26 Do you usually lose stool beyond your control if your stool is loose?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.27 Do you usually lose gas from the rectum beyond your control?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.28 Do you usually have pain when you pass your stool?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.29 Do you experience a strong sense of urgency and have to rush to the bathroom to have a bowel movement?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

5.30 Does part of your bowel ever pass through the rectum and bulge outside during or after a bowel movement?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

---

16/04/2024, 16:41

Surveys of Health monitoring of female retired professional footballers

5.31 Do you usually experience frequent urination?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.32 Do you usually experience urine leakage associated with a feeling of urgency, that is, a strong sensation of needing to go to the bathroom?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.33 Do you usually experience urine leakage related to coughing, sneezing or laughing?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.34 Do you usually experience small amounts of urine leakage (that is, drops)?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.35 Do you usually experience difficulty emptying your bladder?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.36 Do you usually experience pain or discomfort in the lower abdomen or genital region?

Not present  
 Not at all  
 Somewhat  
 Moderately  
 Quite a bit

5.37 Have you ever had a stress fracture?

Yes  
 No

5.38 Please indicate when it occurred and what bone(s).

5.39 Have you even been told you have low bone density (e.g., osteopenia or osteoporosis)?

Yes  
 No