Serum cardiac markers are inversely associated with VO_2max of amateur athletes in response to endurance training adaptations

Gashaw Tesema,^{1,2} Mala George,³ Soumitra Mondal,⁴ D Mathivana⁴

To cite: Tesema G, George M, Mondal S, *et al.* Serum cardiac markers are inversely associated with VO_2max of amateur athletes in response to endurance training adaptations. *BMJ Open Sport & Exercise Medicine* 2019;5:e000537. doi:10.1136/bmjsem-2019-000537

Accepted 20 March 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹Department of Sport Science, College of Natural and Computational Science, Mizan-Tepi University, Mizan, Ethiopia

²Department of Sport Science, College of Natural and Computational Science, Mekelle University, Mekelle, Ethiopia

³Department of Biochemistry, College of Health Sciences, Mekelle University, Mekelle, Ethiopia

⁴Department of Sport Science, College of Natural and Computational Science, Mekelle University, Mekelle, Ethiopia

Correspondence to

Dr Gashaw Tesema, Department of Sport Science, College of Natural and Computational Science, Mizan-Tepi University, Mizan, Ethiopia; gashatesema@gmail.com

ABSTRACT

Background The influence of endurance training intensity and adaptation on serum cardiac markers is poorly understood and controversial; however, no enough data observed the association of serum cardiac markers with VO_2max . Therefore, we aimed to investigate whether serum cardiac markers are associated with maximum oxygen consumption (VO_2max) in response to 12-week endurance training on amateur athletes.

Methods 15 apparently healthy male amateur athletes with 19.47 ± 1.30 years of age were recruited and participated in endurance training with 70%–80% maximal heart rate intensity for 35 min per session for the first week and 2 min increments each week from the second to the last week for a period of 12 weeks. VO_2max and serum cardiac markers (lactate dehydrogenase [LDH], creatine kinase myocardial band (CK-MB) and cardiac troponin I [CTnI]) were assessed at the beginning of the training and after 12-week endurance training.

Results The result of CTnI indicated significantly ($p < 0.01$) and inversely ($r = -0.466$) correlated with VO_2max and CK-MB indicated significantly ($p < 0.01$) and inversely associated with VO_2max ($r = -0.536$) with moderate relationship. However, we did not find a significant association on LDH ($p > 0.05$) with VO_2max in response to endurance training adaptation.

Conclusion Our finding confirms our hypothesis that serum cardiac markers are inversely associated with estimated VO_2max in response to endurance training adaptation.

INTRODUCTION

Endurance training causes muscle damage, fatigue and muscle pain.^{1 2} This damage can be linked to large artery wall stiffening, cartilage and muscle damage,³ atrial fibrillation as well as left and right ventricular dysfunction⁴ and systemic inflammatory reaction⁵ resulting in elevated serum muscle damage markers. A study indicates seven continuous days of training did not increase incidences of muscle damage as compared with the group performed only one session of the training⁶ and this is attributed to adaptation of the stress caused by the training.⁶ Adaptations of

What are the new findings?

- ▶ Serum cardiac markers and maximum oxygen consumption have inverse relationship in young amateur moderately trained athletes.
- ▶ Gradual training intensity adaptation results in decreased serum cardiac markers.
- ▶ Increased serum cardiac markers are dependent on adaptation of training.

How might it impact on clinical practice in the near future?

- ▶ Elevation of serum cardiac markers in young, moderately trained amateur athletes may not be pathological rather it may be due to training experience, genetic factor and extreme intensity as well as training duration which needs further considerations.

training stress are best explained by maximum oxygen consumption (VO_2max) as the single indicator of cardiorespiratory fitness. In line with this, researchers⁷ suggested the time course of the training related to muscle fibre damage, pain, as well as creatine kinase (CK), cardiac troponin I (CTnI) and lactate dehydrogenase (LDH) release. However, there are no evidences that indicate the association of serum cardiac markers and VO_2max of amateur athletes in response to endurance training adaptation.

Moreover, studies reported serum cardiac markers (LDH, CK myocardial band [CK-MB] and CTnI) are not significantly correlated with VO_2max ,⁸ a measure of endurance training adaptation, which is a more accurate and reproducible predictor of cardiovascular outcomes.⁹ Conversely, a study reported muscle damage markers are inversely related to VO_2max .¹⁰ Besides, the prevalence of elevated CTnI is associated with inexperienced athletes than experienced athletes.¹¹ This difference might strongly related to previous

training experience¹² and age¹⁰ of an athlete. Moreover, a 12-week soccer training with 80% (maximal heart rate, HRmax) average intensity: reported increased VO₂max and inversely decreased CK-MB at the end of the 12-week training.¹³

However, there is not enough data observed the association of serum cardiac markers with VO₂max in response to 12-week endurance training adaptation of amateur athletes. Moreover, there is a lack of prospective studies that observe the relationship of high intensity endurance training to exercise-induced cardiac remodelling.¹⁴ Thus, since the athletes are exposed for elevated serum cardiac markers as a result of endurance training, it helps them to understand the level of serum markers in response to adaptation and preventing overtraining for monitoring fitness and enhancing cardiac health.

The aim of this study was to examine the association between serum cardiac markers and VO₂max in response to chronic endurance training on amateur athletes. Thus, we have applied high intensity endurance training at 70%–80% HRmax with duration of 35 min per session for the first week and 2 min increments each week from the second to the last week for a period of 12 weeks. Consequently, we hypothesised that serum cardiac markers are inversely associated with estimated VO₂max because of endurance training adaptation.

METHODS

Study area

This study was conducted in Bahir Dar, located about 578 km north-northwest of Addis Ababa-Ethiopia. It has an altitude of 1840 m above the sea level and within latitude and longitude of 11°36'N 37°23'E coordinates. While the annual average temperature is 25°C–32°C, with an average humidity of 58%.¹⁵

Study population

In all, 15 apparently healthy male amateur moderately trained athletes within 18–25 years of age were selected randomly from Bahir Dar University sport academy students. The sample size was based on similar studies done before¹⁶ using the following equations which considers 80% power and $p \leq 0.05$, where n =sample size needed, σ^2_T is variance of the treatment, σ^2_R is variance of the reference, δ is the maximum percent of no clinical importance, CVT is the coefficient of variations of the treatment and CVR is the coefficient of variations of the reference:

$$n = \frac{(z\alpha + z\beta/2)^2 (\sigma_T^2 + \sigma_R^2)}{(\delta - CVT - CVR)^2}$$

Consequently, a validated physical activity readiness questionnaire¹⁷ was used to evaluate conditions that may prohibit participants from practising in endurance training. Therefore, participants that reported problems in their health (heart dysfunction and chest pain during exercise) and physical conditions (loss of balance,

consciousness and bone or joint problem) and are currently taking anti-hypertensive drugs were excluded from the study. In addition, smokers and alcoholics were also excluded from participating in the study due to reported influences on both metabolic and physical performance biomarkers.^{18,19}

Study design

A single group pre–post design was used in this study. In all, 15 subjects selected and placed to high intensity endurance training. VO₂max and serum markers of muscle damage were assessed at the beginning of the training and after 12-week endurance training. High intensity endurance training was done at 70%–80% HRmax 35 min per session for the first week and 2 min increments each week from the second to the last week for a period of 12 weeks.

Training protocol and ethical approval

This field-based study conducted based on the protocol specified here. One professional athletics coach assigned to the experiment. Five minutes of warming up and stretching exercise are given at the beginning of the training. The participants were performed at 70%–80% HRmax, a continuous endurance running in the open field track for 35 min per session for the first week and 2 min increments each week from the second to the last week. The training intensity was monitored with polar heart rate monitor throughout the entire session. The heart rate monitor adjusted to the designed percent maximum heart rate. Therefore, the beep sound of the heart rate monitor guided us if it is below and above the designed % HRmax to adjust the heart rate drift throughout the training. Five minutes of cooling down training were also given at the end of each training time for 3 days per week and it continued for 12 weeks aimed to observe chronic effects of the training on serum cardiac markers and VO₂max. Participants were instructed not to participate in other exercise training and to continue the usual habit of nutrition throughout the experiment.

Ethical approval was obtained from Institutional Research Ethics Review Committee of Mekelle University conformed to the 1975 Declaration of Helsinki. Written consent was delivered to the participants and they were informed about the objectives of the study. Participation in this study was voluntary and their right not to participate was respected. Confidentiality and anonymity were also ensured.

Assessment of VO₂max

VO₂max of subjects was estimated using 3 min step test. Subjects stepped up and down a 16.5 inch stepping bench with a rate of 96 steps per min for 3 min in time of metronome beep sound. Five seconds after exactly 3 min pulse was taken from radial artery with heart rate monitor tied around the wrist. Finally, maximal aerobic power (VO₂max) was calculated using the equation VO₂max = 111.33 – (0.42 × maximum pulse measured 5

s after stepping for 3 min)²⁰ and reported in (mL/kg/min).

Assays of serum cardiac markers

Blood samples of 5 mL were collected just before training (pretest) from each participant. Then, post-training samples collected 4 hours after the final training session at the end of 12-week endurance training. Since the peak cardiometabolic markers have been achieved 3–4 hours after training time.^{21 22} Blood was taken from an antecubital vein using vacutainer serum separator tube containing blood clotting accelerant gel. The serum was separated by centrifugation of the blood sample at 4000 rpm (revolution per minute) for 3 min and stored at –20°C until analysis. The level of LDH in U/L was measured using a spectrophotometric assay with BS-2E chemistry analyzer according to the guidelines of the International Federation of Clinical Chemistry (Beckman Coulter, Krefeld, Germany). The levels of cTnI band in ng/mL and CK-MB in ng/mL were measured by chemiluminescence immunoassay using Maglumi 800 fully automated chemiluminescence immunoassay analyzer.

Data analysis

Analysis of the data obtained from the study inferential statistics was employed. To evaluate whether or not the data are normally distributed, the Kolmogorov-Smirnov and Shapiro-Wilk^{23 24} normality tests were applied. To identify the presence of any significant relationships between variables, Pearson's correlation analysis was used. IBM-SPSS V.20 packages were used to analyse the data. Variables were expressed as Pearson's correlation coefficient (r) and probability value (p). In all cases, $P < 0.05$ was accepted significant.

RESULTS

The Kolmogorov-Smirnov and Shapiro-Wilk's normality tests^{23 24} indicate ($p > 0.05$) and visual inspection of their histograms, normal Q–Q plots and box plots showed that the data were approximately normally distributed. General characteristics of participants in mean±SD were age in years (19.47±1.30), height in metres (1.67±0.07), body mass in kg (53.96±5.67) and body mass index in kg/m² (19.31±1.41).

Serum cardiac markers correlation results with estimated VO_2max in response to 12-week endurance training are shown in table 1. Value of CTnI with 12-week endurance training was significantly ($p < 0.01$) and inversely correlated with estimated VO_2max ($r = -0.466$) indicating low relationship. Moreover, CK-MB was statistically significant ($p < 0.01$) and inversely associated with estimated VO_2max ($r = -0.536$) with moderate relationship. Furthermore, CTnI and LDH showed significant ($p < 0.05$) and positive association with CK-MB. However, we did not find a significant association of LDH with VO_2max ($r = -0.23$) and CTnI in response to endurance training adaptation in amateur athletes.

Table 1 Serum cardiac muscle damage markers correlation result with VO_2max

Pearson correlations		VO_2max	CK-MB	CTnI	LDH
VO_2max	Pearson's r	–			
	P value	–			
CK-MB	Pearson's r	–0.536**	–		
	P value	0.002	–		
CTnI	Pearson's r	–0.466**	0.362*	–	
	P value	0.009	0.050	–	
LDH	Pearson's r	–0.236	0.368*	0.066	–
	P value	0.209	0.045	0.728	–

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

CK-MB, creatine kinase myocardial band; CTnI, Cardiac troponin I band; LDH, Lactate dehydrogenase; VO_2max , maximum oxygen consumption.

DISCUSSION

The purpose of this study was to examine the association between serum cardiac markers and VO_2max in response to chronic endurance training on amateur athletes so as to consider cardiac health. The main findings of the study support our hypothesis that serum cardiac markers are inversely associated with estimated VO_2max in response to endurance training adaptation. Consequently, serum cardiac markers (CTnI and CK-MB) in response to 12-week endurance training indicated a statistically significant ($p < 0.01$) association with estimated VO_2max . However, we did not find a significant association on LDH ($p > 0.05$) with estimated VO_2max in response to endurance training adaptation.

CTn and VO_2max are important factors in the study of cardiac muscle damage markers and cardiorespiratory fitness. Consequently, our study confirmed a statistically significant and inverse associations of CTnI with estimated VO_2max . Consistently, our study reported CTnI is inversely related to VO_2max .¹⁰ The reason might be adaptation of the stress caused by chronic endurance training.⁶ Conversely, a study⁸ reported CTnI has no relation with VO_2max . This disparity could be attributed to training experiences of the athletes.^{2 11}

Recent evidences reported increased VO_2max and inversely decreased CK-MB at the end of the 12-week training¹³ with 80% (HRmax) average intensity. It supports our finding of CK-MB is significantly and inversely associated with estimated VO_2max in response to endurance training adaptation. Similarly, a study reported an inverse relationship between cardiac markers and VO_2max .¹⁰ On the contrary, CK-MB is not significantly correlated with VO_2max .⁸ The variation of results between studies might be training experience of athletes¹² and age.¹⁰

Changes in serum cardiac muscle damage markers can be influenced by gender, age, different training intensity, training method differences and in this study, we used calculated or estimated VO_2max rather than measured VO_2max which was the limitation of our study

design. Thus, further study is needed regarding the fore mentioned points. Nevertheless, we can see training experience effect with gradual increment in training duration for 12 weeks improved VO_2max and reduced serum cardiac markers. Therefore, these factors should be considered in designing training to improve cardiorespiratory fitness and maintain cardiovascular health.

CONCLUSION

In conclusion, serum cardiac markers, CTnI band and CK-MB, are significantly and inversely associated with estimated VO_2max in amateur athletes in response to 12-week endurance training. These findings add to the growing body of evidence linking between cardiac damage markers to cardiorespiratory fitness. Indicating gradual adaptation reduces serum cardiac marker concentration after exercise.

Acknowledgements The authors thank Mekelle University and Mizan-Tepi University for their support.

Contributors I am the contributor of this work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

1. Lee YH, Lee YH, Kim CK. Biomarkers of muscle and cartilage damage and inflammation during a 200 Km run. *Eur J Appl Physiol* 2007;99:443–7.
2. Lieber RL, Shah SFJ. Cytoskeletal disruption after eccentric contraction-induced muscle injury. *Clin Orthop Relat Res* 2002;403:S90–9.
3. Kim HJ, Lee YH, Kim CK, et al. Changes in serum cartilage oligomeric matrix protein (COMP), plasma CPK and plasma hs-CRP in relation to running distance in a marathon (42.195 Km) and an ultra-marathon (200 Km) race. *Eur J Appl Physiol* 2009;105:765–70.
4. Mascia G, Perrotta L, Galanti GPL. Atrial fibrillation in athletes. *Int J Sport Med* 2013;34:379–84.
5. Wallberg L, Mikael Mattsson C, Enqvist JK, et al. Plasma IL-6 concentration during ultra-endurance exercise. *Eur J Appl Physiol* 2011;111:1081–8.
6. Chen TC, Hsieh SS. Effects of a 7-day eccentric training period on muscle damage and inflammation. *Med Sci Sports Exerc* 2001;33:1732–8.
7. Evans GF, Haller RG, Wyrick PS, et al. Submaximal delayed-onset muscle soreness: correlations between MR imaging findings and Clinical measures. *Radiology* 1998;208:815–20.
8. Ponte AD, Giovanelli N, Antonutto G. Changes in cardiac and muscle biomarkers following an uphill-only marathon. *Res Sport Med An Int J* 2017;26:1–12.
9. Myers J, Gullestad L, Vagelos R, et al. Hemodynamic, and cardiopulmonary exercise test determinants of survival in patients referred for evaluation of heart failure. *Ann Intern Med* 1998;129:286–93.
10. Manfredi TG, Fielding RA, O'Reilly KP, et al. Plasma creatine kinase activity and exercise-induced muscle damage in older men. *Med Sci Sports Exerc* 1991;23:1028–34.
11. Shave R, George KP, Atkinson G, et al. Exercise-induced cardiac troponin T release: a meta-analysis. *Med Sci Sport Exerc* 2007;39:2099–106.
12. Mehta R, Gaze D, Mohan S, et al. Post-exercise cardiac troponin release is related to exercise training history. *Int J Sports Med* 2012;33:333–7.
13. Bekris E, Gissis I, Sambanis M, et al. The effect of soccer training on biochemical indices of young soccer players during the game season. *Phys Train* 2012.
14. La Gerche A, Gerche L. The potential cardiotoxic effects of exercise. *Can J Cardiol* 2016;32:421–8.
15. Haile Z. Assessment of climate change impact on the net Basin supply of Lake Tana water balance. The Netherlands: International institute for geo- information science and earth observation ENSCHEDE, 2009.
16. Thalheimer W, Cook S. How to calculate effect sizes from published research articles: a simplified methodology. *Work Res* 2002.
17. Warburton DER, Jamnik VK, Bredin SSD. The physical activity readiness questionnaire for everyone (PAR-Q+) and electronic physical activity readiness medical examination (ePARmed-X+). *Heal Fit J Canada* 2011;4:3–23.
18. Lee C-L, Chang W-D. The effects of cigarette smoking on aerobic and anaerobic capacity and heart rate variability among female university students. *Int J Womens Health* 2013;5:667–79.
19. Barnes MJ. Alcohol: impact on sports performance and recovery in male athletes. *Sports Med* 2014;44:909–19.
20. Marquaria R, Aghema PRE. Determination of maximal oxygen consumption in man. *J Appl Physiol* 1965;20:1070–3.
21. Tian Y, Nie J, Huang C, et al. The kinetics of highly sensitive cardiac troponin T release after prolonged treadmill exercise in adolescent and adult athletes. *J Appl Physiol* 2012;113:418–25.
22. Legaz-Arrese A, López-Laval I, George K, et al. Impact of an endurance training program on exercise-induced cardiac biomarker release. *Am J Physiol Heart Circ Physiol* 2015;308:H913–20.
23. Razli M, Yap W. Power comparisons of some selected normality tests. in: proceedings of the regional conference on statistical sciences 2010:126–38.
24. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika* 1965;52:591–611.