



Concussion Symptom Reporting in Rugby League – Player Survey



Please mark the one most accurate answer for the following 9 questions:

1. **Years playing for an NRL club (U20s/NRL/Reserve Grade)**
 - 0-2
 - 3-5
 - >5
2. **What level are you currently playing at (most recent game)?**
 - NRL
 - Reserve Grade
 - U20s
3. **Age**
 - <18
 - 19 – 22
 - 23 – 25
 - 26 – 28
 - 28 – 30
 - >30
4. **Regular playing position**
 - Wing
 - Fullback
 - Halves
 - Hooker
 - Centre
 - 2nd Row
 - Front row
 - Lock
5. **Number of diagnosed concussions (by a doctor) in the past 2 seasons (game or training)**
 - None
 - 1-2
 - 3-5
 - 5+
6. **Have you felt you have suffered a concussion in the past two seasons and not reported to a member of the medical staff (game or training)**
 - No never
 - Yes, 1-2 occasions
 - Yes 3-5 occasions
 - Yes 5+ occasions
7. **Have you ever not reported a symptom to the doctor during a concussion assessment?**
 - I always report my symptoms
 - Yes, during pre-season baseline testing
 - Yes, during or after a game
 - Yes, during return to play testing
 - Yes, multiple different occasions
8. **For what reason have you not reported a symptom(s) to the doctor during the assessment?**
 - I always report my symptoms
 - I didn't want to be ruled out of the game/training
 - I didn't want to let down the team and/or coaches
 - I didn't think the symptom(s) were related to concussion
 - Despite the symptom(s), I didn't think they were significant
 - It occurred during an important game or time of the season
 - Fear that having a concussion diagnosis would affect future selections or contracts.
9. **Have you had annual education from your NRL club surrounding concussion symptoms and the potential consequences of under-reporting concussion?**
 - Yes
 - No
 - Don't know

Principle investigator: **Dr Thomas Longworth**, BMed, PG Dip SEM (Bath)