Is there a role for physical activity interventions in the treatment pathway of bladder cancer? A scoping review of the literature

Agustina Bessa ¹, Cecilia Bosco,² Sneha Mehrotra,¹ Megan Rowland,³ Hanyu Zhang,³ Beth Russell,³ Louis Fox ¹, Katharina Beyer,³ Elke Rammant,⁴ Suzanne Amery,⁵ Kathryn Chatterton,⁵ Nicola Peat,⁵ Christel Haggstrom,⁶ Mieke Van Hemelrijck¹

To cite: Bessa A, Bosco C, Mehrotra S, *et al*. Is there a role for physical activity interventions in the treatment pathway of bladder cancer? A scoping review of the literature. *BMJ Open Sport & Exercise Medicine* 2021;**7**:e000951. doi:10.1136/bmjsem-2020-000951

Accepted 18 February 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Translational Oncology Urology Research, King's College London, London, UK

²Faculty of Life Sciences and Medicine, King's College London, London, UK

³King's College London – Strand Campus, London, UK

⁴Ghent University, Ghent, Belgium

⁵Guy's and St Thomas' NHS Foundation Trust, London, UK

⁶Uppsala Universitet, Uppsala, Sweden

Correspondence to

Agustina Bessa;
agustina.m.bessa@gmail.com

ABSTRACT

Introduction Physical activity (PA) interventions have been introduced in patients with cancer as they may contribute to better treatment outcomes and quality of life (QoL). However, little is known about the impact of PA on patients with bladder cancer (BC). This scoping review aimed to explore efficacy and feasibility of existing PA interventions in the BC care pathway.

Methods and analysis Preferred Reporting Items for Systematic Reviews and Meta-Analyses Scoping Review guidelines and the Levac methodology framework were used; electronic databases were searched. Two independent reviewers screened all titles, abstracts and full-text publications for inclusion. The feasibility of integrating a PA intervention in the BC treatment pathway was discussed in a consultation phase with healthcare professionals and patient and public representatives.

Results A total of 675 records were identified through database searching of which 14 studies were included in our scoping review. An additional 17 clinical trials were identified of which 12 were included for which no results have been published yet. The included studies looked at the feasibility of a PA intervention programme, the associations between PA, obesity and BC, but also the determinants of PA engagement for BC patients and the assessment of QoL.

Conclusion This scoping review highlights that despite the general recognition on the role of PA in the BC treatment pathway, there is a gap regarding the understanding of the impact of PA interventions in BC care pathways as well as the limited understanding of factors underlying possible benefits of PA. No clear conclusions could be made regarding structure and processes of PA interventions that may lead to better outcomes. Further PA studies for patients with BC are needed to understand how to incorporate exercise guidelines recommendations.

BACKGROUND

Recognition of the impact of physical activity (PA) for people affected by cancer has expanded in recent years. Several studies

What are the new findings?

- ▶ Physical activity (PA) should be included in the bladder cancer (BC) preoperative and postoperative treatment pathway.
- ▶ The benefit of multimodal PA interventions in BC care pathways should be further investigated.
- ▶ Biological mechanisms and factors underlying possible benefits of PA in the BC care pathway have to be explored.

have shown the potential of PA at all stages of the cancer care pathway.^{1–3}

More specifically, PA interventions have been introduced in the cancer patient pathway to help improve treatment outcomes and increase health-related quality of life (HRQoL).⁴ The WHO defines HRQoL as a multidimensional concept that includes domains related to physical, mental, emotional and social functioning and focuses on the impact health status has on quality of life.⁵ A related concept of HRQoL is well-being, which is a broader concept that assesses the positive aspects of a person's life including life satisfaction.⁶

Several studies have demonstrated that PA should be part of the standard cancer care and treatment as it may improve disease-related physiological and psychological outcomes in cancer patients; this is mostly due to the reduction on drug toxicity levels and increases treatment completion rates.² For example, an analysis by Holmes *et al*⁷ in the Nurses' Health Study highlighted that engagement in sufficient vigorous PA following breast cancer diagnosis was inversely associated with breast cancer specific mortality risk. A recent Cochrane review looked at the association between the intensity of PA and the effects on HRQoL; it was concluded that the greater the

intensity, better HRQoL scores including factors such as social functioning and fatigue.⁸ The potential benefit of PA in treating patients with cancer has been linked to several mechanisms such as enhanced immune response, body composition, tumour vascularisation and tryptophan metabolism regulation.²

The potential benefits of PA on bladder cancer (BC) management has not been studied comprehensively as in other cancers; there is currently one review focusing specifically on those patients who have undergone a radical cystectomy, which is only affecting a subgroup of patients with BC.⁹ As highlighted in our recent research prioritisation work, although it is ninth most common cancer worldwide, with about 18 000 people being diagnosed annually in the UK alone,¹⁰ BC is an under-researched cancer.¹¹ The 2015 National Institute for Health and Care Excellence BC guidelines concluded: 'There is considerable variation in provision of BC care and evidence that the patient experience for people with BC is worse than that for people with other cancer'.^{12 13}

Scoping reviews are a useful tool to determine the coverage of the literature on a given topic and have an indication of the studies available as well as an overview of its focus. Scoping reviews help to clarify emerging evidence and pose more specific questions.¹⁴ Therefore, by conducting this scoping review, we aimed to explore efficacy and feasibility of existing PA interventions in the BC care pathway and to identify the gaps in the current work on the influence of PA on BC management.

METHODS

To explore the efficacy and feasibility of existing PA interventions in the BC care pathway and to identify the gaps in the current research on the influence of PA on BC management, a detailed protocol of this scoping review has been published elsewhere¹⁵ and a brief summary is presented in figure 1. Briefly, our methods were developed based on the Joanna Briggs Institute guidelines,¹⁶ and the methodological guidelines were developed by Levac *et al.*¹⁷ The Preferred Reporting Items for Systematic Reviews and Meta-Analyses Scoping Review extension for scoping reviews¹⁸ was followed to ensure that all suggested items are reported. A clinical research team was consulted to define the research questions. MEDLINE (using the PubMed interface) and Ovid Gateway (Embase and Ovid), the Cochrane Library, and PsycInfo and Global Health and Healthcare Management Information Consortium were searched for relevant studies between⁶ A separated search was conducted on ClinicalTrials.gov using the same search strategy. All types of study design were considered for inclusion if they assessed clinical/HRQoL and feasibility outcomes in patients who had a primary diagnosis of BC and PA had been measured (either as part of their lifestyle or as part of an intervention in the treatment pathway). Studies will be excluded if the publication is any other language than English or if the study was not relevant or had a wrong study design. Two other independent reviewers analysed

the studies included/excluded and discussed any conflict until a final consensus was obtained; however, no conflicts occurred, and the results of both reviewers were the same. The studies found were analysed using a qualitative and quantitative method. For qualitative aspects, the Donabedian conceptual framework¹⁹ was used to assess the possible efficacy of the PA interventions reported by analysing the structure, process and outcomes.

To ensure that our findings are relevant to those affected by BC and their healthcare professionals, the results were synthesised, presented and discussed in a consultation phase with a clinical oncologist, a medical oncologist, a BC nurse specialist, clinical research nurses, a psychologist as well as patients and patient and public involvement representative.

RESULTS

A total of 675 records were identified through database searching. After screening by title and abstracts, 54 were assessed for eligibility. In total, 14 studies matched the inclusion criteria and were included in this scoping review for qualitative synthesis (table 1). Figure 2 shows the flow chart of this scoping review. Four studies were

Figure 1 Stages of the scoping review. BC, bladder cancer; HRQoL, health-related quality of life; PA, physical activity.

Table 1 Descriptive table of the studies included in the scoping review

Ref	Author	Year of publication	Country	Aim/purpose	Study population	Study design	PA type (measurement)	Outcomes	Key findings
47	Kristina H Karvinen, Kerry S Courneya and Scott North	2007	Canada	To evaluate the relationship between exercise and QoL in BC survivors.	BC survivors (n=525).	Cohort/questionnaires sent to BC survivors.	A modified version of the Leisure Score Index from the Godin Leisure Time Exercise Questionnaire.	22.3% were meeting public health exercise recommendation; 16.0% were insufficiently active and 61.7% were completely sedentary.	QoL is positively influenced by exercise in BC survivors.
36	Chris M Blanchard and Kevin Stein	2010	Canada	To understand the associations among BMI, PA, and HRQoL in cancer survivors.	Cancer survivors (n=3306). Bladder cancer (n=210).	Cohort/questionnaires sent to cancer survivors.	Godin Leisure-Time Exercise Questionnaire.	Obese cancer survivors were significantly less likely to meet the PA recommendations. This indicates an association between BMI and PA and HRQoL.	Understand the relationship between BMI and PA will help identify target groups for PA interventions
48	Koebnick C, Michael D, Moore S C and Park Y	2008	USA	To clarify the association between BMI and PA.	Patients with BC (n=471 760).	Prospective cohort study.	Questionnaire.	No direct correlation was found between BC and PA.	The protective influence of PA on BC could occur through a mechanism involving reduction of body mass.
32	Michael A Liss, Martha White and Loki Natarajan	2017	USA	To understand the associations of BMI, PA and smoking with BC mortality.	Patients with BC (n=222 163).	Interview survey.	National Health Information Survey.	Exercise associated with reduced BC mortality.	BC mortality was lower in those patients who had the amount of exercise as per the guidelines as compared with no exercise.
38	Kristina H Karvinen and Kerry S Courneya	2007	Canada	To understand the exercise programmes and counselling preferences of patients with BC.	BC survivors (n=397).	Survey study.	Questionnaire.	81.1% of participants interested in an exercise programme designed for BC survivors.	Results indicate that the most common preferences were for exercise counselling in a presentational format.
33	Elke Rammant et al	2018	Belgium	To explore, for patients with BC, before and after cystectomy, the determinants of PA.	BC patients candidates for radical cystectomy (n=30).	Semistructured interviews.		PA motivational factors in patients with BC is multifactorial.	The development of theory-based behaviour change interventions to increase PA in patients with BC, should include multifactorial components.
49	Marilyn L Kwan, Lawrence H Kushi and Kim N Danforth	2019	USA	The goal is to investigate diet and lifestyle factors in recurrence and progression of NMIBC.	NMIBC (n=1282).	Prospective cohort study.	Questionnaire.	NA	NA
41	Jensen Laustsen, Jensen, Borre, Petersen	2016	Denmark	To investigate the feasibility, the adherence and the efficacy of a short-term physical prehabilitation programme.	Patients with BC awaiting radical cystectomy (n=107).	RCT	Standardised exercise programme consisting of both muscle strength exercises and endurance training.	A total of 66% adhered more than 75% of the recommended progressive standardised exercise programme.	In patients awaiting RC, a short-term exercise-based prehabilitation intervention is feasible and effective.

Continued

Table 1 Continued

Ref	Author	Year of publication	Country	Aim/purpose	Study population	Study design	PA type (measurement)	Outcomes	Key findings
42	Poserud A, Sherif A and Tollbäck A	2014	Sweden	Assessment of feasibility and effects of an exercise training programme.	Patients with BC after cystectomy with ileal conduit.	RCT	Exercise programme included group exercise training twice a week and daily walks.	The intervention group increased walking distance and the role physical domain in SF-36 more than the control group.	A 12-week group exercise training programme was not feasible for most cystectomy patients. Improvement in functional capacity and the role physical domain in HRQoL.
39	Minnella E M, Awasthi, Bousquet-Dion and Ferreira, Austin	2019	Canada	To investigate if a preoperative multimodal intervention is feasible and effective in radical cystectomy.	Patients with BC following radical cystectomy (n=70).	RCT	Preoperative conditioning intervention included aerobic and resistance exercise, diet therapy and relaxation techniques.	There is significant difference in functional capacity 4 weeks after surgery.	After radical cystectomy, multimodal prehabilitation resulted in faster functional recovery.
43	Connor Chestnuta, Woodson Smelser, Travis Duma and Misty Bechtela	2019	USA	To evaluate the feasibility of a diet and exercise programme and its impact in patients with DM2 and NMIBC.	NMIBC and pre-diabetes or type II diabetes mellitus.	Prospective clinical trial.	Participants were given diet and exercise counselling.	Average fasting blood glucose demonstrated a 9.65% relative reduction.	Incorporating an exercise regimen in patients with diabetes mellitus and NMIBC is feasible.
37	Bente Thoft Jensen, Jørgen Bjerggaard Jensen, Sussie Laustsen, annemette	2014	Denmark	To test if a standardised preoperative and postoperative physical exercise programme and enhanced mobilisation can impact on HRQoL and inpatient satisfaction in RC.	Patients with BC after radical cystectomy (n=107).	RCT	The intervention group received standard FT and an exercise-based intervention that involved both preoperative and postoperative exercises.	There was no difference in the global health score or in the scores of the five functional scales between treatment groups.	Preoperative and postoperative physical rehabilitation results on better HRQoL.

BC, bladder cancer; BMI, body mass index; CPET, cardiopulmonary exercise testing; Diabetes Mellitus type 2; DM2, Diabetes Mellitus type 2; NMIBC, non-muscle invasive bladder cancer; PA, physical activity; QoL, quality of life; RC, radical cystectomy; RCT, randomised controlled trial.

Figure 2 PRISMA flow chart diagram. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

conducted in Canada, two in Denmark, four in the USA, two in the UK, one in Sweden and one in Belgium. Of those, six were randomised controlled trials (RCTs), six were cohort questionnaires/surveys and two were interviews/semistructured interviews. From the ClinicalTrials.gov search, 17 studies were identified and assessed for eligibility and 14 studies coincide with the inclusion and exclusion criteria; of those, three had no information on location,^{20–22} one was in Sweden,²³ five in the USA,^{24–28} one in France,²⁹ one in Belgium and one in Canada.³⁰

Table 1 summarises the included studies. All RCTs had feasibility of a PA programme as their primary outcome; one also included cardiovascular outcomes.³¹ The interview studies evaluated the associations between PA, obesity and BC and the determinants of PA engagement in patients with BC.^{32–33} The questionnaires/surveys evaluated the associations of PA and QoL. The prospective cohort studies evaluated underlying biological factors such as body mass index (BMI) and glucose and one study also evaluated the cardiovascular component, in addition to the effect of PA.^{34–36}

Overall, the studies in table 1 reported that PA helped to provide improved QoL, but in one study, a physical rehabilitation programme, it was found to not impact on HRQoL.³⁷ However, the same study showed that preoperative and postoperative physical rehabilitation

provide better results on HRQoL. In addition, another study showed that BC survivors are more interested in an exercise programme specially designed for them in a physical presence format.³⁸ One study supported that multimodal prehabilitation resulted in faster functional recovery after radical cystectomy.³⁹ Regarding oncological outcomes, BC mortality was noted lowest in those patients who obtained the recommended amount of exercise as compared with no exercise. Patients with BC responded well to aerobic exercise, and it is suggested that improvements in cardiopulmonary/cardiorespiratory fitness variables could have important implications for postoperative recuperation after RC.³¹

Table 2 shows the data extraction for studies that looked at PA and suggested hypotheses for potential underlying biological mechanisms. Both studies looked at preoperative cardio exercise^{31–34}; patients with BC responded well to the aerobic exercise training, including the elderly population; patients were compliant with a supervised exercise regime, and it was concluded that the improvements in cardiopulmonary fitness variables could have important implications for postoperative recuperation after RC.³¹

Table 3 summarises the qualitative Donabedian conceptual framework of the included studies where information on process and structure of a PA intervention

Table 2 Studies that evaluate PA and proposed potential underlying biological mechanisms

Ref	Author	Year of publication	Country	Aim/purpose	Study population	Study design	PA type+duration	Outcomes	Proposed biological mechanism
31	Banerjee S, Manley K, Shaw B, Lewis L, Cucato G, Mills R, Rochester M, Clark A and Saxton J M	2018	UK	To evaluate the feasibility of vigorous intensity aerobic interval exercise in patients with BC prior to radical cystectomy.	Patients with BC patients prior to radical cystectomy (n=60).	RCT	Twice-weekly preoperative supervised vigorous intensity aerobic interval exercise in addition to standard treatment.	Improvements in oxygen pulse and power output were observed at the follow-up CPET in the exercise group vs controls.	Presurgical aerobic interval exercise and the improvements in cardiopulmonary fitness variables could have important implications for postoperative recuperation after RC.
34	Banerjee, S, Manley, K, Thomas, L, Shaw B, Saxton J, Mills R and Rochester M	2013	UK	Test the effect of a short preoperative course of supervised exercise in cardiopulmonary fitness.	Patients with BC who underwent radical cystectomy (n=30).	RCT	Patients in the intervention arm were offered twice weekly supervised exercise sessions preoperatively for 4 weeks, while the control arm had standard care.	In the control group, $\dot{V}O_2$ max was 22.8 (12.9–41.0) mL/kg/min, and the mean change was 0.48 mL/kg/min (95% CI -0.71 to 1.67) following the intervention. In the intervention arm, initial average $\dot{V}O_2$ max was 20.38 mL/kg/min, which increased to an average of 25.5 mL/kg/min.	Patients comply with a supervised exercise regime, and there seems to be an improvement in their cardiorespiratory fitness.

BC, bladder cancer; CPET, cardiopulmonary exercise testing; PA, physical activity; RC, radical cystectomy; RCT, randomised clinical trial; $\dot{V}O_2$, maximum rate of O_2 consumption.

Table 3 Donabedian conceptual framework of the included studies

Ref	Author	Aim/purpose	Structure	Process	Outcome
45	Jensen Laustsen <i>et al</i>	To evaluate the feasibility, the adherence and the efficacy of a short-term physical prehabilitation programme.	2-week preoperative exercise-based intervention were instructed by highly specialised physiotherapists dedicated to educate radical cystectomy patients.	It was recommended to repeat each exercise 10–15 times twice a day. Daily achievements were patient-reported in a personal diary. Patients were contacted by ways of telephone call 1 week before surgery in order to answer questions regarding any concerns related to the programme or documentation of achievements.	A total of 66% adhered to the recommended progressive standardised exercise programme.
42	Porserud A, Sherif A and Tollbäck A	Test the feasibility and benefits of an exercise programme.	Exercise programme included group exercise training twice a week and daily walks.	The intervention group took part in a group exercise training programme. The training programme started within a week after the baseline assessment and its main aim was to increase physical function. Music was used as inspiration. The daily walks were reported by the patients in an exercise training diary.	A 12-week group exercise training programme was not feasible for most cystectomy patients.
39	Minnella <i>et al</i> .	To evaluate the feasibility of a preoperative multimodal intervention in radical cystectomy patients	Preoperative conditioning intervention included aerobic and resistance exercise, diet therapy, and relaxation techniques.	This study was conducted at an academic tertiary healthcare institution and enrolled adult patients scheduled for radical cystectomy.	Multimodal prehabilitation resulted in faster functional recovery after radical cystectomy.
37	Bente Thoft Jensen Jørgen Bjerggaard Jensen, Sussie laustsen, annemette	To test if a standardised preoperative and postoperative physical exercise programme can impact on HRQoL and inpatient satisfaction.	The intervention group received multimodal rehabilitation and an exercise-based intervention that involved both preoperative and postoperative exercises.	Patients were randomised to fast-track RC and intervention (n=50) or fast-track standard treatment (n=57). HRQoL and inpatient satisfaction was measured using valid questionnaires.	There was no difference in the global health score.
31	Banerjee S, Manley K, Shaw B, Lewis L, Cucato G, Mills R, Rochester M, Clark A and Saxton J M.	To test vigorous intensity aerobic interval exercise in BC patients prior to radical cystectomy.	Twice-weekly preoperative supervised vigorous intensity aerobic interval exercise in addition to standard treatment.	The exercise group was offered twice-weekly preoperative supervised vigorous intensity aerobic interval exercise in addition to standard treatment. The controls received standard treatment only. A repeat CPET was undertaken before surgery and postoperative recovery outcomes were recorded.	Improvements in peak values of oxygen pulse and power output were observed at the follow-up CPET in the exercise group versus controls.
34	Banerjee, S, Manley K, Thomas L, Shaw B, Saxton J, Mills R and Rochester M.	To evaluate if short preoperative course of supervised exercise sessions is tolerated and whether it can lead to an improvement in cardiopulmonary fitness.		Patients in the intervention arm were offered twice weekly supervised exercise sessions preoperatively for 4 weeks, while the control arm had standard care.	In the control group, $\dot{V}O_2$ max was 22.8 (12.9–41.0) mL/kg/min, and the mean change was 0.48 mL/kg/min (–0.71 to 1.67) following the intervention. In the intervention arm initial average $\dot{V}O_2$ max was 20.38 mL/kg/min, which increased to an average of 25.5 mL/kg/min.

CPET, cardiopulmonary exercise testing; HRQoL, health-related quality of life; $\dot{V}O_2$, maximum rate of O_2 consumption.

was provided. Positive outcomes were shown in the studies that included a multimodal approach and aerobic exercise,³⁹ but no conclusions could be made regarding what structure and processes lead to the better outcomes for patients with BC.

Table 4 summarises the clinical trials currently listed on ClinicaTrials.gov. Four have been completed, but no results have yet been published.^{22 25 26 40} Two still need to start recruiting,^{41 42} four are currently recruiting^{20 21 29 43} and two have an unknown status^{23 29} (table 4).

To guarantee that our assessment (tables 1–4) is relevant to those affected by BC and their healthcare professionals, we also organised a consultation with patients, their family members and healthcare professionals. We held a face-to-face meeting in January 2020 with a clinical oncologist, a medical oncologist, urological surgeon, three nurse specialists, a physiotherapist, a patient and public involvement specialist and three patients (one male and two female) to discuss the feasibility of a PA component in an intervention for patients with BC. The findings of the scoping review were discussed, and it was concluded that PA should be included as part of a multimodal approach to support patients' well-being by combining PA support with other components of well-being such as mental or sexual well-being. A well-being intervention including PA, sexual and mental well-being support is currently in development.

DISCUSSION

Of the 14 studies included in our scoping review, six were RCTs, six were cohort questionnaires/surveys and two were interviews/semistructured interviews. All RCTs had feasibility of a PA programme as their primary outcome, and the interview studies evaluated the associations between PA, obesity and BC as well as the determinants of PA engagement in patients with BC. The questionnaires/surveys evaluated the associations of PA and quality of life. The prospective cohort studies evaluated underlying biological factors such as BMI and glucose in addition to PA. Nevertheless, no conclusions could be made regarding structure and processes of a PA intervention that could lead to better outcomes for BC patients. Moreover, although some clinical trials have been identified through ClinicalTrials.gov, no results have yet been published.

There is a increasing number of studies suggesting a potential benefits of PA at all stages of the cancer care pathway.¹ Looking at patients with BC in particular, our findings support this recommendation with studies showing the important implications of a preoperative and postoperative physical rehabilitation with most studies on aerobic/cardiovascular exercises. More studies are needed to evaluate the benefits of other types of exercises in relation to different BC treatments and time points. As PA may be part of BC treatment pathway, PA integrating exercises to strengthen pelvic floor and improve continence should be considered.

The benefits of prehabilitation activities for patients with cancer in general are well documented.^{35 44} In the context of BC, preoperative interventions have been found to be feasible and acceptable. However, few studies evaluated the underlying biological mechanisms of PA effects in the context of radical cystectomy and survival. The studies that aimed to clarify the underlined biological mechanisms focused on increasing the aerobic/cardiovascular/cardiopulmonary capacity of those patients.^{16 17} However, they mainly focused on the short or intense implications of PA, and it would be interesting to evaluate the effect of long-term PA in terms of biological mechanisms and its implications in BC response to treatment and survival.

It is known that after a cancer diagnosis and during medical treatments, it is difficult to become physically active.⁴⁵ Rammant *et al*⁹ explored the determinants of PA in patients with BC before and after radical cystectomy and, as per other cancer types, concluded that the PA behaviour in patients with BC is multifactorial. Minnella *et al*³⁹, in their RCT, also concluded that a multimodal prehabilitation approach resulted in faster functional recovery after radical cystectomy. These observations also support the points highlighted during our consensus meeting regarding the need for a multidisciplinary approach that also incorporates mental and sexual well-being to support patients with BC. PA activity is positively associated with HRQoL in BC survivors, and therefore, it would be interesting to further evaluate the underlying reasons that support that association. As explained above, HRQoL is a multidimensional⁵ concept that includes domains related to physical and mental well-being; it focuses on the impact health status has on quality of life.

From the literature review and the consultation phase, we can conclude that there are many questions to be answered concerning PA and BC. For example, which patients with BC would benefit the most from engagement in PA in terms of survival and quality of life?⁴⁶ Moreover, there is a need to understand which types of PA are needed and when they should be included during the BC treatment pathway. Given the high level of physical inactivity in the population, especially in a more elderly population such as patients with BC, it should be considered priority to understand how to implement PA in this group of patients.⁶ Our consultation phase with patients and HCPs confirmed the need to include a multimodal approach developing PA interventions.

Our extensive search through different databases, allowing for the inclusion of all types of interventions focused on PA, quality of life and biological mechanisms, is a strength of this scoping review. Another strength is the inclusion of a consultation phase that allowed to discuss the findings with patients and healthcare professionals. However, due to the nature of the methods used of the included studies and the lack of information in these published study designs, we were unable to provide a conclusion regarding the structure and processes of PA interventions that lead to better clinical outcomes

Table 4 Clinical trials on PA

Ref	Study results	Conditions	Interventions	Location	Year/status
20	No results available.	Bladder cancer. Urothelial carcinoma.	Behavioural: exercise programme.		2020. Not recruiting.
21	No results available.	Postoperative complications. Bladder cancer.	Behavioural: prehabilitation.		2019. Not recruiting.
23	No results available.	Urinary bladder cancer.	Other: The CanMoRe programme. Other: home exercise.	Karolinska University Hospital, Stockholm, Sweden.	2019. Recruiting.
24	No results available.	Bladder cancer.	Behavioural: activities of daily living. Behavioural: instrumental activities of daily living. Other: hand grip strength test. Other: timed up and go walking test. Behavioural: SF-8. Behavioural: FACT-VCI.	University of Miami, Miami, Florida, USA.	2018. Recruiting.
29	No results available.	Sarcopenia. Bladder cancer.	Diagnostic Test: sarcopenia and mobility measurement.	Caen University Hospital, Caen, France.	2017. Recruiting.
22	No results available.	Bladder cancer. Nutritional aspects of cancer.	Combination Product: Trimodal Prehab and ERP.		2017. Unknown.
40	No results available.	Bladder cancer. Urologic cancer surgery.		Ghent University, Ghent, Belgium.	2017. Unknown.
25	No results available.	Bladder cancer. Telomere shortening. Colorectal adenoma.	Behavioural: dietary intervention. Behavioural: PA intervention.	UPMC Shadyside, Pittsburgh, Pennsylvania, USA.	2017. Recruiting.
26	No results available.	Bladder cancer. Diabetes mellitus.	Other: CR diet. Behavioural: exercise.	University of Kansas Cancer Centre, Kansas City, Kansas, USA. University of Kansas Medical Centre, Kansas City, Kansas, USA.	2016. Published, no results regarding PA – completed.
27	No results available.	Bladder cancer.	Other: baseline strength test. Other: exercise.	University of Michigan, Ann Arbor, Michigan, USA.	2013. No results published – completed.
30	No results available.	Bladder cancer. Entire ileal conduit. Urinary bladder cancer	Other: prehabilitation.	Montreal General Hospital, Montreal, Quebec, Canada.	2013. No results published – completed.

Continued

Table 4 Continued

Ref	Study results	Conditions	Interventions	Location	Year/status
28	No results available.	Bladder cancer. Breast cancer. Colorectal cancer. Oesophageal cancer. Fatigue lung cancer. Lymphoma. Ovarian cancer. Prostate cancer.	Behavioural: exercise intervention Procedure: complementary or complementary medicine procedure Procedure: fatigue assessment and management Procedure: physical therapy.	Huntsman Cancer Institute at University of Utah, Salt Lake City, Utah, USA.	Includes all cancer but not bladder specific, results published – completed.

CR, Calorie restriction; FACT-VCI, Functional Assessment of Cancer Therapy-Vanderbilt Cystectomy Index; PA, physical activity.

for patients with BC. Future studies should better report on the development of their intervention, the different components and the processes followed. Additionally, the studies that analyse the possible underlying biological mechanisms to support the benefits of PA are limited, restricted to short period of time before BC treatment and no conclusion could be made on the long-term benefits of PA. Moreover, it is important to highlight that we did not identify any studies conducted in low-income or middle-income countries or non-western countries. Also, no multinational studies have been conducted.

CONCLUSION

This scoping review highlights the gap regarding the understanding of the impact of PA interventions in the BC pathway. It also highlights the limited understanding of the factors that are underlying the possible PA benefits, such as psychological aspects and biological components. Further PA studies for patients with BC are needed to understand where and how to incorporate the global WHO recommendations on PA.

Twitter Louis Fox @_LouisFox

Acknowledgements The authors gratefully acknowledge the helpful comments from the bladder cancer patient support group.

Contributors Conception and design: AB, CB and MVH. Acquisition of data: AB, CB, SM, MR and HZ. Analysis and interpretation of data: AB, CB and MVH. Drafting of the manuscript: AB and CB. Critical revision of the manuscript for important intellectual content: BR, LF, KB, ER, SA, KC, NP, CH and MVH.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Agustina Bessa <http://orcid.org/0000-0001-7196-2280>

Louis Fox <http://orcid.org/0000-0002-8280-1797>

REFERENCES

- Physical activity and cancer - A concise evidence review 2017.
- Christensen JF, Simonsen C, Hojman P. Exercise training in cancer control and treatment. *Compr Physiol* 2018;9:165–205.
- Ashcraft KA, Warner AB, Jones LW, et al. Exercise as adjunct therapy in cancer. *Semin Radiat Oncol* 2019;29:16–24.
- Albrecht TA, Taylor AG. Physical activity in patients with advanced-stage cancer: a systematic review of the literature. *Clin J Oncol Nurs* 2012;16:293–300.
- Who | WHOQOL: measuring quality of life. who 2014.
- Health-Related quality of life and well-being | healthy people, 2020. Available: <https://www.healthypeople.gov/2020/about/foundation-health-measures/Health-Related-Quality-of-Life-and-Well-Being> [Accessed 1 Oct 2019].

- 7 Holmes MD, Chen WY, Feskanich D, *et al.* Physical activity and survival after breast cancer diagnosis. *JAMA* 2005;293:2479–86.
- 8 Mishra SI, Scherer RW, Geigle PM, *et al.* Exercise interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev* 2012;CD007566.
- 9 Rammant E, Decaestecker K, Bultjnick R, *et al.* A systematic review of exercise and psychosocial rehabilitation interventions to improve health-related outcomes in patients with bladder cancer undergoing radical cystectomy. *Clin Rehabil* 2018;32:594–606.
- 10 Kockelbergh R, Hounsom L, Mayer E. The epidemiology of urological cancer 2001–2013. *J Clin Urol* 2017;10:3–8.
- 11 Bessa A, MacLennan S, Enting D, *et al.* Consensus in bladder cancer research priorities between patients and healthcare professionals using a four-stage modified Delphi method. *Eur Urol* 2019;0.
- 12 Bladder cancer: diagnosis and bladder cancer: diagnosis and management management NICE guideline 2015.
- 13 Edmondson AJ, Birtwistle JC, Catto JWF, *et al.* The patients' experience of a bladder cancer diagnosis: a systematic review of the qualitative evidence. *J Cancer Surviv* 2017;11:453–61.
- 14 Munn Z, Peters MDJ, Stern C, *et al.* Systematic review or scoping review? guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018;18:143.
- 15 Mehrotra S, Rowland M, Zhang H, *et al.* Scoping review protocol: is there a role for physical activity interventions in the treatment pathway of bladder cancer? *BMJ Open* 2019;9:e033518.
- 16 et al Peters MDJ, Godfrey C, McInerney P. Chapter 11: Scoping reviews - JBI Reviewer's Manual - JBI GLOBAL WIKI n.d. Available: <https://wiki.joannabriggs.org/display/MANUAL/Chapter+11%3A+Scoping+reviews> [Accessed 21 Jan 2020].
- 17 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69.
- 18 Tricco AC, Lillie E, Zarin W, *et al.* PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- 19 Gardner G, Gardner A, O'Connell J. Using the Donabedian framework to examine the quality and safety of nursing service innovation. *J Clin Nurs* 2014;23:145–55.
- 20 Exercise-based Pre-habilitation in Bladder Cancer Patients Prior to Radical Cystectomy: a Feasibility Study - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT04223063> [Accessed 30 Apr 2020].
- 21 Against All Odds -Prehabilitation in Urologic Cancer Surgery - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT04088968> [Accessed 30 Apr 2020].
- 22 Trimodal Prehabilitation for Cystectomy Patients to Enhance Post-operative Care - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT03347045> [Accessed 1 May 2020].
- 23 Physical Rehabilitation Among Patients Undergoing Radical Cystectomy Due to Urinary Bladder Cancer - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT03998579> [Accessed 30 Apr 2020].
- 24 NCT03469362. Intra-corporal vs Extra-corporal urinary diversion after robot assisted radical cystectomy, 2018. Available: <https://ClinicalTrialsGov/Show/Nct03469362>
- 25 Lifestyle Change for Better Health - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT03286699> [Accessed 1 May 2020].
- 26 The Feasibility of a Diet and Exercise Intervention in Diabetics During Treatment for Non-muscle Invasive Bladder Cancer - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT02716623> [Accessed 1 May 2020].
- 27 Trial of Prehabilitation in Vulnerable Patients Undergoing Cystectomy for Bladder Cancer - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT01840137> [Accessed 1 May 2020].
- 28 LaStayo PC, Marcus RL, Dibble LE, *et al.* Eccentric exercise versus usual-care with older cancer survivors: the impact on muscle and mobility--an exploratory pilot study. *BMC Geriatr* 2011;11:5.
- 29 Sarcopenia, mobility, physical activity and post-operative risk of bladder carcinoma in the elderly (SAMPHYR) 2017.
- 30 Li C, Carli F, Lee L, *et al.* Impact of a trimodal prehabilitation program on functional recovery after colorectal cancer surgery: a pilot study. *Surg Endosc* 2013;27:1072–82.
- 31 Banerjee S, Manley K, Shaw B, *et al.* Vigorous intensity aerobic interval exercise in bladder cancer patients prior to radical cystectomy: a feasibility randomised controlled trial. *Support Care Cancer* 2018;26:1515–23.
- 32 Liss MA, White M, Natarajan L, *et al.* Exercise decreases and smoking increases bladder cancer mortality. *Clin Genitourin Cancer* 2017;15:391–5.
- 33 Rammant E, Fonteyne V, Decaestecker K, *et al.* Understanding physical activity behavior in patients with bladder cancer before and after radical cystectomy: a qualitative interview study. *Clin Rehabil* 2019;33:750–61.
- 34 Banerjee S, Manley K, Thomas L, *et al.* O2 preoperative exercise protocol to aid recovery of radical cystectomy: results of a feasibility study. *Eur Urol Suppl* 2013;12:125.
- 35 Pouwels S, Stokmans RA, Willigendael EM, *et al.* Preoperative exercise therapy for elective major abdominal surgery: a systematic review. *Int J Surg* 2014;12:134–40.
- 36 Blanchard CM, Stein K, Courneya KS. Body mass index, physical activity, and health-related quality of life in cancer survivors. *Med Sci Sports Exerc* 2010;42:665–71.
- 37 Jensen BT, Jensen JB, Laustsen S, *et al.* Multidisciplinary rehabilitation can impact on health-related quality of life outcome in radical cystectomy: secondary reported outcome of a randomized controlled trial. *J Multidiscip Healthc* 2014;7:301–11.
- 38 Karvinen KH, Courneya KS, Venner P, *et al.* Exercise programming and counseling preferences in bladder cancer survivors: a population-based study. *J Cancer Surviv* 2007;1:27–34.
- 39 Minnella EM, Awasthi R, Bousquet-Dion G, *et al.* Multimodal prehabilitation to enhance functional capacity following radical cystectomy: a randomized controlled trial. *Eur Urol Focus* 2019. doi:10.1016/j.euf.2019.05.016. [Epub ahead of print: 08 Jun 2019].
- 40 Unmet Supportive Care Needs in Bladder Cancer Patients Undergoing Radical Cystectomy - Full Text View - ClinicalTrials.gov n.d. Available: <https://clinicaltrials.gov/ct2/show/NCT03307200> [Accessed 1 May 2020].
- 41 Jensen BT, Laustsen S, Jensen JB, *et al.* Exercise-based prehabilitation is feasible and effective in radical cystectomy pathways-secondary results from a randomized controlled trial. *Support Care Cancer* 2016;24:3325–31.
- 42 Porsrud A, Sherif A, Tollbäck A. The effects of a physical exercise programme after radical cystectomy for urinary bladder cancer. A pilot randomized controlled trial. *Clin Rehabil* 2014;28:451–9.
- 43 Chestnut C, Smelser W, Dum T, *et al.* Glycemic impact of a diet and lifestyle intervention on diabetics and prediabetics during treatment for non-muscle invasive bladder cancer. *Nutr Cancer* 2020;72:1219–24.
- 44 Tsimopoulou I, Pasquali S, Howard R, *et al.* Psychological prehabilitation before cancer surgery: a systematic review. *Ann Surg Oncol* 2015;22:4117–23.
- 45 Clifford BK, Mizrahi D, Sandler CX, *et al.* Barriers and facilitators of exercise experienced by cancer survivors: a mixed methods systematic review. *Support Care Cancer* 2018;26:685–700.
- 46 Wilcox S, King AC, Brassington GS, *et al.* Physical activity preferences of middle-aged and older adults: a community analysis. *J Aging Phys Act* 1999;7:386–99.
- 47 Karvinen KH, Courneya KS, North S, *et al.* Associations between exercise and quality of life in bladder cancer survivors: a population-based study. *Cancer Epidemiol Biomarkers Prev* 2007;16:984–90.
- 48 Koebnick C, Michaud D, Moore SC, *et al.* Body mass index, physical activity, and bladder cancer in a large prospective study. *Cancer Epidemiol Biomarkers Prev* 2008;17:1214–21.
- 49 Kwan ML, Kushi LH, Danforth KN, *et al.* The Be-Well study: a prospective cohort study of lifestyle and genetic factors to reduce the risk of recurrence and progression of non-muscle-invasive bladder cancer. *Cancer Causes Control* 2019;30:187–93.