Fidelity, tolerability and safety of acute high-intensity interval training after hospitalisation for COVID-19: a randomised cross-over trial

Frederik Foged,¹ Iben Elmerdahl Rasmussen,¹ Josephine Bjørn Budde,^{1,2} Rasmus Syberg Rasmussen,¹ Villads Rasmussen,¹ Mark Lyngbæk,¹ Simon Jønck,¹ Rikke Krogh-Madsen,^{1,3,4} Birgitte Lindegaard,^{1,4,5} Mathias Ried-Larsen ^{1,6}, Ronan Martin Griffin Berg ^{1,2,7,8}, Regitse Højgaard Christensen^{1,9}

To cite: Foged F, Rasmussen IE, Bjørn Budde J, *et al.* Fidelity, tolerability and safety of acute high-intensity interval training after hospitalisation for COVID-19: a randomised cross-over trial. *BMJ Open Sport & Exercise Medicine* 2021;**7**:e001156. doi:10.1136/bmjsem-2021-001156

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2021-001156>).

FF and IER contributed equally.

Accepted 2 August 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Ronan Martin Griffin Berg; ronan@sund.ku.dk

ABSTRACT

Objectives. Many patients with COVID-19 suffer from persistent symptoms, many of which may potentially be reversed by high-intensity interval training (HIIT). Yet, the safety and tolerability of HIIT after COVID-19 is controversial. This study aimed to investigate the fidelity, tolerability and safety of three different HIIT protocols in individuals that had recently been hospitalised due to COVID-19.

Methods. The study was a randomised cross-over trial. We compared three supervised HIIT protocols (4×4, 6×1, 10-20-30) in 10 individuals recently discharged after hospitalisation for severe COVID-19. Each HIIT protocol had a duration of 38 min and was performed with a 1-week washout between them. Outcomes included adverse events, exercise training intensity and tolerability assessed by the Likert scale (1–10).

Results. All 10 participants aged 61 (mean, SD 8) years (5 males) completed all three HIIT protocols with no adverse events. High intensities were achieved in all three protocols, although they differed in terms of time spent with a heart rate ≥85% of maximum (mean (SD); 4×4: 13.7 (6.4) min; 10-20-30: 12.1 (3.8) min; 6×1: 6.1 (5.6) min; p=0.03). The three protocols were all well tolerated with similar Likert scale scores (mean (SD); 4×4: 8 (2), 10-20-30: 8 (2), 6×1: 9 (2), p=0.72).

Conclusion. Our findings indicate that recently hospitalised individuals for severe COVID-19 may safely tolerate acute bouts of supervised HIIT as per protocol. This warrants future studies testing the potential of regular HIIT as a rehabilitation strategy in this context.

INTRODUCTION

It has become increasingly clear that a substantial proportion of patients suffering from COVID-19 develop so-called 'long COVID-19', that is, often invalidating symptoms including dyspnoea, chest pain, exercise intolerance and extreme fatigue, to mention a few, all of which persist way beyond the acute viral infection.¹⁻⁴ The underlying

Key messages

What is already known

- Many patients suffering from COVID-19 develop so-called 'long COVID-19' which may be prevented or alleviated by targeted rehabilitation.
- High-intensity interval training (HIIT) is known to be a time-efficient rehabilitation intervention in several patient populations, notably in those with heart and lung disease.

What are the new findings

- Recently hospitalised patients who were mildly to slightly affected by long COVID-19 can satisfactorily complete acute exercise by the three most widely used HIIT protocols, so-called 4×4, 6×1 and 10-20-30 HIIT.
- Patients find HIIT both enjoyable and tolerable and we observed no adverse events.
- In patients recently hospitalised with COVID-19, the fidelity, tolerability and safety are comparable between the three HIIT protocols.

mechanisms of long COVID-19 are unknown but likely involve residual inflammation in the heart, lungs and vasculature.⁵ While the need for implementing evidence-based rehabilitation schemes that include exercise training following COVID-19 is imminent, no consensus exists as to how such programmes should be designed.

It is widely appreciated that high-intensity interval training (HIIT) promotes greater health-related benefits than traditional moderate-intensity training in various patient populations despite a reduced time commitment.⁶⁻⁸ However, in the context of COVID-19 rehabilitation and long COVID-19 prevention, it has emerged as a particularly controversial training modality. Hence, several opinion papers and guidelines favour low-intensity exercise with gradual increases

in intensity, mostly due to safety concerns.^{6 9–15} There is nonetheless currently no firm evidence base to argue in favour of this. A recent retrospective study on 28 discharged COVID-19 survivors conversely concluded that HIIT might safely be performed in these patients.¹⁶ Furthermore, in contrast to low-intensity and classical endurance exercise training, HIIT may more specifically target residual vascular inflammation, a putative central mechanism of long COVID-19.⁵ However, it is unknown to which extent it is feasible to implement HIIT in recently discharged patients with COVID-19, that is, whether they can tolerate HIIT without adverse effects.

In the present study, we assessed the fidelity, tolerability and safety of three widely used HIIT schemes in individuals recently discharged after hospitalisation for severe COVID-19.

METHODS

Study design, participants and randomisation

The study was performed from September to December 2020 at the Centre for Physical Activity Research at Rigshospitalet in Copenhagen, Denmark.

Ten recently hospitalised individuals who had recovered from the acute phase of severe COVID-19 infection were included. Severe acute respiratory distress syndrome coronavirus 2 was confirmed by a PCR from a pharyngeal swap in all cases, and the COVID-19 was considered as severe when it had required hospitalisation with oxygen supplementation. Recovery from COVID-19 was defined as the time of hospital discharge. In terms of eligibility, the inclusion criteria were hospitalisation-requiring COVID-19, age ≥ 40 years and ≤ 10 L oxygen requirement during hospitalisation. Exclusion criteria were atrial fibrillation/flutter, acute myocarditis, health conditions that prevent from participating in the exercise training intervention, and treatment with interleukin-6 receptor antagonists.

Participants underwent baseline testing, including a medical health interview and examination, arterial office blood pressure measurement, 12-lead ECG, a whole-body dual-energy X-ray absorptiometry, pulmonary function testing (dynamic spirometry, whole-body plethysmography and single-breath diffusing capacity to carbon monoxide). A maximal oxygen consumption ($\dot{V}O_{2\max}$) test was performed on a bicycle ergometer with a 5 min warm-up, immediately followed by a 15 Watt increase every minute until exhaustion. Participants completed a Post-COVID-19 Functional Status Scale (PFASS) questionnaire,¹⁷ which yields a score from 0 to 4, where 0 indicates no functional limitations, 1 negligible functional limitations, 2 slight functional limitations, 3 moderate functional limitations and 4 severe functional limitations.

The participants were allocated to the random sequence by which order they would complete and evaluate the three HIIT protocols in a cross-over design as described below. Randomisation was performed by a researcher with no other role in the study who generated a block randomisation schedule with balanced

permutations stratified by sex. The block sizes were not disclosed to ensure concealment, and participants were blinded for the sequence allocation until just before initiating the given HIIT session. To maintain blinding, the randomised sequence was delivered by email to a study investigator. The study investigator informed the participant before each training session which HIIT protocol to perform.

No changes in the trial design were made after the start of the experiments. It was not appropriate or possible to involve patients or the public in designing, conducting, reporting or disseminating our research. The original data that support the findings of this contribution can be obtained from the corresponding author on reasonable request.

Exercise training interventions

The three different HIIT protocols, including the specific training designs, are detailed in [table 1](#). The duration of the three HIIT protocols was matched in exercise duration (38 min). A 10 min warm-up was followed by a block of interval exercises, with a duration between 21 and 25 min depending on the specific protocol, aiming at reaching a specific percentage of maximal heart rate ($HR_{\max}$) for 4×4, maximal workload ($Watt_{\max}$) for 6×1 or subjective effort for 10-20-30 (further details in [table 1](#)). This was followed by a variable cool-down period (3–7 min) depending on the protocol. All exercise sessions were supervised by educated personnel.

Fidelity and tolerability

Fidelity was assessed by participant adherence to the prescribed training (dose reduction or dose modification) and the achieved intensity ([table 1](#)). The latter was evaluated by rate of perceived exertion (RPE) after each interval (6–20 Borg scale)¹⁸ and after completed training (1–10 Borg scale),¹⁹ as well as from minutes above 85% of $HR_{\max}$. Furthermore, enjoyment and perceived tolerance were evaluated by the 10-point Likert scale.²⁰

The primary qualitative evaluation of the HIIT protocols was an a priori defined composite endpoint based on three parameters: (1) participant adherence (fidelity), (2) a 10-point Likert scale (tolerability) and (3) adverse events (safety). This was displayed by a red-amber-green system inspired by a previous publication²¹ ([table 2](#)). The lowest of the three parameters determined the overall red-amber-green rank of the given HIIT protocol in the individual participant. No changes in outcomes were made after the start of the experiments.

Safety

At the baseline visit, all participants underwent a medical health interview and examination to assess health status. The following contraindications and precautions were considered: acute or recurring chest pain or systolic blood pressure >200 mm Hg±diastolic blood pressure >100 mm Hg. Safety during exercise training was assessed as the following, which also served as termination criteria:

Table 1 Structure of the three different high-intensity interval training protocols

	4×4	6×1	10-20-30
Total time	38 min	38 min	38 min
Warm-up	10 min warm-up with a heart rate of 60%–70% of HR_{max}	10 min warm-up on 30% of $Watt_{max}$	10 min warm-up at 60%–70% of HR_{max}
Intervals	Four intervals of 4 min at an intensity that will induce at least 85% of HR_{max} (starting at 75% of $Watt_{max}$). Each interval is separated by 3 min active pauses, biking at 50%–70% of HR_{max} .	Six intervals of 1 min at 100% of the $Watt_{max}$. Each interval is interspersed by 3 min active pauses at 30% of $Watt_{max}$.	Three intervals of 5 min interspersed by 3 min at 50%–70% of HR_{max} . Each interval set consists of 5 min of five repeated 30-20-10 intervals, consisting of 30 s at easy pace, 20 s at medium pace and 10 s at all-out.
Cool-down	3 min cool-down at ~warm up intensity	7 min cool-down at ~warm-up intensity	7 min cool-down at ~warm-up intensity
Success criteria	The participant must spend at least 25% of the interval above 85% of HR_{max} . The participant needs to reach 85% of HR_{max} during the first 3 min of each 4 min interval. If the participant does not reach 85% of HR_{max} , the last minute of the 4 min interval, the workload will be increased based on how far the heart rate is from the prescribed target. Likewise, the workload will be reduced if the RPM falls below 60 or in the next interval if the HR exceeds 95% of HR_{max} .	The participant must keep a pace above 60 rounds per minute during each interval. If the participant is not capable of keeping the right pace, the workload will be downgraded by 10% from the next interval.	The participant must spend at least 25% of each interval above 85% of HR_{max} .

HR_{max} , maximal heart rate; RPM, rounds per minute; $Watt_{max}$, maximal workload (in Watt).

onset of angina, signs of poor perfusion (confusion, nausea, lightheadedness, central cyanosis), requests to terminate (eg, due to intolerable dyspnoea), and physical or verbal manifestations of severe fatigue. A 12-lead ECG was obtained after each HIIT session to assess whether there was evidence of arrhythmia or myocardial ischaemia.

Statistics

The HIIT protocols were compared using analysis of covariance (ANCOVA) modelling for training intensity (minutes above 85% of HR_{max}), the Likert scale, RPE and PFASS. Unless otherwise stated, data are reported as mean (SD) or median (IQR). The sample size was

Table 2 Red-amber-green system designed to evaluate the three high-intensity interval training protocols

	Red	Amber	Green
Dose reduction	If ≤50% of the total training duration is completed	If 50%–85% of the training duration is completed	If ≥85% of the training duration is completed
Specific dose reduction for 4×4 min	If HR has been above 85% of max <15% of the time during the intervals (<2:20)	If HR has been above 85% of max 15%–24% of the time during intervals (2:20–3:50)	If HR has been above 85% of max ≥25% of the time during intervals (≥3:50)
Specific dose reduction for 6×1 min	If the intensity must be downgraded ≥10% in five or more of the intervals	If the intensity must be downgraded 10% in 3–4 of the intervals	If the intensity must be downgraded 10% in <3 of the intervals
Specific dose reduction for 10-20-30	If HR has been above 85% of max <15% of the time during the intervals (<2:15)	If HR has been above 85% of max 15%–24% of the time during intervals (2:15–3:40)	If HR has been above 85% of max ≥25% of the time during intervals (≥3:40)
Likert scale (tolerability)	1–2	3–4	5+
Adverse events	If the participant experiences angina, must stop the training due to any adverse event or has ST depression or arrhythmia on the post-training ECG	If ≥3 adverse events are registered during the training session	If ≤2 adverse events are registered during the training session

HR, heart rate.

Figure 1 Study flow chart. Flow chart of participants included in the study.

determined on what was feasible within the local context under considerations of reducing the number of participants to exposure of potential harms. All analyses were performed using GraphPad Prism, V.7.02 (GraphPad Software) or STATA/SE (StataCorp) V.16.1, and a two-tailed $p < 0.05$ was considered statistically significant.

RESULTS

The study flow diagram is shown in [figure 1](#). A total of 10 participants were enrolled and randomised and complied with the three HIIT protocols, and all participants were thus included in the analysis (attendance 100%). Baseline characteristics, including lung function and PFASS scores, are presented in [table 3](#). All participants except for one with a restrictive pattern had normal ventilatory capacity, while five had reduced diffusing capacity (mild reduction in three cases, moderate-to-severe in two cases). Generally, the participants were only mildly to slightly affected by persisting COVID-19 symptoms according to PFASS scores, with the highest score of 3 in one individual. The primary reason for not participating was feeling too weak to exercise after the infection.

Outcomes of the three HIIT training protocols were compared by a red-amber-green design ([table 2](#)). No 'red' training sessions occurred, and only 1/10 training session in 4×4 was amber due to a low Likert score (tolerable: 3/10), and 1/10 in the 10-20-30 was amber due to inadequate time spent $>85\%$ of HR_{max} as prescribed. The rest of the training sessions were reported as green, which means all three HIIT protocols are tolerable and safe ([figure 2](#)). No adverse events were reported during the study, and no angina or ECG changes were seen after HIIT training.

There was a between-protocol difference in terms of the exercise intensity when dichotomously assessing for time spent with a heart rate $\geq 85\%$ of HR_{max} ([figure 3A](#)). In contrast, tolerability as assessed by Likert scale scores ([figure 3B](#)) and RPE (4×4; 5 (2), 6×1; 5 (2), 10-20-30; 6 (2); $p=0.28$) did not differ between protocols. When considering HR and RPE changes non-dichotomously,

Table 3 Baseline characteristics

Measurement (mean (SD) or median (IQR))	
Age (years)	61 (8)
Height (cm)	172 (7)
Weight (kg)	78.3 (10.8)
Fat mass (%)	35.5 (7.7)
BMD (T-score)	1.22 (0.07)
FEV ₁ (%-pred)	110 (19)
FVC (%-pred)	112 (16)
FEV ₁ /FVC	81 (10)
RV (%-pred)	89 (21)
TLC (%-pred)	94 (15)
D _L COc (%-pred)	78 (20)
VO _{2max} (mL/min)	1968 (766)
VO _{2max} (mL/min/kg)	24.9 (8.8)
Duration of hospitalisation (days)	7 (4)
Time since discharge (days)	40 (22–145)
PFASS	2 (0–2)

Anthropometry measured at baseline.

BMD, bone mass density; D_LCOc, carbon monoxide diffusing capacity corrected for haemoglobin; PFASS, Post-COVID-19 Functional Status Scale; RPE, rate of perceived exertion; RV, residual volume; TLC, total lung capacity; VO_{2max}, maximal oxygen consumption.

largely similar objective and subjective measures of intensities were observed throughout each protocol (online supplemental figures 1 and 2).

DISCUSSION

This study compared the fidelity, tolerability and safety of three HIIT protocols conducted on individuals recently hospitalised with COVID-19. Our findings indicate that HIIT is feasible regarding fidelity, tolerability and safety, regardless of the specific protocol.

No adverse events occurred in any of the HIIT protocols. These results are in line with a previous study on COVID-19 survivors,¹⁶ as well as several studies on other 'high risk' patient groups such as ischaemic heart disease and heart failure in whom it is considered safe.^{6 22} Thus, we consider it unlikely that HIIT should be associated with a high risk of acute adverse cardiovascular events. This concern was raised in The Stanford Consensus Statement for Post-COVID-19 Rehabilitation as one of the primary arguments against HIIT use in these individuals.²³ However, studies with regular exercise are needed to determine the long-term safety of HIIT in this context.

We found that while all three HIIT protocols were well tolerated, the greatest intensity, given by the time spent above 85% of HR_{max} , was achieved by 4×4. Time spent above 85% of HR_{max} was chosen since HIIT is widely defined as training intensities at 85%–90% of HR_{max} .^{7 24} The cut-off, 85% of HR_{max} , is arbitrarily defined, and the

Figure 2 Red-amber-green. Number of 'red', 'amber' and 'green' training sessions of the three high-intensity interval training protocols.

supplemental graphic overview shows that all protocols performed well with high RPE and HR reaching maximum in all protocols. These findings, combined with the strong physiological rationale for HIIT, highlight it as an attractive option for COVID-19 rehabilitation. This is further invigorated when considering the widespread and disease-specific vascular inflammation in COVID-19 that persists after the acute viral infection and which is thought to contribute to the very diverse cluster of symptoms associated with long COVID-19.²⁵ Potentially, this may be specifically counteracted with HIIT, where the pronounced changes in vascular shear stress that occur between intervals combined with the effects of anti-inflammatory myokines that are released from working skeletal muscle in an intensity-dependent manner may enhance endothelial repair and dampen proinflammatory processes within the vascular wall.⁵

Figure 3 Intensity and tolerability. (A) Intensity: time spent with a heart rate >85% of maximum for each protocol. (B) Tolerability rated by the Likert scale. Data are presented as mean (95% CI); * $p=0.028$. HR_{max}, maximal heart rate.

The high composite intensity achieved is the best-documented means for increasing maximal oxygen consumption ($\dot{V}O_{2max}$) and exerting beneficial cardiovascular effects. HIIT has thus been reported to be superior to continuous endurance training in several patient groups.^{7,26} These include patients with hypertension, ischaemic heart disease and heart failure, in which $\dot{V}O_{2max}$ has furthermore been noted as the single best predictor of mortality.^{27,28} It remains to be determined whether this is also the case in various lung diseases, such as chronic obstructive pulmonary disease. Still, HIIT has been noted to be associated with higher adherence in these patients, conceivably because the intermittent nature of HIIT reduces ventilatory demand and thus exertional dyspnoea and perceived breathing effort.^{29,30} This may also apply to patients recovering from COVID-19. Even though the present study is small scale in this context, and larger randomised controlled studies are warranted, we posit that HIIT should also be considered among the preferred training modalities in the rehabilitation post-COVID-19.

A strength of the study is the randomised sequence of the exercise training protocols, which eliminated systematic carry-over effects of the previous training protocol. Another strength is that this is the first study testing HIIT protocols post-COVID-19 in a supervised manner. Moreover, our research investigated the fidelity, tolerable and safety of three various HIIT protocols with different work or intensity periods, thus enabling between-protocol comparisons.

Some limitations should be considered when interpreting the results of this study. First, our results may be affected by selection bias because the main reason for declining participation was feeling too weak to exercise, which may indeed be a symptom of long COVID-19. Our results regarding fidelity and tolerability may thus have been different if we had studied these patients, which arguably represent the group that may benefit the most from targeted rehabilitation. Another important limitation is the small study size ($n=10$) which limits the conclusions, particularly when considering the heterogeneity of the participants. Furthermore, we only tested the fidelity, tolerability and safety to acute HIIT and cannot draw any firm conclusions regarding training over extended periods. Even though we expect a HIIT-based rehabilitation scheme over an extended period to be feasible, this remains speculative.

In conclusion, we found that individuals recently recovered from COVID-19 could successfully and safely complete three acute bouts of the most widely used HIIT protocols. Our findings highlight that HIIT should be evaluated as a rehabilitation strategy for post-COVID-19 in future randomised controlled trials.

Author affiliations

¹Centre for Physical Activity Research, Rigshospitalet, Copenhagen, Denmark

²Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

³Department of Infectious Diseases, Hvidovre Hospital, Hvidovre, Denmark

⁴Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁵Department of Pulmonary Medicine and Infectious Diseases, Hillerød Hospital, Hillerød, Denmark

⁶Institute of Sports Science and Clinical Biomechanics, University of Southern Denmark Faculty of Health Sciences, Odense, Denmark

⁷Department of Clinical Physiology, Nuclear Medicine & PET, Rigshospitalet, Copenhagen, Denmark

⁸Neurovascular Research Laboratory, Faculty of Life Sciences and Education, University of South Wales, Pontypridd, UK

⁹Department of Cardiology, Rigshospitalet, Copenhagen, Denmark

Twitter Ronan Martin Griffin Berg @cresp_research @ronan_berg @cfas_dk

Contributors Conceptualisation: RHC, RMGB, MR-L. Methodology: RHC, RMGB, VR, MR-L, RKM, BLM. Formal analysis: FF, IER, RHC, RMGB, MR-L. Investigation: FF, IER, JBB, RSR, ML, SJ, RHC, RMGB, BLM, RKM. Supervision: RMGB, RHC. Resources: RMGB, RHC. Writing—original draft: FF, IER. Writing—review and editing: All authors contributed. Visualisation: FF, IER, MR-L.

Funding The Centre for Physical Activity Research (CFAS) is supported by a grant from TrygFonden, and the Centre of Inflammation and Metabolism (CIM) is a member of DD2—the Danish Center for Strategic Research in Type 2 Diabetes (the Danish Council for Strategic Research, grant nos. 09-067009 and 09-075724).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval The study was conducted per the Declaration of Helsinki with oral and written informed consent from all participants before inclusion and approved by the Scientific Ethical Committee of the Capital Region of Denmark (H-2003733 including amendment 75068 and 75799) and the Danish Data Protection Agency (P-2020-781) and pre-registered at ClinicalTrials.gov (NCT04549337).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The original data that support the findings of this contribution can be obtained from the corresponding author on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Mathias Ried-Larsen <http://orcid.org/0000-0002-8388-5291>

Ronan Martin Griffin Berg <http://orcid.org/0000-0002-5757-9506>

REFERENCES

- Alwan NA, Attree E, Blair JM, *et al.* From doctors as patients: a manifesto for tackling persisting symptoms of covid-19. *BMJ* 2020;370:m3565.
- Bos LDJ, Brodie D, Calfee CS. Severe COVID-19 Infections—Knowledge gained and remaining questions. *JAMA Intern Med* 2021;181:9–11.
- Rubin R. As Their Numbers Grow, COVID-19 “Long Haulers” Stump Experts. *JAMA* 2020;324:1381–5.
- Carfi A, Bernabei R, Landi F, *et al.* Persistent symptoms in patients after acute COVID-19. *JAMA* 2020;324:603–5.
- Christensen RH, Berg RMG. Vascular Inflammation as a Therapeutic Target in COVID-19 “Long Haulers”: HIITing the Spot? *Front Cardiovasc Med* 2021;8:643626.
- Batacan RB, Duncan MJ, Dalbo VJ, *et al.* Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies. *Br J Sports Med* 2017;51:494–503.
- Karlsen T, Aamot I-L, Haykowsky M, *et al.* High intensity interval training for maximizing health outcomes. *Prog Cardiovasc Dis* 2017;60:67–77.
- Gillen JB, Gibala MJ. Is high-intensity interval training a time-efficient exercise strategy to improve health and fitness? *Appl Physiol Nutr Metab* 2014;39:409–12.
- Leandro CG, Ferreira E Silva WT, Lima-Silva AE. Covid-19 and exercise-induced immunomodulation. *Neuroimmunomodulation* 2020;27:75–8.
- Rahmati-Ahmadabad S. Exercise against SARS-CoV-2 (COVID-19): does workout intensity matter? *Obes Med* 2020;19:100245.
- Bhatia RT, Marwaha S, Malhotra A, *et al.* Exercise in the Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) era: A Question and Answer session with the experts Endorsed by the section of Sports Cardiology & Exercise of the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2020;27:1242–51.
- Verwoert GC, de Vries ST, Bijsterveld N, *et al.* Return to sports after COVID-19: a position paper from the Dutch Sports Cardiology Section of the Netherlands Society of Cardiology. *Neth Heart J* 2020;28:391–5.
- Dores H, Cardim N. Return to play after COVID-19: a sport cardiologist's view. *Br J Sports Med* 2020;54:1132–3.
- Denay KL, Breslow RG, Turner MN. ACSM call to action statement: COVID-19 considerations for sports and physical activity. *N Engl J Med* 2020;383:120–8.
- Kennedy FM, Sharma S. COVID-19, the heart and returning to physical exercise. *Occup Med* 2020;70:19–21.
- Hermann M, Pekacka-Egli A-M, Witassek F, *et al.* Feasibility and efficacy of cardiopulmonary rehabilitation after COVID-19. *Am J Phys Med Rehabil* 2020;99:865–9.
- Klok FA, Boon GJAM, Barco S, *et al.* The Post-COVID-19 functional status scale: a tool to measure functional status over time after COVID-19. *Eur Respir J* 2020;56:2001494.
- Borg GAV. Psychophysical bases of perceived exertion. *Plast Reconstr Surg* 1954;14:377–81.
- Shariat A, Cleland JA, Danaee M, *et al.* Borg CR-10 scale as a new approach to monitoring office exercise training. *Work* 2018;60:549–54.
- Afthanorhan A. The Likert scale analysis using parametric based structural equation modeling (SEM). *Computational Methods in Social Sciences* 2016;4:13–21.
- Avery KNL, Williamson PR, Gamble C, *et al.* Informing efficient randomised controlled trials: exploration of challenges in developing progression criteria for internal pilot studies. *BMJ Open* 2017;7:e013537.
- Giallauria F, Piccioli L, Vitale G, *et al.* Exercise training in patients with chronic heart failure: a new challenge for cardiac rehabilitation community. *Monaldi Arch Chest Dis* 2018;88:38–44.
- Barker-Davies RM, O'Sullivan O, Senaratne KPP, *et al.* The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *Br J Sports Med* 2020;54:949–59.
- Wisløff U, Ellingsen Øyvind, Kemi OJ. High-Intensity interval training to maximize cardiac benefits of exercise training? *Exerc Sport Sci Rev* 2009;37:139–46.
- Sollini M, Ciccarelli M, Cecconi M, *et al.* Vasculitis changes in COVID-19 survivors with persistent symptoms: an [¹⁸F]FDG-PET/CT study. *Eur J Nucl Med Mol Imaging* 2021;48:1460–6.
- Helgerud J, Høydal K, Wang E, *et al.* Aerobic high-intensity intervals improve VO₂max more than moderate training. *Med Sci Sports Exerc* 2007;39:665–71.
- Myers J, Prakash M, Froelicher V, *et al.* Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346:793–801.
- Kavanagh T, Mertens DJ, Hamm LF, *et al.* Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. *Circulation* 2002;106:666–71.
- Sawyer A, Cavalheri V, Hill K. Effects of high intensity interval training on exercise capacity in people with chronic pulmonary conditions: a narrative review. *BMC Sports Sci Med Rehabil* 2020;12:22.
- Kortianou EA, Nasis IG, Spetsioti ST, *et al.* Effectiveness of interval exercise training in patients with COPD. *Cardiopulm Phys Ther J* 2010;21:12–19.