

# Association of sedentary and physical activity behaviours with body composition: a genome-wide association and Mendelian randomisation study

Ferris A Ramadan <sup>1</sup>, Jennifer W Bea<sup>2,3</sup>, David O Garcia,<sup>4</sup> Katherine D Ellingson,<sup>1</sup> Robert A Canales,<sup>5</sup> David A Raichlen,<sup>6</sup> Yann C Klimentidis<sup>1</sup>

**To cite:** Ramadan FA, Bea JW, Garcia DO, *et al.* Association of sedentary and physical activity behaviours with body composition: a genome-wide association and Mendelian randomisation study. *BMJ Open Sport & Exercise Medicine* 2022;**8**:e001291. doi:10.1136/bmjsem-2021-001291

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2021-001291>).

Accepted 7 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Ferris A Ramadan;  
ferris@email.arizona.edu

## ABSTRACT

**Objectives** Studies suggest that body composition can be independently improved through physical activity (PA). We performed a Mendelian randomisation (MR) study to test the incremental benefits of sedentary behaviour and various PA exposures on body composition outcomes as assessed by anthropometric indices, lean body mass (kg), body fat (%) and visceral adipose tissue (VAT) (kg).

**Methods** Genetic instruments were identified for both self-reported and accelerometer-measured sedentary behaviour and PA. Outcomes included anthropometric and dual-energy X-ray absorptiometry measures of adiposity, extracted from the UK Biobank and the largest available consortia. Multivariable MR (MVMR) included educational attainment as a covariate to address potential confounding. Sensitivity analyses were evaluated for weak instrument bias and pleiotropic effects.

**Results** We did not identify consistent associations between genetically predicted self-reported and accelerometer-measured sedentary behaviour and body composition outcomes. All analyses for self-reported moderate PA were null for body composition outcomes. Genetically predicted PA at higher intensities was protective against VAT in MR and MVMR analyses of both accelerometer-measured vigorous PA (MVMR  $\beta$ =−0.15, 95% CI: −0.24 to −0.07,  $p$ <0.001) and self-reported participation in strenuous sports or other exercises (MVMR  $\beta$ =−0.27, 95% CI: −0.52 to −0.01,  $p$ =0.034) was robust across several sensitivity analyses.

**Conclusions** We did not identify evidence of a causal relationship between genetically predicted PA and body composition, with the exception of a putatively protective effect of higher-intensity PA on VAT. Protective effects of PA against VAT may support prior evidence of biological pathways through which PA decreases risk of downstream cardiometabolic diseases.

## INTRODUCTION

Obesity has reached critical levels and continues to exacerbate pervasive public health concerns, including cardiovascular diseases, cancers, type 2 diabetes, mental health and discrimination.<sup>1</sup> This high prevalence has been ascribed to an increasingly

## WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Previous studies have shown a limited protective association between physical activity (PA) and body composition measures.
- ⇒ Observational data suggest PA may protect against visceral adipose tissue (VAT), a stronger predictor of morbidity and mortality compared with subcutaneous adipose tissue.
- ⇒ Recent Mendelian randomisation studies have suggested a bidirectional positive association between body mass index and sedentary behaviour, with some evidence suggesting a protective effect from PA on body composition measures.

## WHAT THIS STUDY ADDS

- ⇒ We did not identify consistent protective associations between genetically predicted sedentary behaviour or PA exposures and body composition outcomes.
- ⇒ Genetically predicted accelerometer-measured PA and self-reported strenuous sports and other exercises were protective against VAT.
- ⇒ Our findings align with previous evidence suggesting enhanced lipolysis in VAT due to increased secretion of catecholamines during vigorous PA.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ National and international health agencies' emphasis on moderate PA for weight management may be insufficient for independently improving anthropometric indices, lean body mass or total body fat per cent.
- ⇒ Higher-intensity levels of PA may independently reduce VAT and improve an individual's metabolic profile.
- ⇒ Future investigations into the effect of PA on other fat measures such as subcutaneous adipose tissue, ectopic fat or hepatic fat may reveal additional insights into the specific cardiometabolic benefits of PA.

obesogenic environment, characterised by reduced physical activity (PA), increased occupational and recreational sedentary activities, and dietary options inundated by highly palatable energy-dense foods.<sup>2</sup> In an effort to

reduce body fat, weight management programmes have principally targeted both a reduction in energy intake and increased PA levels. Although PA offers substantial benefits to overall health,<sup>3</sup> these efforts have largely fallen short in attenuating obesity rates.<sup>4–6</sup>

Interventions have examined the putative effect of PA on various body composition measures, such as body mass index (BMI), waist circumference (WC), waist-to-hip ratio (WHR) and total body fat per cent (%TBF). Combined diet and exercise programmes are optimal for improving body composition versus diet-only programmes<sup>7</sup>; however, PA alone inconsistently protects against clinically significant weight gain.<sup>8</sup> Previous studies examining the effect of PA on lean body mass (LBM)—an aggregate body mass measurement excluding fat—have similarly yielded mixed outcomes.<sup>9–11</sup> Current evidence suggests that at least 150 min/week spent in moderate PA offers the greatest protection against weight gain.<sup>12–13</sup> Although there is limited evidence indicating a dose–response relationship between PA and weight gain, PA programmes prescribing higher-intensity exercise have resulted in greater mass reduction, compared with lower-intensity programmes.<sup>14–15</sup> Of particular importance for improving body composition is reducing visceral adipose tissue (VAT), which is a stronger predictor of mortality compared with non-specific anthropometric and subcutaneous fat measures.<sup>16</sup> Observational studies have supported significant benefits of PA on VAT reduction, independently of diet.<sup>17–20</sup> High-intensity interval training interventions have further demonstrated VAT reductions with similar results to mixed diet and PA programmes.<sup>21</sup> Understanding the effect of PA on VAT thus holds particular value in attenuating the risk of associated downstream diseases.

Mendelian randomisation (MR) is a form of instrumental variable analysis that leverages genetic variants


from genome-wide association studies (GWAS) to infer causality between an exposure and a given outcome.<sup>22</sup> The assignment of genetic variants during gamete formation confers a lifelong exposure for individuals randomised to a given allelic variant. Unlike exposures measured at the phenotype level, genetic variants are not typically associated with a multitude of behavioural, social or physiological factors that may confound an epidemiological association.<sup>23</sup> MR may thus avoid many of the pitfalls seen in previous observational studies, including confounding<sup>24</sup> and reverse causation.<sup>25</sup> Previous MR analyses have suggested a positive bidirectional association between sedentary behaviour and BMI,<sup>26</sup> and a negative bidirectional association between accelerometer-measured overall activity<sup>27</sup> and moderate and vigorous PA on BMI.<sup>28</sup>

In the present study, we performed MR using both self-reported and accelerometer-measured sedentary behaviour and PA on a wider set of body composition outcomes. Current MR studies have not evaluated self-reported and accelerometer-measured activity in parallel, or the effect of sedentary behaviour and PA on more specific body composition measures. Here, we aim for a more granular understanding of the benefit, if any, of activity levels on body composition.

## METHODS

### Study design

We performed MR and multivariable MR (MVMR) analyses to evaluate the dose–response relationship between sedentary behaviour, PA and (1) whole LBM, (2) BMI, (3) WC, (4) WHR, (5) %TBF and (6) VAT (figure 1). Exposure and outcome instruments were derived from non-overlapping samples to facilitate application of a two-sample MR study, which is less prone to weak instrument bias compared with one-sample MR. Both



**Figure 1** Sedentary behaviour and physical activity (PA) exposures and body composition outcomes evaluated in Mendelian randomisation analyses. <sup>†</sup>GWAS author and publication year are reported adjacent to exposure and outcome phenotypes where applicable. <sup>†</sup>Self-reported exposure. <sup>§</sup>Accelerometer-measured exposure. GWAS, genome-wide association studies.

self-reported and accelerometer-measured exposures were used to evaluate PA associations with body composition measures. Self-reported and fraction acceleration thresholds were selected based on metabolic equivalent of tasks (METs), including sedentary behaviour ( $\leq 1.5$  METs), moderate PA (3.0–5.9 METs), moderate-to-vigorous PA (MVPA) ( $\geq 3.0$  METs) and vigorous PA ( $\geq 6.0$  METs). We did not consider the light PA phenotype due to the limited measurement sensitivity of these measurements. Self-reported strenuous sports or other exercises (SSOEs) and overall acceleration measures were further included in analyses due to evidence suggesting strong genetic markers for these exposures.<sup>29</sup> MVMR analyses included educational attainment as a covariate, as described below. This study is reported according to the Strengthening the Reporting of Observational Studies in Epidemiology using MR guidelines.<sup>30</sup>

Genetic instruments were identified from GWAS from the UK Biobank (UKB), GENetic Factors for Osteoporosis (GEFOS) consortium<sup>31</sup> and the Genetic Investigation of Anthropometric Traits (GIANT) consortium.<sup>32</sup> The UKB is an ongoing prospective cohort study in the UK. Approximately half a million adults, aged 37–73 years, have been enrolled and provided genotype and phenotype data to investigators. Neither patients nor the public were involved in the design, conduct, reporting or dissemination of this research.

### Exposure measures

Self-reported PA was previously captured for over 377 000 UKB participants via in-person questionnaires at the baseline examination (2006–2010). Self-reported total sedentary behaviour was not measured directly in the UKB. Therefore, we used a combined proxy measure of sedentary behaviours, including average self-reported hours of leisure television watching, leisure computer use and driving time ( $n=422\,218$ ).<sup>33</sup> These behaviours show higher validity than total sedentary behaviour and are easier to recall.<sup>34</sup>

Self-reported light PA was not included in analyses due to low specificity of available UKB phenotypes. Self-reported moderate and vigorous PAs were captured directly in the UKB. MVPA was calculated using the International Physical Activity Questionnaire scoring protocol by taking the sum of total minutes/week of moderate PA multiplied by 4 and the total number of vigorous PA minutes/week multiplied by 8.<sup>35 36</sup> A summary of these phenotypes and quality control procedures has been previously described by Klimentidis *et al.*<sup>29</sup>

Acceleration data in the UKB were captured for over 90 000 participants over 1-week duration with the Axivity AX3 wrist-worn triaxial accelerometer.<sup>27</sup> Data were calibrated, resampled and summarised using previously outlined procedures.<sup>37</sup> Accelerometer-derived PA can be converted to METs using fraction of acceleration time, measured in 5 s epochs of milligravities (mg). Sedentary and moderate PA METs were identified by Doherty *et al.*<sup>27</sup> on accelerometer data using predicted behaviour states

from a machine-learning model trained on free-living participants. We did not derive a light PA phenotype due to limited accelerometer sensitivity at lower-level accelerations, which may have introduced ambiguity in differentiating between activity states. MVPA was defined as the fraction of accelerations above 100 mg.<sup>38</sup> Vigorous PA was defined as the fraction of accelerations above 425 mg.<sup>38</sup> In summary, accelerometer-measured sedentary and moderate PA phenotypes were isolated using predicted behaviour states, while MVPA and vigorous PA phenotypes were established from acceleration cut-off points corresponding to METs.

Summary statistics were previously unavailable for self-reported moderate PA and accelerometer-measured PA. We conducted GWAS on 360 911 participants with self-reported moderate PA, and 97 737 participants with accelerometer-measured MVPA using the BOLT-LMM V.2.3.6 software tool.<sup>39</sup> Briefly, participants were excluded based on  $>5\%$  missing rates, unusually high heterozygosity, and a mismatch between self-reported and genetically inferred sex. To limit confounding from population-level stratification, only self-reported European participants were included in GWAS. Single nucleotide polymorphisms (SNPs) with high missingness ( $>1.5\%$ ), genetic versus self-reported sex mismatches, low minor allele frequency ( $<0.1\%$ ) and low imputation quality were excluded from analyses.

In all models, we controlled for covariates including genotyping chip, centre, age-squared, age, sex, age–sex interaction, and the first 10 principal components to correct for population stratification and genotyping array. Educational attainment was collected from multiple cohorts as a continuous measure of years of school completed ( $n=293\,723$ ).<sup>40</sup> We previously identified strong genetic correlations of education with PA.<sup>29 41</sup> MVMR analyses included educational attainment as a covariate to minimise potential confounding.

### Outcome measures

Outcome measurements were extracted from non-overlapping studies. The LBM phenotype used in the GEFOS consortium was measured in kilograms (kg) using bioimpedance analysis (BIA) or dual energy X-ray absorptiometry (DEXA) ( $n=38\,292$ ).<sup>42</sup> Despite lower specificity from BIA measurements, BIA and DEXA show strong agreement and can be used interchangeably at the population level.<sup>43</sup> Summary statistics of GWAS (not including the UKB) for anthropometric outcomes including BMI ( $\text{kg}/\text{m}^2$ ) ( $n=249\,796$ ), WC adjusted for BMI (cm) and WHR adjusted for BMI ( $n=224\,459$ ) were obtained from the GIANT consortium.<sup>32</sup> We also used a GWAS meta-analysis that did not overlap with the UKB to extract summary statistics for %TBF ( $n=89\,297$ ).<sup>44</sup> The %TBF phenotype was measured with either BIA or DEXA. Large-scale GWAS of VAT has been limited, however, due to the need for advanced imaging phenotypes on large numbers of individuals to reveal significant loci. To reach a sample size that would yield greater genetic signal, we



applied a recently developed non-linear VAT prediction model developed by Karlsson *et al.*<sup>45</sup> Briefly, the model was trained on 4000 UKB participants with DEXA-measured VAT (kg). BIA measures on the other UKB participants were used as the primary predictive features within models. Despite a reduced sensitivity, the consistent relationship between BIA-derived measures and measured VAT suggests heightened model performance and indicated strong predictive capacity (coefficient of determination ( $r^2$ )=0.76).<sup>45</sup> We performed GWAS for VAT on a subset of non-overlapping UKB participants including those without self-reported PA ( $n=56\,908$ ), and accelerometer measurements ( $n=323\,769$ ) (online supplemental table S1).

### Instrument selection

We selected SNPs at  $p < 5 \times 10^{-6}$  for exposures. SNPs in linkage disequilibrium with the top SNP were removed to account for correlated genetic variants ( $r^2 < 0.001$ ). Proxies for genetic variants missing in outcome GWAS were identified using the 1000 Genomes European sample data. Alleles were harmonised to ensure that the same allele was being referenced for exposure and outcome effect sizes. We attempted to infer the positive strand allele using allele frequencies for palindromic SNPs. We did not manually 'prune' for genetic variants with suspected associations with outcome phenotypes as it risks the removal of only instrumental variables with a directional effect.<sup>46</sup> We were unable to identify an adequate number of SNPs for accelerometer-measured moderate PA exposures, and for sedentary behaviour with LBM, and therefore removed these associations from analyses.

We performed Steiger filtering as a sensitivity analysis to remove genetic variants explaining greater variance in the exposure trait compared with the outcome. To account for likely biases from the *APOE* variant as previously described,<sup>29</sup> we performed sensitivity analyses removing instrumental variables within 500 kb of the *APOE* rs429358 variant. Briefly, the rs429358 variant is strongly associated with low-density lipoprotein cholesterol and Alzheimer's disease meaning that participants (especially older ones) with these risk variants and who are still alive and incorporated in the UKB are likely enriched for healthy behaviours that have counterbalanced their genetic risk from *APOE*, leading to a form of survival/selection bias.<sup>29</sup> It is therefore strongly suspected that the association of the *APOE* variant with self-reported MVPA is an artefact of this bias.

### Analyses

MR associations were first estimated with the inverse variance weighted (IVW) linear regression. MVMR analyses were also conducted to address potential confounding of the PA–body composition association by educational attainment. Causal estimates were reported as regression coefficients with 95% CIs. Body composition outcomes reflect a one-unit incremental change in self-reported and

accelerometer-based exposure variables. Post hoc sensitivity analyses were performed using MR-Egger, median and mode-based estimates, MR pleiotropy residual sum and outlier (MR-PRESSO)<sup>47</sup> and causal analysis using summary effect estimates (CAUSE).<sup>48</sup>


The validity of MR estimates is dependent on three core assumptions: (1) the relevance assumption, meaning selected variants must be directly associated with the exposure; (2) the independence assumption that unmeasured confounders are not associated with the outcome; (3) and the exclusion restriction assumption that genetic variants are not pleiotropically associated with the exposure.

We used the conventional F-statistic threshold of 10 to evaluate instrument strength. We performed visual assessments of IVW regression, funnel plots and leave-one-out (LOO) analyses to identify individual variants influencing causal effects. Heterogeneity was assessed for all MR and MVMR analyses using a modified Cochran's Q statistic.<sup>49</sup> Outcome and exposure measures were reversed to evaluate for bidirectionality in MR. Associations were considered (statistically) significant at the  $p < 0.05$  threshold with no correction for multiple testing. Although we recognise that multiple tests are being performed, we evaluate our results based on the consistency of estimates across multiple exposures, outcomes and sensitivity tests. MR analyses and quality control procedures were performed using the 'TwoSampleMR',<sup>50</sup> 'MVMR',<sup>51</sup> and 'cause' R packages.<sup>48</sup> Analyses were performed in R V.3.6.3 (R Foundation for Statistical Computing).<sup>52</sup>

### RESULTS

GWAS for self-reported moderate PA revealed one novel genome-wide significant SNP, rs6545389 ( $p=3.5 \times 10^{-8}$ ). GWAS of accelerometer-measured MVPA revealed two novel loci including rs17443704 ( $p=3.8 \times 10^{-9}$ ) and rs4480415 ( $p=8.7 \times 10^{-9}$ ), as well as a near genome-wide significant effect for rs791273 ( $p=5.0 \times 10^{-8}$ ). Additional details on these variants are provided in the online supplemental figures S1 and S2.

Univariable MR analyses suggested self-reported sedentary behaviour was positively associated with anthropometric indices and VAT, while no association was identified with LBM (figure 2). After controlling for educational attainment in MVMR, sedentary behaviour associations were null for all measures except an inverse association with WHR ( $\beta=-0.25$ , 95% CI:  $-0.02$  to  $-0.48$ ), and a positive association with VAT ( $\beta=0.38$ , 95% CI:  $0.27$  to  $0.49$ ) (figure 3). Self-reported sedentary behaviour was significantly associated with VAT across sensitivity analyses including CAUSE ( $\gamma=0.23$ , 95% CI:  $0.19$  to  $0.28$ ) (table 1). MR analyses for the accelerometer-measured sedentary phenotype similarly identified a significant positive association with BMI ( $\beta=0.36$ , 95% CI:  $0.02$  to  $0.71$ ), while other anthropometric outcomes and VAT were null (figure 4). CAUSE did not identify an association between accelerometer-measured sedentary behaviour




**Figure 2** Univariable Mendelian randomisation IVW estimates for self-reported physical activity (PA) levels and body composition. %TBF, total body fat per cent; BMI, body mass index; IVW, inverse variance weighted; LBM, lean body mass; METs, metabolic equivalent of tasks; MVPA, moderate-to-vigorous PA; SNPs, single nucleotide polymorphisms; SSOE, strenuous sports or other exercises; VAT, visceral adipose tissue; WC, waist circumference; WHR, waist-to-hip ratio.

and VAT (table 1). After controlling for educational attainment, however, no associations were identified for the accelerometer-measured sedentary phenotype (figure 5).

MR for self-reported moderate PA revealed a singular positive association with WC ( $\beta=0.22$ , 95% CI: 0.03 to 0.41) (figure 2), while no associations were identified in MVMR. MR for self-reported MVPA was not associated with WC ( $\beta=-0.10$ , 95% CI:  $-0.22$  to  $0.02$ ) while a significant association was identified with VAT ( $\beta=-0.09$ , 95% CI:  $-0.14$  to  $-0.03$ ). However, MVMR analyses did not identify a protective association of MVPA with any body composition measure (figure 3). MR analysis of accelerometer-measured MVPA was negatively associated with %TBF and VAT (figure 4). After controlling for educational attainment, accelerometer-measured MVPA was only protective against VAT ( $\beta=-0.12$ , 95% CI:  $-0.19$  to  $-0.04$ ) (figure 5). We additionally identified a negative association between accelerometer-measured MVPA and VAT in CAUSE ( $\gamma=-0.11$ , 95% CI:  $-0.15$  to  $-0.07$ ) (table 1).

MR analysis of self-reported vigorous PA suggested a protective association with VAT, while no relationship was identified in MVMR analysis ( $\beta=-0.02$ , 95% CI:  $-0.34$  to  $0.29$ ) (figure 3). CAUSE did not suggest an association between vigorous PA and VAT ( $\gamma=-0.13$ , 95% CI:  $-0.33$  to  $0.06$ ) (table 1). In contrast to VAT, no association was identified between vigorous PA and BMI in MR, while MVMR indicated a positive association with BMI ( $\beta=0.44$ , 95% CI: 0.03 to 0.85) (figure 3). No associations were identified between self-reported vigorous PA and other body composition measures. MR analysis of accelerometer-measured vigorous PA demonstrated a protective association with %TBF ( $\beta=-0.22$ , 95% CI:  $-0.38$  to  $-0.06$ ) and VAT ( $\beta=-0.19$ , 95% CI:  $-0.25$  to  $-0.13$ ) (figure 4). In MVMR analysis, only vigorous PA remained protective against VAT ( $\beta=-0.15$ , 95% CI:  $-0.24$  to  $-0.07$ ), while no association was identified for %TBF ( $\beta=0.09$ , 95% CI:  $-0.11$  to  $0.29$ ) (figure 5). CAUSE similarly suggested a positive association between accelerometer-measured vigorous PA and VAT ( $\gamma=-0.14$ , 95% CI:  $-0.19$  to  $-0.08$ ) (table 1).



**Figure 3** Multivariable Mendelian randomisation IVW estimates for self-reported physical activity (PA) levels and body composition. Due to an insufficient number of overlapping SNPs for sedentary behaviour and LBM with educational attainment, no estimate was identified. %TBF, total body fat per cent; BMI, body mass index; IVW, inverse variance weighted; LBM, lean body mass; METs, metabolic equivalent of tasks; MVPA, moderate-to-vigorous PA; SNPs, single nucleotide polymorphisms; SSOE, strenuous sports or other exercises; VAT, visceral adipose tissue; WC, waist circumference; WHR, waist-to-hip ratio.

MR analyses with the self-reported SSOE measure demonstrated protective effects for all body composition measures except LBM (figure 2). In MVMR analyses, all SSOE–body composition associations were null except for VAT ( $\beta=-0.27$ , 95% CI:  $-0.52$  to  $-0.01$ ) (figure 3). CAUSE analyses indicated a negative association between SSOE and VAT ( $\gamma=-0.28$ , 95% CI:  $-0.46$  to  $-0.11$ ) (table 1). MR analyses of the overall activity phenotype (average accelerations) identified significant positive associations with %TBF ( $\beta=-0.31$ , 95% CI:  $-0.54$  to  $-0.08$ ) and VAT ( $\beta=-0.28$ , 95% CI:  $-0.49$  to  $-0.06$ ) (figure 4). MVMR and CAUSE did not support a significant protective effect for overall activity on VAT (table 1).

The F-statistics were greater than 20 for all genetic instruments suggesting instruments were sufficiently associated with the exposure (online supplemental table S2). We did not identify genetic variants in visual assessments of regressions, funnel plots or LOO which may invalidate the independence assumption. The MR-Egger test detected significant pleiotropy for self-reported

sedentary behaviour and %TBF, self-reported vigorous PA and WHR, and accelerometer-measured vigorous PA and LBM (online supplemental table S3). We did not identify other self-reported or accelerometer-measured associations which may have invalidated the exclusion criteria. However, heterogeneity estimates measured by Cochran's Q indicated widespread pleiotropy across self-reported and accelerometer-measured MR analyses (online supplemental table S4). An exception was noted in univariable MR analyses for self-reported PA with %TBF outcome, in which there was little evidence of heterogeneity. In MVMR analyses, there was a notably higher degree of heterogeneity.

MR-PRESSO sensitivity analyses identified some pleiotropic SNPs, which were removed and evaluated for significance after outlier correction (online supplemental table S5). Overall, MR-PRESSO estimates were consistent with our MVMR findings. Self-reported sedentary behaviour was positively associated with all body composition outcomes. Although no association was observed

**Table 1** Mendelian randomisation (MR) results for the putative effect of sedentary behaviour and physical activity on visceral adipose tissue


Exposure	Self-report			Accelerometer		
	SNPs (#)	Effect	95% CI	SNPs (#)	Effect	95% CI
Sedentary behaviour						
IVW	128	0.24 <sup>***</sup>	0.20 to 0.29	4	0.15	-0.01 to 0.31
MVMR		0.38 <sup>***</sup>	0.27 to 0.49		0.04	-0.14 to -0.23
MR-Egger		0.34 <sup>**</sup>	0.13 to 0.56		0.00	-0.69 to 0.69
Weighted mode		0.24 <sup>***</sup>	0.13 to 0.36		0.03	-0.17 to 0.24
Weighted median		0.24 <sup>***</sup>	0.20 to 0.28		0.07	-0.07 to 0.22
MR-PRESSO		0.30 <sup>***</sup>	0.21 to 0.38		0.12	-0.06 to 0.30
CAUSE		0.23 <sup>***</sup>	0.19 to 0.28		0.1	-0.01 to 0.21
Moderate PA						
IVW	32	0.08	-0.01 to 0.17		—	
MVMR		0.14	-0.07 to 0.36		—	
MR-Egger		0.08	-0.21 to 0.38		—	
Weighted mode		0.04	-0.13 to 0.21		—	
Weighted median		0.03	-0.05 to 0.12		—	
MR-PRESSO		0.13	-0.08 to 0.33		—	
CAUSE		0.25	0.02 to 0.47		—	
MVPA						
IVW	93	-0.09 <sup>**</sup>	-0.14 to -0.03	46	-0.14 <sup>***</sup>	-0.19 to -0.08
MVMR		0.05	-0.09 to 0.19		-0.12 <sup>**</sup>	-0.19 to -0.04
MR-Egger		-0.18	-0.4 to 0.03		-0.09	-0.24 to 0.06
Weighted mode		-0.04	-0.17 to 0.09		-0.03	-0.18 to 0.11
Weighted median		-0.07	-0.12 to -0.01		-0.09 <sup>**</sup>	-0.16 to -0.03
MR-PRESSO		-0.08	-0.21 to 0.05		0.00	-0.03 to 0.03
CAUSE		-0.06	-0.16 to 0.04		-0.11 <sup>***</sup>	-0.15 to -0.07
Vigorous PA						
IVW	61	-0.31 <sup>***</sup>	-0.31 to -0.13	26	-0.19 <sup>***</sup>	-0.25 to -0.13
MVMR		-0.02	-0.34 to 0.29		-0.15 <sup>***</sup>	-0.24 to -0.07
MR-Egger		0.05	-0.2 to 0.3		-0.24	-0.51 to 0.02
Weighted mode		-0.09	-0.34 to 0.15		-0.23	-0.39 to -0.06
Weighted median		-0.14 <sup>**</sup>	-0.24 to -0.04		-0.19 <sup>***</sup>	-0.28 to -0.1
MR-PRESSO		-0.12	-0.40 to 0.16		-0.09	-0.16 to -0.02
CAUSE		-0.13	-0.33 to 0.06		-0.14 <sup>***</sup>	-0.19 to -0.08
SSOE*/overall PA†						
IVW	98	-0.37 <sup>***</sup>	-0.47 to -0.27	6	-0.28	-0.49 to -0.06
MVMR		-0.27	-0.52 to -0.01		-0.21	-0.44 to 0.02
MR-Egger		-0.09	-0.45 to 0.26		-0.25	-1.08 to 0.57
Weighted mode		-0.37	-0.73 to 0.0		-0.17	-0.39 to 0.06
Weighted median		-0.31 <sup>***</sup>	-0.41 to -0.21		-0.21	-0.38 to -0.03
MR-PRESSO		-0.33 <sup>**</sup>	-0.55 to -0.11		-0.05	-0.16 to 0.05
CAUSE		-0.28	-0.46 to -0.11		-0.08	-0.19 to 0.03

\*\*\*P<0.001, \*\*p<0.01.

\*Analyses only conducted for self-report PA.

†Analyses only conducted for accelerometer-measured PA.

.CAUSE, causal analysis using summary effect; IVW, inverse variance weighted; MVMR, multivariable MR; MVPA, moderate-to-vigorous PA; PA, physical activity; PRESSO, pleiotropy residual sum and outlier; SNPs, single nucleotide polymorphisms; SSOE, strenuous sports and other exercises.



**Figure 4** Univariable Mendelian randomisation IVW estimates for accelerometer-measured physical activity (PA) levels and body composition. %TBF, total body fat per cent; BMI, body mass index; IVW, inverse variance weighted; LBM, lean body mass; METs, metabolic equivalent of tasks; MVPA, moderate-to-vigorous PA; SNPs, single nucleotide polymorphisms; VAT, visceral adipose tissue; WC, waist circumference; WHR, waist-to-hip ratio.


for any self-reported vigorous PA exposure, negative associations were identified in accelerometer-measured vigorous PA for %TBF and VAT. Further, we identified robust associations of SSOE with all body composition measures, including VAT.

Other sensitivity analyses including median and mode-based estimates were typically consistent with our main findings (online supplemental tables S6 and S7), while MR-Egger estimates trended towards the null (online supplemental table S8). CAUSE suggested robust protective effects for accelerometer-measured MVPA and vigorous PA on VAT, with limited evidence indicating a protective effect of SSOE on VAT (online supplemental table S9). CAUSE additionally suggested a positive association of self-reported sedentary behaviour with BMI, WHR, WC and VAT, while no effects were identified using the accelerometer-measured sedentary phenotype (online supplemental table S9). Reversing our phenotypes did not reveal consistent associations between body composition and PA outcomes (online supplemental table S10). Sensitivity analyses excluding the *APOE* rs429358 variant attenuated multiple associations, including the protective effect of accelerometer-measured MVPA on VAT. However, accelerometer-measured vigorous PA remained protective for VAT (online supplemental figures S3 and S4), and lower-intensity levels of PA continued to show no association with anthropometric or body composition measures. We did not identify directional changes for

any associations after Steiger filtering sensitivity analyses. Across self-report and accelerometer PA exposures sensitivity analyses, including CAUSE and MVMR, our results suggest an association between genetically predicted higher-intensity PA and lower VAT (table 1).

## DISCUSSION

Ambiguity currently exists in our understanding of whether PA protects against obesity and/or central adiposity, which has been driven by fundamental limitations to both observational studies and intervention trials.<sup>8 9 23</sup> The relationship between PA and body composition traits is highly complex, and genetic variants underlying each of these have been shown to have pleiotropic effects, including with each other (eg, *CADM2* associated with both BMI and PA). Therefore, efforts are needed to better elucidate the shared genetics and pathways that link PA with weight and body composition.<sup>26</sup> We applied MR using a comprehensive set of PA and body composition measures to increase confidence in causal estimations between PA and body composition, considering self-reported and accelerometer-based PA exposures in parallel analyses. Our results provide some evidence of a protective effect on VAT from genetically predicted levels of accelerometer-measured MVPA and vigorous PA, as well as self-reported SSOE. In contrast to these findings with VAT, we find limited evidence of a



**Figure 5** Multivariable Mendelian randomisation IVW estimates for accelerometer-measured physical activity (PA) levels and body composition. Due to an insufficient number of overlapping SNPs for sedentary behaviour and LBM with educational attainment, no estimate was identified. %TBF, total body fat per cent; BMI, body mass index; IVW, inverse variance weighted; LBM, lean body mass; METs, metabolic equivalent of tasks; MVPA, moderate-to-vigorous PA; SNPs, single nucleotide polymorphisms; VAT, visceral adipose tissue; WC, waist circumference; WHR, waist-to-hip ratio.

causal relationship between genetically predicted PA and other body composition indices.

MR estimates were not always consistent across self-reported and accelerometer-based PA measures. Self-reported measures may not capture some aspects of activity or sedentary behaviour, while accelerometer measurements were designed to record uninterrupted activity.<sup>53</sup> Further, accelerometer measurements reflected activity from one specific week and may not capture some forms of PA such as resistance training.<sup>36</sup> Overall, each of these methods likely suffers from both shared and unique biases. These two types of measures can thus be thought of as complementary, and evidence supported by both measures may be viewed as stronger.

MR and MVMR analyses suggested consistent protective associations for accelerometer-measured MVPA, vigorous PA and self-reported SSOE with VAT. Sensitivity analyses with MR-PRESSO, median and mode-based estimates were consistent with the direction of IVW estimates despite the reduced sensitivity of these methods. CAUSE estimates were marginally attenuated compared with other methods, but similarly suggested a protective association from higher-intensity PA. Although we detected a strong positive association between self-reported sedentary behaviour and VAT, acceleration-measured sedentary exposure did not reveal any association with VAT. Although the directional effect was stable between

self-report and accelerometer measurements, we find limited evidence suggesting a consistent positive association between sedentary behaviour and VAT. Further, we did not observe any consistent protective associations between moderate PA and VAT. Overall, these associations suggest an independent effect of high-intensity PA on VAT.

Our results align with previous evidence suggesting an independent protective effect of PA on VAT. The overproportional reduction of VAT compared with subcutaneous tissues may be explained by heterogeneous lipolysis regulation, which has been shown to be dependent on the location of adipose tissue in the body.<sup>54</sup> Increasing levels of PA intensity and duration have been shown to increase secretion of catecholamines.<sup>55</sup> In contrast to subcutaneous tissue types, higher concentrations of  $\beta$ -adrenoceptors and  $\alpha$ 2A-adrenoceptors in VAT enhance lipolysis through binding of hormones including catecholamines, thereby upregulating cyclic AMP production and increasing hydrolysis of triglycerides.<sup>56</sup> Other physiological features such as reduced insulin receptor substrate-1 in VAT limit antilipolytic processes by lowering insulin sensitivity in this region compared with subcutaneous tissue.<sup>56</sup> Increased blood flow and higher concentration of cells in the abdominal cavity may further amplify regional lipolytic activity. In a previous intervention by Johnson *et al*,<sup>57</sup> aerobic exercise was shown to reduce



visceral adiposity without change to BMI or %TBF. A systematic review of randomised PA control trials further identified a small number of studies where VAT was reduced independently from body mass changes.<sup>58</sup> Here, we find evidence suggesting that PA at sufficient levels of intensity may independently reduce VAT and improve an individual's metabolic profile.

We did not identify consistent associations between genetically predicted PA exposures and other body composition outcomes. Although MR analyses indicated positive associations of sedentary behaviour with multiple body composition measures, no positive associations were identified for body composition measures (with the exception of VAT) after controlling for educational attainment. Similarly, MR estimates of accelerometer-measured sedentary behaviour were null. MR estimates of higher-intensity MET-based measures, including MVPA and vigorous PA, revealed few protective effects for anthropometric measures. While both accelerometer-measured MVPA and vigorous PA demonstrated negative associations with %TBF, these effects were again null after controlling for educational attainment. MVMR estimates for exposures not based on METs, including SSOE and overall activity, did not reach significance. Sensitivity analyses using MR-PRESSO, median and mode-based estimates were similarly inconsistent or null across sedentary behaviour, moderate PA, MVPA and vigorous PA exposures. In contrast to other null body composition findings, we identified a positive association with genetically predicted self-reported vigorous PA and BMI, and accelerometer-measured PA and LBM in MVMR analysis. These results may suggest that PA has a direct positive effect on BMI and LBM. We interpret the robustness of these findings with caution, however, due to the null effects in parallel analyses and other sensitivity tests.

Our findings partially contrast with MR analyses conducted by Carrasquilla *et al*, who identified a protective effect of accelerometer-measured vigorous PA on BMI, and a bidirectional positive association between accelerometer-measured sedentary time and BMI using the IVW and CAUSE methods.<sup>28</sup> Our IVW estimates similarly suggested a positive association for self-reported and accelerometer-measured sedentary behaviour with BMI, however, our MVMR analyses and accelerometer-based CAUSE analyses did not identify positive associations. Further, we did not identify any protective association for vigorous PA with BMI. These differences may have been a result of our application of two-sample MR across analyses, differing instrument selection thresholds, and/or the use of different priors in CAUSE analyses. Similar to Carrasquilla *et al*, a recent preprint from den Hoed *et al*<sup>59</sup> reported a positive association for sedentary time with both BMI and %TBF in CAUSE analyses. Although we also report a positive association between self-reported sedentary behaviour and BMI, we did not identify an association with %TBF. This discrepancy may have been a result of the use of leisure screen time as the proxy sedentary behaviour used by den Hoed *et al*, which reflects a

narrower phenotype compared with our combined self-reported sedentary measure. Like den Hoed *et al*, we did not identify consistent protective associations of MVPA with body composition measures.

A previous MR meta-analysis from Shnurr *et al*<sup>26</sup> using a smaller cohort of both children and adults has further supported a positive association of BMI with sedentary time. Two previous one-sample MR studies among children have similarly identified a positive association between higher BMI and sedentary time.<sup>60 61</sup> In our bidirectional analyses, we identified a positive association for BMI with self-reported sedentary behaviour, while estimates of BMI on accelerometer-measured sedentary behaviour were null. Similar to Shnurr *et al*,<sup>26</sup> we did not identify an effect of genetically predicted BMI on overall activity or MVPA. A previous MR analysis from Doherty *et al*<sup>27</sup> additionally reported an inverse bidirectional relationship between BMI and accelerometer-measured overall activity. However, we did not identify a protective association for PA measures with BMI after controlling for educational attainment in MVMR analyses. Further, bidirectional IVW estimates in our study did not suggest a causal association of BMI, %TBF or LBM with PA measures. In contrast to our study, Doherty *et al* used an accelerometer-measured overall activity exposure distinct from our own exposure, as well as a likelihood-based method to estimate causal effects. Overall, our findings conflicting with previous literature may have been driven by our use of a two-sample MR design, differing instrumental variable inclusion criteria, as well as our application of more conservative methodologies such as CAUSE and MVMR.

Health agencies uniformly emphasise the importance of moderate PA for weight management.<sup>62 63</sup> Some evidence suggests, however, that these PA recommendations are insufficient.<sup>64</sup> While there is little doubt that PA offers immeasurable benefits to overall health, our results suggest a limited protective effect for anthropometric indices, LBM or %TBF from PA. Our null findings may be plausible under the recently proposed constrained total energy expenditure (TEE) model.<sup>65</sup> In contrast to conventional additive models, the constrained TEE model accounts for potential downregulation of non-musculoskeletal physiological activity during higher levels of PA, resulting in TEE within a homeostatically constrained range.<sup>66</sup> This model may provide an explanatory framework for the lack of associations observed between PA and most body composition measures in our analyses.

### Limitations

Our study includes several critical limitations. First, we were unable to identify SNPs at the genome-wide significance level in GWAS and used a reduced instrumental variable threshold ( $p < 5 \times 10^{-6}$ ) across our analyses. The lower predictive capacity of these SNPs may have limited our ability to detect significant sedentary and PA effects on body composition. Despite this limitation, all

instrumental variables indicated sufficient strength, as measured by the F-statistic. Additionally, while we applied two-sample MR to limit weak instrument bias, stratification of VAT into samples not overlapping with sedentary behaviour or PA may have introduced a form of selection bias. The inherent assumptions of MR may be violated when evaluating polygenic risk factors, such as PA, as the influence of genetic variants on such exposures is unlikely to be specific. We therefore applied MVMR to estimate the direct causal effect of sedentary and PA exposures, which is valuable when instruments are suspected to violate the second and third instrumental variable assumptions.<sup>49</sup> The lower number of instruments identified for accelerometer-measured exposures may have reduced the ability to detect significant associations in MVMR; however, we note that null findings were consistent with CAUSE analyses.

A key design limitation was our inability to evaluate light PA exposures in self-reported or accelerometer-based measurements. Despite the large consortia used in this investigation, we were unable to perform sex-stratified analysis due to limited instrument availability. Our results may have been further attenuated by effect modification, as both regional adiposity<sup>67</sup> and PA<sup>37</sup> have been shown to differ between sexes. We acknowledge a potential increase in the type I error rate in analyses on account of no correction for multiple testing. Here we report and consider each association individually.<sup>68</sup> Importantly, we do not rely on a single estimate for our inferences, but more broadly consider the consistency of associations across measures and methods to draw conclusions. We emphasise that MR does not provide an infallible test of causation and encourage replication in future studies.

In conclusion, MR and MVMR analyses of various PA levels did not demonstrate consistent protective effects on body composition measures. Our null findings reaffirm the equivocal effectiveness of weight management programmes relying on PA alone, in contrast to programmes using both dietary restriction and PA. We do find evidence of an association between genetically predicted high-intensity PA and lower VAT. At the time of this work, no peer-reviewed predictive models for subcutaneous adipose tissue, ectopic fat or hepatic fat were identified to evaluate as an outcome with PA. MR analyses of these and other more detailed body composition outcomes may provide greater insight into specific cardiometabolic benefits of PA.

#### Author affiliations

<sup>1</sup>Department of Epidemiology and Biostatistics, The University of Arizona, Tucson, Arizona, USA

<sup>2</sup>Department of Medicine, The University of Arizona, Tucson, Arizona, USA

<sup>3</sup>Department of Nutritional Sciences, University of Arizona, Tucson, Arizona, USA

<sup>4</sup>Department of Health Promotion Sciences, University of Arizona, Tucson, Arizona, USA

<sup>5</sup>Milken Institute of Public Health, The George Washington University, Washington, District of Columbia, USA

<sup>6</sup>Department of Biological Sciences, University of Southern California, Los Angeles, California, USA

**Twitter** Ferris A Ramadan @ferris\_ram

**Acknowledgements** We would like to recognise the UK Biobank, and the GIANT and GEFOS consortia for their incredible efforts in developing and curating the data sources used in this study.

**Contributors** FAR and YCK conceived the project. FAR processed the experimental data, performed the analyses and designed the figures. YCK, DAR and JWB contributed to the interpretation of the results. YCK and RAC helped guide replication of visceral adipose tissue models. FAR and YCK led manuscript writing. All authors provided critical feedback and helped shape the research, analysis and manuscript. YCK supervised and is the guarantor of the project.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Ethics approval** This study involves human participants and ethical approval of the UKB was provided by the Northwest Multicentre Research Ethics Committee, the National Information Governance Board for Health & Social Care, and the Community Health Index Advisory Group (IRAS project ID: 299116). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be publicly available from a third party. Summary statistics from the GENetic Factors for Osteoporosis (GEFOS) consortium, the Genetic Investigation of ANthropometric Traits (GIANT) consortium, self-reported sedentary behaviour and total body fat per cent are publicly available. Summary statistics for physical activity, educational attainment and visceral adipose tissue were obtained from the UK Biobank and are not publicly available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Ferris A Ramadan <http://orcid.org/0000-0003-2039-5241>

#### REFERENCES

- Meldrum DR, Morris MA, Gambone JC. Obesity pandemic: causes, consequences, and solutions-but do we have the will? *Fertil Steril* 2017;107:833–9.
- Townshend T, Lake A. Obesogenic environments: current evidence of the built and food environments. *Perspect Public Health* 2017;137:38–44.
- Powell KE, King AC, Buchner DM, et al. The scientific foundation for the physical activity guidelines for Americans, 2nd edition. *J Phys Act Health* 2018;1–11.
- Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019;15:288–98.
- Popkin B, Monteiro C, Swinburn B. Overview: Bellagio conference on program and policy options for preventing obesity in the low- and middle-income countries. *Obes Rev* 2013;14 Suppl 2:1–8.
- Roberto CA, Swinburn B, Hawkes C, et al. Patchy progress on obesity prevention: emerging examples, entrenched barriers, and new thinking. *Lancet* 2015;385:2400–9.
- Franz MJ, VanWormer JJ, Crain AL, et al. Weight-Loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc* 2007;107:1755–67.

- 8 Swift DL, McGee JE, Earnest CP, *et al.* The effects of exercise and physical activity on weight loss and maintenance. *Prog Cardiovasc Dis* 2018;61:206–13.
- 9 Chin S-H, Kahathuduwa CN, Binks M. Physical activity and obesity: what we know and what we need to know. *Obes Rev* 2016;17:1226–44.
- 10 Peterson MD, Sen A, Gordon PM. Influence of resistance exercise on lean body mass in aging adults: a meta-analysis. *Med Sci Sports Exerc* 2011;43:249–58.
- 11 Kelley GA, Kelley KS, Tran ZV. Exercise and bone mineral density in men: a meta-analysis. *J Appl Physiol* 2000;88:1730–6.
- 12 Jakicic JM, Powell KE, Campbell WW, *et al.* Physical activity and the prevention of weight gain in adults: a systematic review. *Med Sci Sports Exerc* 2019;51:1262–9.
- 13 Johnson NA, Sultana RN, Brown WJ, *et al.* Physical activity in the management of obesity in adults: a position statement from exercise and sport science Australia. *J Sci Med Sport* 2021;24:1245–54.
- 14 Slentz CA, Aiken LB, Houmard JA, *et al.* Inactivity, exercise, and visceral fat. STRRIDE: a randomized, controlled study of exercise intensity and amount. *J Appl Physiol* 2005;99:1613–8.
- 15 Slentz CA, Duscha BD, Johnson JL, *et al.* Effects of the amount of exercise on body weight, body composition, and measures of central obesity: STRRIDE—a randomized controlled study. *Arch Intern Med* 2004;164:31–9.
- 16 Hamdy O, Porramatikul S, Al-Ozairi E. Metabolic obesity: the paradox between visceral and subcutaneous fat. *Curr Diabetes Rev* 2006;2:367–73.
- 17 Vissers D, Hens W, Taeymans J, *et al.* The effect of exercise on visceral adipose tissue in overweight adults: a systematic review and meta-analysis. *PLoS One* 2013;8:e56415.
- 18 Verheggen RJHM, Maessen MFH, Green DJ, *et al.* A systematic review and meta-analysis on the effects of exercise training versus hypocaloric diet: distinct effects on body weight and visceral adipose tissue. *Obes Rev* 2016;17:664–90.
- 19 Whitaker KM, Pereira MA, Jacobs DR, *et al.* Sedentary behavior, physical activity, and abdominal adipose tissue deposition. *Med Sci Sports Exerc* 2017;49:450–8.
- 20 Murabito JM, Pedley A, Massaro JM, Alison P, Massaro Joseph M, *et al.* Moderate-to-vigorous physical activity with accelerometry is associated with visceral adipose tissue in adults. *J Am Heart Assoc* 2015;4:e001379.
- 21 Maillard F, Pereira B, Boisseau N. Effect of high-intensity interval training on total, abdominal and visceral fat mass: a meta-analysis. *Sports Med* 2018;48:269–88.
- 22 Lawlor DA, Harbord RM, Sterne JAC, *et al.* Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Stat Med* 2008;27:1133–63.
- 23 Smith GD, Ebrahim S. Mendelian Randomization: Genetic Variants as Instruments for Strengthening Causal Inference in Observational Studies [Internet]. *Biosocial Surveys. National Academies Press* 2008 <https://www.ncbi.nlm.nih.gov/books/NBK62433/> (cited 2020 Jun 17).
- 24 Wareham NJ, van Sluijs EMF, Ekelund U. Physical activity and obesity prevention: a review of the current evidence. *Proc Nutr Soc* 2005;64:229–47.
- 25 Ekelund U, Brage S, Besson H, *et al.* Time spent being sedentary and weight gain in healthy adults: reverse or bidirectional causality? *Am J Clin Nutr* 2008;88:612–7.
- 26 Schnurr TM, Stallknecht BM, Sørensen TIA, *et al.* Evidence for shared genetics between physical activity, sedentary behaviour and adiposity-related traits. *Obes Rev* 2021;22:e13182 <https://onlinelibrary.wiley.com/doi/>
- 27 Doherty A, Smith-Byrne K, Ferreira T, *et al.* Gwas identifies 14 loci for device-measured physical activity and sleep duration. *Nat Commun* 2018;9:5257.
- 28 Carrasquilla GD, Garcia-Ureña M, Fall T, *et al.* Mendelian randomization suggests a bidirectional, causal relationship between physical inactivity and obesity [Internet]. *Genetics* 2021 <http://biorxiv.org/lookup/doi/> (cited 2021 Aug 5).
- 29 Klimentidis YC, Raichlen DA, Bea J, *et al.* Genome-Wide association study of habitual physical activity in over 377,000 UK Biobank participants identifies multiple variants including CADM2 and APOE. *Int J Obes* 2018;42:1161–76.
- 30 Skrivankova VW, Richmond RC, Woolf BAR, *et al.* Strengthening the reporting of observational studies in epidemiology using Mendelian randomisation (STROBE-MR): explanation and elaboration. *BMJ* 2021;375:n2233.
- 31 Zillikens MC, Demissie S, Hsu Y-H, *et al.* Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. *Nat Commun* 2017;8:80.
- 32 GIANT consortium - Giant Consortium [Internet]. [cited 2021 Jan 7]. Available: [https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT\\_consorrtium](https://portals.broadinstitute.org/collaboration/giant/index.php/GIANT_consorrtium)
- 33 van de Vegte YJ, Said MA, Rienstra M, *et al.* Genome-Wide association studies and Mendelian randomization analyses for leisure sedentary behaviours. *Nat Commun* 2020;11:1770.
- 34 Prince SA, LeBlanc AG, Colley RC, *et al.* Measurement of sedentary behaviour in population health surveys: a review and recommendations. *PeerJ* 2017;5:e4130.
- 35 Committee IR. Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ)—short and long forms, 2005. Available: <http://www.ipaq.ki.se/scoring.pdf>
- 36 Cleland C, Ferguson S, Ellis G, *et al.* Validity of the International physical activity questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol* 2018 ;;18:176. Dec 22.
- 37 Doherty A, Jackson D, Hamerla N, *et al.* Large scale population assessment of physical activity using wrist worn Accelerometers: the UK Biobank study. *PLoS One* 2017;12:e0169649.
- 38 Hildebrand M, VAN Hees VT, Hansen BH, *et al.* Age group comparability of raw accelerometer output from wrist- and hip-worn monitors. *Med Sci Sports Exerc* 2014;46:1816–24.
- 39 Loh P-R, Tucker G, Bulik-Sullivan BK, *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat Genet* 2015;47:284–90.
- 40 Okbay A, Beauchamp JP, Fontana MA, *et al.* Genome-Wide association study identifies 74 loci associated with educational attainment. *Nature* 2016;533:539–42.
- 41 Droomers M, Schrijvers CT, Mackenbach JP. Educational level and decreases in leisure time physical activity: predictors from the longitudinal globe study. *J Epidemiol Community Health* 2001;55:562–8.
- 42 Zillikens MC, Demissie S, Hsu Y-H, *et al.* Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. *Nat Commun* 2017;8:80.
- 43 Achamrah N, Colange G, Delay J, *et al.* Comparison of body composition assessment by DXA and BIA according to the body mass index: a retrospective study on 3655 measures. *PLoS One* 2018;13:e0200465.
- 44 Lu Y, Day FR, Gustafsson S, *et al.* New loci for body fat percentage reveal link between adiposity and cardiometabolic disease risk. *Nat Commun* 2016;7:10495.
- 45 Karlsson T, Rask-Andersen M, Pan G, *et al.* Contribution of genetics to visceral adiposity and its relation to cardiovascular and metabolic disease. *Nat Med* 2019;25:1390–5.
- 46 Davies NM, Holmes MV, Davey Smith G, Smith GD. Reading Mendelian randomisation studies: a guide, glossary, and checklist for clinicians. *BMJ* 2018;362:k601.
- 47 Verbanck M, Chen C-Y, Neale B, *et al.* Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018;50:693–8.
- 48 Morrison J, Knoblauch N, Marcus JH, *et al.* Mendelian randomization accounting for correlated and uncorrelated pleiotropic effects using genome-wide summary statistics. *Nat Genet* 2020;52:740–7.
- 49 Sanderson E, Davey Smith G, Windmeijer F, *et al.* An examination of multivariable Mendelian randomization in the single-sample and two-sample summary data settings. *Int J Epidemiol* 2019;48:713–27.
- 50 Yavorska OO, Burgess S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. *Int J Epidemiol* 2017;46:1734–9.
- 51 Sanderson E, Spiller W, Bowden J. Testing and Correcting for Weak and Pleiotropic Instruments in Two-Sample Multivariable Mendelian Randomisation [Internet]. *Genetics*; 2020 <http://biorxiv.org/lookup/doi/> (cited 2021 Mar 22).
- 52 R: The R Project for Statistical Computing [Internet]. [cited 2020 Nov 2]. Available: <https://www.r-project.org/>
- 53 Prince SA, Adamo KB, Hamel ME, *et al.* A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008;5:56.
- 54 Wajchenberg BL. Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev* 2000;21:697–738.
- 55 Zouhal H, Jacob C, Delamarche P, *et al.* Catecholamines and the effects of exercise, training and gender. *Sports Med* 2008;38:401–23.
- 56 Arner P. Human fat cell lipolysis: biochemistry, regulation and clinical role. *Best Pract Res Clin Endocrinol Metab* 2005;19:471–82.
- 57 Johnson NA, Sachinwalla T, Walton DW, *et al.* Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Hepatology* 2009;50:1105–12.

- 58 Kay SJ, Fiatarone Singh MA, Singh MAF. The influence of physical activity on abdominal fat: a systematic review of the literature. *Obes Rev* 2006;7:183–200.
- 59 den Hoed M, Wang Z, Emmerich A, *et al*. Physical activity and sedentary behavior; mechanistic insights and role in disease prevention 2021.
- 60 Richmond RC, Davey Smith G, Ness AR, *et al*. Assessing causality in the association between child adiposity and physical activity levels: a Mendelian randomization analysis. *PLoS Med* 2014;11:e1001618.
- 61 Schnurr TM, Viitasalo A, Eloranta A-M, *et al*. Genetic predisposition to adiposity is associated with increased objectively assessed sedentary time in young children. *Int J Obes* 2018;42:111–4.
- 62 WHO. Physical activity [Internet]. World Health Organization, 2020. Available: <https://www.who.int/news-room/fact-sheets/detail/physical-activity> [Accessed cited 2021 May 9].
- 63 Choosing a Safe and Successful Weight-loss Program | NIDDK [Internet]. National Institute of Diabetes and Digestive and Kidney Diseases. [cited 2021 Mar 19]. Available: <https://www.niddk.nih.gov/health-information/weight-management/choosing-a-safe-successful-weight-loss-program>
- 64 Lee I-M, Djoussé L, Sesso HD, *et al*. Physical activity and weight gain prevention. *JAMA* 2010;303:1173–9.
- 65 Pontzer H. Constrained total energy expenditure and the evolutionary biology of energy balance. *Exerc Sport Sci Rev* 2015;43:110–6.
- 66 Pontzer H. Energy constraint as a novel mechanism linking exercise and health. *Physiology* 2018;33:384–93.
- 67 Smith HA, Storti KL, Arena VC, *et al*. Associations between accelerometer-derived physical activity and regional adiposity in young men and women. *Obesity* 2013;21:1299–305.
- 68 Perneger TV. What's wrong with Bonferroni adjustments. *BMJ* 1998;316:1236–8.

## Supplemental Tables and Figures

<b>Table S1.</b> Participant Characteristics Included in the Genome-Wide Association Studies.....	2
<b>Table S2.</b> F Statistics for Mendelian Randomization Analyses for Physical Activity to Body Composition.....	3
<b>Table S3.</b> MR-EGGER Intercept Estimates for Physical Activity to Body Composition.....	4
<b>Table S4.</b> Heterogeneity estimates Mendelian Randomization Analyses for Physical Activity to Body Composition.....	5
<b>Table S5.</b> MR-PRESSO Estimates for Physical Activity to Body Composition .....	6
<b>Table S6.</b> MR Median-based Estimates for Physical Activity to Body Composition.....	7
<b>Table S7.</b> MR Mode-based Estimates for Physical Activity to Body Composition.....	8
<b>Table S8.</b> MR-EGGER Estimates for Physical Activity to Body Composition.....	9
<b>Table S9.</b> CAUSE estimates for Physical Activity to Body Composition .....	10
<b>Table S10.</b> Bidirectional MR IVW Estimates for Body Composition to Physical Activity .....	11
<b>Table S11.</b> SNP Exposure and Outcome Associations .....	Online Supplemental
<b>Figure S1.</b> Manhattan Plot and genome-wide significant SNPs for self-reported moderate PA .....	142
<b>Figure S2.</b> Manhattan Plot and genome-wide significant SNPs for accelerometer-measured MVPA .....	153
<b>Figure S3.</b> Univariable Mendelian Randomization IVW Estimates for Self-Reported Physical Activity Levels and Body Composition removing the <i>APOE rs429358</i> variant .....	<b>Error! Bookmark not defined.</b> 4
<b>Figure S4.</b> Univariable Mendelian Randomization IVW Estimates for Accelerometer-Measured Physical Activity Levels and Body Composition removing the <i>APOE rs429358</i> variant .....	15

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

2

**Table S1.** Participant Characteristics Included in the Genome-Wide Association Studies

Phenotype	Study	n	Women (%)	Age, Years (mean ± SD)	Trait Value (mean ± SD)
<b>Self-report (hr./day)</b>					
Sedentary Behavior	UKB / van de Vegte et al.	422,218	54.3	57.4 ± 8.0	4.7 ± 2.7
Moderate PA	UKB / Novel	360,912	53.4	56.7 ± 8.1	1.2 ± 1.3
MVPA	UKB / Klimentidis et al.	377,234	50.6	56.1 ± 8.2	1.9 ± 1.8
Vigorous PA (≥3 vs. 0 dys./wk.) <sup>†</sup>	UKB / Klimentidis et al.	98,060; 162,995	47.3; 58.1	55.4 ± 8.3; 57.4 ± 7.8	1.1 ± 0.9; 0.0 ± 0.0
SSOE (>2-3 vs 0 dys./wk.) <sup>†</sup>	UKB / Klimentidis et al.	124,842; 225,650	38.1; 35.3	52.7 ± 8.1; 52.6 ± 7.9	1.1 ± 0.7; 0.0 ± 0.0
<b>Accelerometer</b>					
Sedentary Behavior <sup>§</sup>	UKB / Doherty et al.	91,105	--	--	--
MVPA <sup>‡</sup>	UKB / Novel	97,737	56.1	56.2 ± 7.8	0.0799 ± 0.0384
Vigorous PA <sup>‡</sup>	UKB / Klimentidis et al.	90,667	56.1	56.2 ± 7.8	0.0026 ± 0.0033
Overall activity ( <i>milli-gravities</i> )	UKB / Klimentidis et al.	91,105	56.1	56.2 ± 7.8	27.1 ± 27.0
<b>Body Comp.</b>					
BMI ( <i>kg/m<sup>2</sup></i> ) <sup>††</sup>	GIANT / Locke et al.	249,796	--	--	--
WHR <sup>††</sup>	GIANT/ Shungin et al.	224,459	--	--	--
WC ( <i>cm</i> ) <sup>††</sup>	GIANT/ Shungin et al.	224,459	--	--	--
LBM ( <i>kg</i> ) <sup>††</sup>	GEFOS / Zillikens et al.	38,292	--	--	--
%TBF	Lu et al.	89,297	48.3	57.8 ± 9.4	31.6 ± 6.7
VAT (Self-Report PA; <i>kg</i> )	UKB / Karlsson et al.	56,908	56.7	57.1 ± 7.8	1.5 ± .86
VAT (Accel. PA; <i>kg</i> )	UKB / Karlsson et al.	323,769	53.8	56.9 ± 8.1	1.4 ± .83
<b>Covariate</b>					
Educational Attainment	UKB / Okbay et al.	293,723	55.1	57.5 ± 8.0	14.3 ± 3.6

Abbreviations: PA, Physical Activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; BMI, body mass index; WC, waist circumference; WHR, waist-to-hip ratio; LBM, lean body mass; %TBF, total body fat percent; VAT, visceral adipose tissue; UKB, UK Biobank; GIANT, Genetic Investigation of ANthropometric Traits consortium; GEFOS, GENetic Factors for Osteoporosis consortium.

<sup>†</sup>Phenotype was evaluated as a binary variable.

<sup>§</sup>Phenotype was trained on 153 free-living individuals to predict metabolic equivalent of tasks.

<sup>‡</sup>Phenotype was derived from fraction accelerations based on metabolic equivalent of tasks.

<sup>††</sup>Consortium demographics unavailable.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

3

**Table S2.** F Statistics for Mendelian Randomization Analyses for Physical Activity to Body Composition

Phenotypes		Self-report MR			Accelerometer MR		
Activity	Body Comp.	Mean F	Median F	Range F	Mean F	Median F	Range F
Sedentary Behavior	LBM	45.75	38.97	30.41, 144.19	27.61	27.61	27.26, 27.95
	BMI	45.32	38.55	30.41-144.19	25.34	25.2	23.01, 27.95
	WHR	45.3	38.48	30.41-144.19	25.34	25.2	23.01, 27.95
	WC	45.23	38.62	30.41-144.19	25.34	25.2	23.01, 27.95
	%TBF	45.77	38.97	30.41-144.19	26.07	27.26	23.01, 27.95
	VAT	42.71	37.3	30.41-144.19	26.4	26.69	23.09, 30.18
Moderate PA	LBM	23.40	23.39	21.52, 26.05	--	--	--
	BMI	23.41	23.37	21.12-30.38	--	--	--
	WHR	23.39	23.13	21.12-30.38	--	--	--
	WC	23.39	23.13	21.12-30.38	--	--	--
	%TBF	23.51	23.37	21.12-30.38	--	--	--
	VAT	23.09	22.77	21.02-30.38	--	--	--
MVPA	LBM	25.85	24.16	20.97, 45.99	25.13	23.32	21.62, 33.12
	BMI	25.94	24.41	20.88-45.99	24.83	23.4	20.91, 34.67
	WHR	25.94	24.41	20.88-45.99	24.83	23.4	20.91, 34.67
	WC	25.9	24.39	20.88-45.99	24.83	23.4	20.91, 34.67
	%TBF	26.02	24.39	20.88-45.99	24.68	23.4	20.91, 34.67
	VAT	25.75	23.9	20.86-45.99	24.06	23.24	20.92, 34.72
Vigorous PA	LBM	27.78	23.05	21.18, 55.26	23.56	22.58	20.88, 26.64
	BMI	26.6	23.48	20.98-55.26	24.98	24.56	20.88, 36.07
	WHR	26.73	23.54	20.98-55.26	24.98	24.56	20.88, 36.07
	WC	26.47	23.42	20.98-55.26	24.98	24.56	20.88, 36.07
	%TBF	27.07	23.54	20.98-55.26	24.98	24.56	20.88, 36.07
	VAT	25.62	23.05	20.9-55.26	24.51	23.03	20.88, 36.76
SSOE	LBM	25.40	24.48	20.98, 37.34	--	--	--
	BMI	26.03	24.52	20.98-75.4	--	--	--
	WHR	25.99	24.65	20.98-75.4	--	--	--
	WC	26.09	24.59	20.98-75.4	--	--	--
	%TBF	26.17	24.65	20.98-75.4	--	--	--
	VAT	25.94	24.48	20.89-82.57	--	--	--
Overall Activity	LBM	--	--	--	25.16	24.84	22.66, 27.98
	BMI	--	--	--	24.54	23.76	22.66, 27.98
	WHR	--	--	--	24.54	23.76	22.66, 27.98
	WC	--	--	--	24.54	23.76	22.66, 27.98
	%TBF	--	--	--	24.54	23.76	22.66, 27.98
	VAT	--	--	--	24.45	24.84	21.28, 27.98

Abbreviations: MR, Mendelian randomization; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: An F statistic of 10 indicates that the bias of the IV estimator is 10% of the bias of the observational estimator.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

4

**Table S3.** MR-EGGER Intercept Estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR		Accelerometer MR	
Activity	Body Comp.	MR-EGGER Intercept	<i>p</i>	MR-EGGER Intercept	<i>p</i>
Sedentary Behavior	LBM	0.025031	0.2958	--	--
	BMI	-0.00449	0.3034	0.017358	0.64
	WHR	-0.00209	0.5284	0.00348	0.94
	WC	-0.00449	0.2279	0.002724	0.96
	%TBF	-0.01355	0.0073	0.052934	0.39
Moderate PA	VAT	-0.00168	0.586	0.005832	0.59
	LBM	0.037154	0.5654	--	--
	BMI	0.006266	0.3227	--	--
	WHR	0.001847	0.7752	--	--
	WC	0.0043	0.4634	--	--
MVPA	%TBF	-0.00217	0.7637	--	--
	VAT	-0.00245	0.6182	--	--
	LBM	0.035138	0.34	0.035717	0.513
	BMI	0.001909	0.5612	-0.00662	0.36
	WHR	-0.00197	0.5577	-0.00404	0.47
Vigorous PA	WC	0.002466	0.496	-0.00533	0.44
	%TBF	0.000303	0.9364	-0.00891	0.18
	VAT	0.004179	0.2116	-0.00181	0.39
	LBM	0.013209	0.753	-0.41894	0.047
	BMI	-0.0054	0.216	0.004192	0.74
SSOE	WHR	-0.00829	0.0331	0.004787	0.7
	WC	-0.0051	0.2259	-0.00019	0.99
	%TBF	-0.00134	0.7813	0.026992	0.13
	VAT	-0.00144	0.7043	-0.00095	0.8
	LBM	-0.00619	0.8902	--	--
Overall Activity	BMI	-0.00428	0.3557	--	--
	WHR	-0.00214	0.549	--	--
	WC	-0.00288	0.5143	--	--
	%TBF	-0.00267	0.5652	--	--
	VAT	-0.00182	0.5187	--	--
Overall Activity	LBM	--	--	-0.3933	0.631
	BMI	--	--	-0.1215	0.32
	WHR	--	--	-0.06486	0.29
	WC	--	--	-0.13557	0.23
	%TBF	--	--	-0.03473	0.64
	VAT	--	--	-0.00733	0.28

Abbreviations: MR, Mendelian randomization; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

5

**Table S4.** Heterogeneity estimates Mendelian Randomization Analyses for Physical Activity to Body Composition

Phenotypes		Self-report MR		Self-report MVMR		Accelerometer MR		Accelerometer MVMR	
Activity	Body Comp.	Q Stat.	p	Q Stat.	p	Q Stat.	p	Q Stat.	p
Sedentary Behavior	LBM	46.9	0.7435	--	--	22.2	0.035	--	--
	BMI	139.5	<0.001	--	--	18.9	<0.001	16.3	0.0121
	WHR	82.4	0.0098	253.7	<0.001	27	<0.001	17.5	0.0077
	WC	74	0.0766	145.1	<0.001	19.9	<0.001	18.2	0.0058
	%TBF	86.3	0.002	141.1	<0.001	11.8	0.0027	5.8	0.5654
Moderate PA	VAT	260.6	<0.001	315.8	<0.001	33.1	<0.001	160.8	<0.001
	LBM	4.2	0.8959	21.3	0.03	--	--	--	--
	BMI	33.2	0.0321	--	--	--	--	--	--
	WHR	18.4	0.4927	279.6	<0.001	--	--	--	--
	WC	24.8	0.1661	166.1	<0.001	--	--	--	--
MVPA	%TBF	13.9	0.788	150.8	<0.001	--	--	--	--
	VAT	52	0.0077	477.9	<0.001	--	--	--	--
	LBM	4.2	0.8959	21.3	0.03	18.9	<0.001	13.1	0.2867
	BMI	110.8	<0.001	--	--	96.9	<0.001	--	--
	WHR	91.2	0.0318	286.4	<0.001	66.6	<0.001	271.7	<0.001
Vigorous PA	WC	94	0.0245	187	<0.001	44.8	0.0089	171.8	<0.001
	%TBF	59.1	0.6155	160.6	<0.001	34.3	0.0613	144.4	<0.001
	VAT	202.9	<0.001	554.2	<0.001	183.5	<0.001	1709.7	<0.001
	LBM	21.7	0.2472	37.5	0.01	13.4	0.037	1.3	0.8616
	BMI	71.4	<0.001	--	--	20.4	0.119	--	--
SSOE	WHR	47.1	0.1242	279.8	<0.001	17.8	0.2154	278.4	<0.001
	WC	47.4	0.1685	153.3	<0.001	13.7	0.4738	165.1	<0.001
	%TBF	34.1	0.3687	165.8	<0.001	19	0.1666	162.6	<0.001
	VAT	148.8	<0.001	538.1	<0.001	90.8	<0.001	1308.6	<0.001
	LBM	25.5	0.8799	42.4	0.453	--	--	--	--
Overall Activity	BMI	190.7	<0.001	--	--	--	--	--	--
	WHR	132.2	<0.001	306.1	<0.001	--	--	--	--
	WC	98	0.0386	173.4	<0.001	--	--	--	--
	%TBF	51.1	0.9472	188.5	<0.001	--	--	--	--
	VAT	182.3	<0.001	458.4	<0.001	--	--	--	--
Overall Activity	LBM	--	--	--	--	4.1	0.1314	1.3	0.8616
	BMI	--	--	--	--	20.6	<0.001	38	<0.001
	WHR	--	--	--	--	14	0.0029	35.4	<0.001
	WC	--	--	--	--	2.6	0.4596	27	0.0046
	%TBF	--	--	--	--	2.5	0.4674	15.8	0.1986
VAT	--	--	--	--	11.5	0.0211	214.7	<0.001	

Abbreviations: MR, Mendelian randomization; MVMR, Multivariable Mendelian randomization; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: A large Q statistic and small p-value indicates instrument heterogeneity and potential violation of the MR assumptions. Where horizontal pleiotropy does not exist, observed values are expected to have smaller residuals, meaning that the variants do not deviate from the slope of the fitted regression. Inversely, variants deviating from the regression line suggest a pleiotropic effect.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

6

**Table S5.** MR-PRESSO Estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR		Accelerometer MR	
Activity	Body Comp.	Causal Estimate	p	Causal Estimate	p
Sedentary Behavior	LBM	0.26	0.3437	0.19*	0.0646
	BMI	0.23	<0.001	0.19*	0.1228
	WHR	0.23	<0.001	0.46**	0.0068
	WC	0.24	<0.001	0.43*	0.1844
	%TBF	0.22	0.0011	0.02	0.8574
	VAT	0.3	<0.001	0.12*	0.1826
Moderate PA	LBM	-0.44	0.5124	--	--
	BMI	0.08	0.4447	--	--
	WHR	0.22	0.0426	--	--
	WC	0.16	0.09	--	--
	%TBF	0.01	0.8836	--	--
	VAT	0.13	0.2193	--	--
MVPA	LBM	0.23	0.5223	0.18	0.6794
	BMI	0.08	0.1298	-0.11*	0.014
	WHR	-0.05	0.3468	-0.06	0.2577
	WC	0	0.9948	-0.06*	0.2203
	%TBF	-0.11	0.0588	-0.17	0.0114
	VAT	-0.08	0.2327	0*	0.9618
Vigorous PA	LBM	0.03	0.9778	0.98	0.3291
	BMI	-0.11	0.3991	-0.08	0.1982
	WHR	-0.23	0.0379	-0.06	0.2683
	WC	-0.16	0.2133	-0.09	0.1648
	%TBF	-0.1	0.4995	-0.22	0.0195
	VAT	-0.12	0.3949	-0.09*	0.0106
SSOE	LBM	-0.4	0.6289	--	--
	BMI	-0.49*	<0.001	--	--
	WHR	-0.21	0.0261	--	--
	WC	-0.3*	0.0124	--	--
	%TBF	-0.41	<0.001	--	--
	VAT	-0.33**	0.003	--	--
Overall Activity	LBM	--	--	0.05	0.205
	BMI	--	--	-0.31	0.0638
	WHR	--	--	-0.16*	0.4411
	WC	--	--	-0.16	0.139
	%TBF	--	--	-0.28*	0.3675
	VAT	--	--	-0.05*	0.3327

Abbreviations: MR, Mendelian randomization; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: MR-PRESSO evaluates for global pleiotropic effects by comparing the residual sum of squares with the expected distance under the null hypothesis for no horizontal pleiotropy. Individual pleiotropic outliers are identified by examining the observed and expected distributions of each variant; \*Value was corrected for outliers; \*\* MR-PRESSO estimate reached significance after outlier correction.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

7

**Table S6.** MR Median-based Estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR			Accelerometer MR		
Activity	Body Comp.	SNPs (#)	beta	95% CI	SNPs (#)	beta	95% CI
Sedentary Behavior	LBM	55	0.32	-0.54, 1.18	4	0.26	0.05, 0.46
	BMI	55	0.22	0.11, 0.33	4	0.35	0.07, 0.64
	WHR	56	0.24	0.12, 0.37	4	0.42	0.21, 0.63
	WC	59	0.22	0.11, 0.33	3	-0.13	-0.5, 0.24
	%TBF	53	0.34	0.18, 0.49	6	0.07	-0.08, 0.22
	VAT	128	0.24	0.2, 0.28	4	0.26	0.05, 0.46
Moderate PA	LBM	10	0.13	-2.26, 2.52	--	--	--
	BMI	21	0.11	-0.11, 0.32	--	--	--
	WHR	20	0.15	-0.1, 0.4	--	--	--
	WC	21	0.22	-0.02, 0.46	--	--	--
	%TBF	20	-0.05	-0.35, 0.25	--	--	--
	VAT	32	0.03	-0.05, 0.12	--	--	--
MVPA	LBM	39	0.57	-0.72, 1.86	14	0.25	-0.9, 1.4
	BMI	72	0.06	-0.06, 0.18	25	-0.1	-0.2, 0
	WHR	72	0.03	-0.11, 0.16	26	-0.03	-0.15, 0.09
	WC	72	-0.02	-0.15, 0.1	26	-0.06	-0.18, 0.06
	%TBF	66	-0.12	-0.28, 0.05	24	-0.15	-0.3, -0.01
	VAT	93	-0.07	-0.12, -0.01	46	-0.09	-0.16, -0.03
Vigorous PA	LBM	36	0.22	-2.43, 2.86	3	-1.07	-3.24, 1.09
	BMI	38	-0.05	-0.33, 0.23	15	-0.12	-0.25, 0.01
	WHR	38	0.09	-0.22, 0.39	15	-0.06	-0.23, 0.11
	WC	40	-0.2	-0.49, 0.1	15	-0.08	-0.23, 0.06
	%TBF	33	-0.07	-0.44, 0.3	15	-0.2	-0.4, 0
	VAT	61	-0.14	-0.24, -0.04	26	-0.19	-0.28, -0.1
SSOE	LBM	19	0.64	-2.57, 3.84	--	--	--
	BMI	74	-0.34	-0.59, -0.1	--	--	--
	WHR	74	-0.12	-0.38, 0.15	--	--	--
	WC	77	-0.06	-0.31, 0.19	--	--	--
	%TBF	72	-0.26	-0.59, 0.07	--	--	--
	VAT	98	-0.31	-0.41, -0.21	--	--	--
Overall Activity	LBM	--	--	--	7	1.12	-0.7, 2.93
	BMI	--	--	--	4	-0.18	-0.43, 0.07
	WHR	--	--	--	4	-0.16	-0.41, 0.1
	WC	--	--	--	4	-0.14	-0.35, 0.07
	%TBF	--	--	--	4	-0.29	-0.59, 0.02
	VAT	--	--	--	6	-0.21	-0.38, -0.03

Abbreviations: MR, Mendelian randomization; SNPs, single nucleotide polymorphisms; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: The weighted median causal estimates are based on the relationship between the strength of association between the SNP and exposure, and the SNP and outcome associations.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

8

**Table S7.** MR Mode-based Estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR			Accelerometer MR		
Activity	Body Comp.	SNPs (#)	beta	95% CI	SNPs (#)	beta	95% CI
Sedentary Behavior	LBM	55	0.38	-1.53, 2.29	4	0.24	0.02, 0.46
	BMI	55	0.21	-0.03, 0.44	4	0.37	-0.08, 0.82
	WHR	56	0.23	-0.01, 0.48	4	0.44	0.19, 0.69
	WC	59	0.16	-0.08, 0.39	3	-0.2	-0.58, 0.18
	%TBF	53	0.46	0.17, 0.75	6	0.03	-0.17, 0.24
	VAT	128	0.24	0.13, 0.36	4	0.24	0.02, 0.46
Moderate PA	LBM	10	0.4	-3.55, 4.35	--	--	--
	BMI	21	0.21	-0.21, 0.64	--	--	--
	WHR	20	0.14	-0.36, 0.63	--	--	--
	WC	21	0.2	-0.29, 0.69	--	--	--
	%TBF	20	-0.19	-0.75, 0.36	--	--	--
	VAT	32	0.04	-0.13, 0.21	--	--	--
MVPA	LBM	39	0.62	-2.2, 3.45	14	0.69	-1.56, 2.93
	BMI	72	-0.01	-0.32, 0.29	25	-0.1	-0.27, 0.07
	WHR	72	0.1	-0.24, 0.44	26	-0.06	-0.3, 0.18
	WC	72	0.05	-0.23, 0.32	26	-0.06	-0.34, 0.23
	%TBF	66	-0.2	-0.57, 0.17	24	-0.19	-0.44, 0.06
	VAT	93	-0.04	-0.17, 0.09	46	-0.03	-0.18, 0.11
Vigorous PA	LBM	36	3.5	-2.73, 9.74	3	-1.73	-4.91, 1.46
	BMI	38	-0.15	-0.86, 0.56	15	-0.17	-0.38, 0.03
	WHR	38	0.36	-0.17, 0.88	15	-0.05	-0.32, 0.22
	WC	40	0.4	-0.27, 1.07	15	-0.09	-0.35, 0.17
	%TBF	33	0.44	-0.44, 1.32	15	-0.27	-0.57, 0.03
	VAT	61	-0.09	-0.34, 0.15	26	-0.23	-0.39, -0.06
SSOE	LBM	19	4.44	-2.06, 10.93	--	--	--
	BMI	74	-0.51	-1.18, 0.16	--	--	--
	WHR	74	0.04	-0.85, 0.93	--	--	--
	WC	77	0.13	-0.48, 0.74	--	--	--
	%TBF	72	0.05	-0.67, 0.78	--	--	--
	VAT	98	-0.37	-0.73, 0	--	--	--
Overall Activity	LBM	--	--	--	7	2.56	-0.86, 5.98
	BMI	--	--	--	4	-0.04	-0.55, 0.46
	WHR	--	--	--	4	-0.04	-0.34, 0.26
	WC	--	--	--	4	-0.28	-0.61, 0.05
	%TBF	--	--	--	4	-0.26	-0.67, 0.15
	VAT	--	--	--	6	-0.17	-0.39, 0.06

Abbreviations: MR, Mendelian randomization; SNPs, single nucleotide polymorphisms; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: The weighted mode causal estimates are based on the relationship between the strength of association between the SNP and exposure, and the SNP and outcome associations. The weighted-mode estimate uses the property that valid instruments should provide the largest number of similar individual-instrument causal estimates even if the majority of instruments is invalid.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

9

**Table S8.** MR-EGGER Estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR			Accelerometer MR		
Activity	Body Comp.	SNPs (#)	beta	95% CI	SNPs (#)	beta	95% CI
Sedentary Behavior	LBM	55	0	-0.24, 0.24	4	-0.25	-1.24, 0.74
	BMI	55	0.53	0, 1.06	4	-0.32	-2.81, 2.17
	WHR	56	0.54	0.1, 0.99	4	0.33	-3.35, 4.02
	WC	59	0.37	-0.03, 0.77	3	0.14	-3.02, 3.29
	%TBF	53	1.03	0.45, 1.62	6	-1.87	-4.7, 0.95
	VAT	128	0.34	0.13, 0.56	4	0	-0.69, 0.69
Moderate PA	LBM	10	-0.05	-0.26, 0.15	--	--	--
	BMI	21	-0.36	-1.23, 0.51	--	--	--
	WHR	20	-0.14	-0.95, 0.67	--	--	--
	WC	21	0.09	-0.81, 0.99	--	--	--
	%TBF	20	0.18	-0.82, 1.18	--	--	--
	VAT	32	0.08	-0.21, 0.38	--	--	--
MVPA	LBM	39	-0.15	-0.35, 0.05	14	0.03	-0.08, 0.15
	BMI	72	-0.08	-0.57, 0.41	25	0.11	-0.44, 0.66
	WHR	72	-0.2	-0.74, 0.35	26	0.1	-0.44, 0.63
	WC	72	0.11	-0.4, 0.61	26	0.1	-0.33, 0.54
	%TBF	66	-0.12	-0.7, 0.46	24	0.18	-0.32, 0.68
	VAT	93	-0.18	-0.4, 0.03	46	-0.09	-0.24, 0.06
Vigorous PA	LBM	36	0.35	-0.05, 0.75	3	0.18	-0.39, 0.75
	BMI	38	0.62	-0.41, 1.65	15	-0.26	-1.31, 0.8
	WHR	38	0.55	-0.44, 1.53	15	-0.09	-1.26, 1.09
	WC	40	0.81	-0.08, 1.71	15	-0.27	-1.27, 0.74
	%TBF	33	0.05	-1.07, 1.17	15	-1.37	-2.76, 0.03
	VAT	61	0.05	-0.2, 0.3	26	-0.24	-0.51, 0.02
SSOE	LBM	19	0.01	-0.3, 0.31	--	--	--
	BMI	74	0.32	-1.08, 1.72	--	--	--
	WHR	74	0.2	-1.13, 1.54	--	--	--
	WC	77	0.1	-0.98, 1.18	--	--	--
	%TBF	72	0.01	-1.41, 1.42	--	--	--
	VAT	98	-0.09	-0.45, 0.26	--	--	--
Overall Activity	LBM	--	--	--	7	0.14	-0.07, 0.35
	BMI	--	--	--	4	4.15	-2.54, 10.84
	WHR	--	--	--	4	4.58	-1.08, 10.24
	WC	--	--	--	4	2.19	-1.08, 5.46
	%TBF	--	--	--	4	0.95	-3.56, 5.47
	VAT	--	--	--	6	-0.25	-1.08, 0.57

Abbreviations: MR, Mendelian randomization; SNPs, single nucleotide polymorphisms; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue.

Note: MR-Egger regression assumes linearity and homogeneity in the associations between the genetic variants, risk factor, and outcome. Egger regression provides an estimate of the causal effect that is consistent asymptotically even if all the genetic variants have pleiotropic effects on the outcome.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

10

**Table S9.** CAUSE estimates for Physical Activity to Body Composition

Phenotypes		Self-Report MR		Accelerometer MR	
Activity	Body Comp.	Effect	95% CI	Effect	95% CI
Sedentary Behavior	LBM	0.00	-0.36, 0.40	0.42	-1.04, 1.85
	BMI	0.14	0.10, 0.18**	0.05	-0.06, 0.17
	WHR	0.13	0.08, 0.17**	0.02	-0.12, 0.17
	WC	0.16	0.11, 0.20**	0.03	-0.15, 0.20
	%TBF	0.17	0.11, 0.24	0.12	-0.08, 0.33
	VAT	0.23	0.19, 0.28***	0.10	-0.01, 0.21
Moderate PA	LBM	-0.69	-2.18, 0.81	--	--
	BMI	0.07	-0.11, 0.26	--	--
	WHR	0.20	0.02, 0.39	--	--
	WC	0.13	-0.10, 0.36	--	--
	%TBF	0.04	-0.24, 0.32	--	--
	VAT	0.25	0.02, 0.47	--	--
MVPA	LBM	0.51	-0.15, 1.24	0.00	-0.47, 0.47
	BMI	-0.02	-0.11, 0.07	-0.13	-0.19, -0.07
	WHR	-0.03	-0.11, 0.06	-0.08	-0.15, -0.02
	WC	-0.03	-0.12, 0.07	-0.12	-0.19, -0.05
	%TBF	-0.08	-0.20, 0.03	-0.11	-0.19, -0.03
	VAT	-0.06	-0.16, 0.04	-0.11	-0.15, -0.07***
Vigorous PA	LBM	0.59	-0.84, 2.05	0.15	-0.42, 0.75
	BMI	0.00	-0.18, 0.19	-0.08	-0.16, 0.01
	WHR	-0.03	-0.20, 0.14	-0.02	-0.10, 0.06
	WC	0.02	-0.16, 0.21	-0.07	-0.16, 0.02
	%TBF	-0.11	-0.34, 0.12	-0.07	-0.19, 0.05
	VAT	-0.13	-0.33, 0.06	-0.14	-0.19, -0.08***
SSOE	LBM	1.36	-0.19, 2.82	--	--
	BMI	-0.19	-0.36, -0.02	--	--
	WHR	-0.22	-0.37, -0.08	--	--
	WC	-0.22	-0.39, -0.05	--	--
	%TBF	-0.34	-0.56, -0.11	--	--
	VAT	-0.28	-0.46, -0.11	--	--
Overall Activity	LBM	--	--	0.18	-0.98, 1.31
	BMI	--	--	-0.12	-0.28, 0.04
	WHR	--	--	-0.09	-0.25, 0.08
	WC	--	--	-0.11	-0.30, 0.07
	%TBF	--	--	-0.17	-0.35, 0.02
	VAT	--	--	-0.08	-0.19, 0.03

Abbreviations: MR, Mendelian randomization; SNPs, single nucleotide polymorphisms; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises; LBM, lean body mass; BMI, body mass index; WHR, waist-to-hip ratio; WC, waist circumference; %TBF, total body fat percent; VAT, visceral adipose tissue. Note: \*\*\*p<.001, \*\*p<.01

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

11

**Table S10.** Bidirectional MR IVW Estimates for Body Composition to Physical Activity

<i>Body Comp.</i>	<b>Phenotypes</b> <i>Activity</i>	<b>Self-Report MR</b>			<b>Accelerometer MR</b>		
		<i>SNPs (#)</i>	<i>beta</i>	<i>95% CI</i>	<i>SNPs (#)</i>	<i>beta</i>	<i>95% CI</i>
LBM <sup>†</sup>	Sedentary Behavior	15	0.00	-0.00, 0.00	--	--	--
	Moderate PA	16	0.00	-0.01, 0.00	--	--	--
	MVPA	16	0.00	-0.00, 0.01	16	0.00	0.00, 0.01
	Vigorous PA	16	0.00	-0.00, 0.00	16	0.01	0.00, 0.02
	SSOE	16	0.00	-0.00, 0.00	--	--	--
	Overall Activity	--	--	--	--	--	--
BMI	Sedentary Behavior	67	0.09	0.08, 0.10	11	0.12	-0.03, 0.26
	Moderate PA	78	0.01	-0.02, 0.03	12	-0.04	-0.16, 0.07
	MVPA	78	-0.01	-0.04, 0.03	71	-0.03	-0.08, 0.02
	Vigorous PA	78	0.01	-0.01, 0.03	71	-0.08	-0.13, -0.04
	SSOE	78	0.00	-0.02, 0.02	--	--	--
	Overall Activity	--	--	--	11	-0.03	-0.17, 0.11
%TBF	Sedentary Behavior	6	0.12	-0.04, 0.29	2	0.11	-0.18, 0.40
	Moderate PA	9	-0.05	-0.10, 0.00	2	0.07	-0.16, 0.31
	MVPA	9	-0.05	-0.17, 0.05	9	-0.02	-0.14, 0.10
	Vigorous PA	9	-0.00	-0.06, 0.05	9	-0.00	-0.11, 0.10
	SSOE	9	0.01	-0.03, 0.04	--	--	--
	Overall Activity	--	--	--	2	0.07	-0.3, 0.45
VAT	Sedentary Behavior	18	0.05	-0.01, 0.11	28	0.23	0.10, 0.36
	Moderate PA	19	-0.01	-0.05, 0.03	29	-0.12	-0.25, 0.01
	MVPA	17	0.02	-0.03, 0.07	236	-0.17	-0.22, -0.12
	Vigorous PA	17	0.03	0.00, 0.06	211	-0.19	-0.23, -0.14
	SSOE	17	0.03	0.02, 0.05	--	--	--
	Overall Activity	--	--	--	27	-0.10	-0.25, 0.05


Abbreviations: MR, Mendelian randomization; SNPs, single nucleotide polymorphisms; LBM, lean body mass; BMI, body mass index; %TBF, total body fat percent; VAT, visceral adipose tissue; PA, physical activity; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises.

<sup>†</sup>LBM contained one SNPs at the genome-wide significance level; a reduced IV threshold ( $p < 5 \times 10^{-6}$ ) was used for analyses. Outcome SNPs could not be identified in GWAS of sedentary behavior, moderate PA, or overall activity.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

12

SNP ID	Chromosome	Position	Gene	Allele	EAF	B	SE	<i>p</i>
rs6545389	2	54386901	ACYP2	A / G	0.4997	-0.0127	0.0023	3.5e-08




**Figure S1.** Manhattan Plot and genome-wide significant SNPs for self-reported moderate PA

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

13


SNP ID	Chromosome	Position	Gene	Allele	EAF	B	SE	<i>p</i>
rs4480415	1	174800121	RABGAP1L	G / A	0.7248	-0.012	0.0023	4.6e-08
rs17443704	5	151668231	--	G / A	0.8231	-0.013	0.0023	3.5e-08



**Figure S2.** Manhattan Plot and genome-wide significant SNPs for accelerometer-measured MVPA

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION


14



**Figure S3.** Univariable Mendelian Randomization IVW Sensitivity Analysis for Self-Reported Physical Activity Levels and Body Composition removing the *APOE* rs429358 variant. Abbreviations: SNPs, single nucleotide polymorphisms; CI, confidence interval; METs, metabolic equivalent of tasks; BMI, body mass index; WC, waist circumference; WHR, waist-to-hip ratio; LBM, lean body mass; %TBF, total body fat percent; VAT, visceral adipose tissue; MVPA, moderate-to-vigorous PA; SSOE, strenuous sports or other exercises.

## MENDELIAN RANDOMIZATION OF PHYSICAL ACTIVITY AND BODY COMPOSITION

15



**Figure S4.** Univariable Mendelian Randomization IVW Sensitivity Analysis for Accelerometer-Measured Physical Activity Levels and Body Composition removing the *APOE* rs429358 variant. Abbreviations: SNPs, single nucleotide polymorphisms; CI, confidence interval; METs, metabolic equivalent of tasks; BMI, body mass index; WC, waist circumference; WHR, waist-to-hip ratio; LBM, lean body mass; %TBF, total body fat percent; VAT, visceral adipose tissue; MVPA, moderate-to-vigorous PA.