


Training interventions to reduce the risk of injury to the lower extremity joints during landing movements in adult athletes: a systematic review and meta-analysis

Chantal Bathe ¹, Lena Fennen,¹ Theresa Heering,^{2,3} Alexander Greif,¹ Rosemary Dubbeldam¹

To cite: Bathe C, Fennen L, Heering T, *et al*. Training interventions to reduce the risk of injury to the lower extremity joints during landing movements in adult athletes: a systematic review and meta-analysis. *BMJ Open Sport & Exercise Medicine* 2023;**9**:e001508. doi:10.1136/bmjsem-2022-001508

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2022-001508>).

Accepted 26 April 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Movement Science, Westfälische Wilhelms University Münster, Münster, Germany

²Coventry University, Coventry, UK

³Deakin University School of Health and Social Development, Melbourne, Victoria, Australia

Correspondence to

Dr Rosemary Dubbeldam;
r.dubbeldam@uni-muenster.de

ABSTRACT

Objective Aim of this systematic review was to summarise training interventions designed to reduce biomechanical risk factors associated with increased risk of lower extremity landing injuries and to evaluate their practical implications in amateur sports.

Design Systematic review and meta-analysis.

Data sources MEDLINE, Scopus and SPORTDiscus.

Eligibility criteria Training intervention(s) aimed at reducing biomechanical risk factors and/or injury rates included the following: (1) prospective or (non-)randomised controlled study design; (2) risk factors that were measured with valid two-dimensional or three-dimensional motion analysis systems or Landing Error Scoring System during jump landings. In addition, meta-analyses were performed, and the risk of bias was assessed.

Results Thirty-one studies met all inclusion criteria, capturing 11 different training interventions (eg, feedback and plyometrics) and 974 participants. A significantly medium effect of technique training (both instruction and feedback) and dynamic strengthening (ie, plyometrics with/without strengthening) on knee flexion angle ($g=0.77$; 95% CI 0.33 to 1.21) was shown. Only one-third of the studies had training interventions that required minimal training setup and additional coaching educations.

Conclusion This systematic review highlights that amateur coaches can decrease relevant biomechanical risk factors by means of minimal training setup, for example, instructing to focus on a soft landing, even within only one training session of simple technique training. The meta-analysis emphasises implementing technique training as stand-alone or combined with dynamic strengthening into amateur sport training routine.

INTRODUCTION

Sport injuries are internationally recognised as a public health problem not only in elite sport, but also at the amateur level.¹ In North Rhine-Westphalia, 40% of all sport injuries occur during sporting activities such as team ball sports or gymnastic and most frequently in the lower extremities, like the

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Lower limb injuries in sports occur due to high-risk movements like jump landing tasks with rapid deceleration and stops.
- ⇒ A reduced knee flexion angle, improved ground reaction force and a knee valgus during landing movements are risk factors for lower extremity injuries.
- ⇒ Performing injury prevention programmes at least twice a week improved neuromuscular and motor performance and reduces the injury risk.

WHAT THIS STUDY ADDS

- ⇒ Technique training, that is, external instruction and feedback on jump landing performance, and dynamic strengthening, that is, plyometrics with increasing intensity, can reduce biomechanical risk factor during landing movements.
- ⇒ Static balance training on a wobble board alone had no significant effect in reducing risk factors for jump landing injuries and is, therefore, not recommended.
- ⇒ Prevention of jump landing injuries requires moderate time consumption (around 30 min), basic training material like cones and hurdles and mostly no specific trainer education.
- ⇒ Future studies should consider analysing jump landing movements using an easy to apply, validated method such as the Landing Error Scoring System to allow for direct comparison with previous research

knee and ankle joint². Fractures (38%) and ligament ruptures (34%) of the ankle and knee form the largest part of these lower extremity injuries.²⁻³ Regardless of gender, the consequences of these lower extremity injuries, such as anterior cruciate ligament (ACL) rupture, ankle sprain or patella tendinopathy, are severe. For instance, an ACL rupture is accompanied by a long and expensive rehabilitation period and increased reinjury rates especially in athletes aged under 20 years.^{4,5} Most lower extremity injuries occur during complex movements



such as a jump-landing and change in direction tasks (eg, side-cutting).⁶⁻⁷ Single or double limb landing manoeuvres with rapid decelerations, stops or repetitive jump landings were identified as frequent injury mechanisms.⁸⁻¹¹ A well-balanced landing strategy is essential for effective absorption of impact forces from landing. Therefore, among others, synergistic lower extremity joint coordination and alignment of the hip, knee and ankle joint in the sagittal plane, with dynamic muscle control of the lower extremity and upper body (eg, core muscle), is required.^{9 12 13}

Biomechanical risk factors for jump landing injuries have been identified in movement analyses studies. A reduced range of motion of the lower extremity joints (eg, less knee or hip flexion) can lead to a stiffer landing technique with a decrease in force absorption, which can subsequently increase the risk of lower extremity injury.^{9 14} In particular, a knee flexion angle smaller than 30° (ie, a more extended knee) at initial contact during a single or double limb landing may affect ACL load.^{15 16} Consequently, at a reduced knee flexion angle, the synergistic hamstring muscle force is directed parallel to the ACL, which is placed vertically to the tibia plateau, thus limiting the hamstring potential to counteract stress on the ACL due to anterior tibial shift.^{9 11 17-21} An increased knee valgus based on simultaneous hip adduction and internal rotation during closed-chain knee flexion with additional ankle eversion is another underlying biomechanical risk factor for jump landing injuries.^{22 23} Lastly, a decrease in ankle plantar flexion during initial contact results in less ankle dorsiflexion during subsequent landing manoeuvre.⁹ This reduced ankle dorsiflexion has been associated with knee overuse injuries such as patellofemoral pain^{8 9 24} and ankle inversions trauma.^{25 26}

Due to the high prevalence and possibly severe consequences of jump landing injuries, adequate injury prevention, that is, reducing injury risk by improving the biomechanical risk factors, is crucial. Injury prevention has already been implemented in amateur and professional sport settings²⁷ and multiple training interventions have been reported. These training interventions are intended, among others, to improve the athlete sports performance, reduce injury risk and costs of injury treatments for the club and athletes.²⁷ For example, training interventions, such as FIFAs 11+ twentymin warm up programme developed for football players of all ages, reduce the general injury risk up to 35%, and have been incorporated into the training routine of some football clubs.^{28 29} Performing the FIFA training intervention at least twice per week leads to improvement in neuromuscular and motor performance.^{28 29} Training interventions have also been developed to specifically reduce the likelihood of injury after jump landing movements. For example, Aerts *et al* advise to perform their training intervention twice a week, including gradually increasing lower extremity strengthening, plyometrics, and technique instructions, that is, teach the athlete how to align the lower extremity joints during landing.³⁰

Despite the need and availability of effective injury prevention programmes, detailed training interventions are often not a part of a normal training routine, especially in an amateur sports setting. As an example, significant injury reduction depends, among others, on the qualification of the coach (eg, knowledge about injury prevention) and medical monitoring, which is mostly limited in an amateur sports setting.³¹ An analysis of the integration of injury prevention in general amateur sports showed that only 21% of 70 amateur coaches used specific training interventions such as FIFA 11+ in football.² Furthermore, new evidence-based training interventions are published almost annually, which is not easy to summarise and integrate into a training routine for amateur and professional coaches and athletes.³² To successfully implement training interventions in amateur sport settings, training materials must be affordable and self-explanatory, as specialised staff such as an educated programme controller (ie, athletic trainer) or physiotherapist are often not available. Considering the limited training time in amateur sports, training interventions should not be time-consuming. Thus, there is a need to summarise evidence-based training interventions that improve jump landing manoeuvre and evaluate whether the programme is implementable in amateur sports.

The aim of this study is to systematically review training interventions for adult athletes that aim to reduce biomechanical risk factors for lower extremity joint injuries during jump landing performances, and to critically evaluate them, regarding their practicability, in terms of required materials, coach education and time consumption, in amateur sports settings.

METHODS

This review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines from 2020.^{33 34}

Search strategy

The electronic databases (search engines) MEDLINE (PubMed), SPORTDiscus (EBSCO) and Scopus were searched from the database's inception to 20 December 2021. The combination of the following keywords was used as search terms: 'athletes OR sports'; 'unilateral landing OR bilateral landing OR jump-landing OR jump task OR landing'; 'intervention protocol OR intervention OR training protocol OR exercise OR training OR prevention'; 'kinematics OR kinetics OR biomechanics OR knee flexion angle OR knee valgus OR injury risk OR injury rate OR landing error scoring system' (online supplemental table 2).

Study selection

The eligibility criteria were created using the PICO(S) model for clinical questions and are summarised in table 1. Included interventions aimed to improve risk factors during jump landing performance and compared an intervention group to a control group with sham or

Table 1 Study criteria for inclusion in the review

Participants	Adult male and female athletes (>18 to <45 years), with a minimum of two training sessions per week or at least active college or comparable athlete, without acute (<6 weeks) or chronic (>6 weeks) injury to the lower extremity. Participants did not receive surgery of the lower extremity joints because of a rupture or fracture.
Intervention	Training programmes or interventions which aimed to improve the jump landing strategy or injury risk and thus the biomechanical risk factors for a single or double leg jump landing injury. Interventions must be integrable into amateur sports setting, that is, no need for specialised staff such as an educated programme controller (ie, athletic trainer) or physiotherapist to implement the programme.
Control group	If there is a control group, it should receive a sham treatment or no intervention.
Outcome	Measurement of lower extremity injury rate, either laboratory based, for example, 3D-motion analysis and force plate data, or by means of in-field measures, for example, valid scoring system and expert ratings like the Landing Error Scoring System, while examining single and double leg landings. The biomechanical risk factors consist of increased knee valgus, smaller hip flexion angle, reduced knee flexion angle/moment, or increased GRF especially of the knee.
Study type	Prospective/retrospective or observational (non-)randomised controlled intervention trials.

3D, three dimensional; GRF, ground reaction force.

without additional intervention ((non)-randomised controlled trials, RCTs) or to the intervention group's baseline levels (eg, pre-post or cross-over trials).³⁵ The outcome measurements should be assessed using reliable tools (eg, VICON Motion Systems, Landing Error Scoring System (LESS)) and analyse jump performance (eg, drop jumps, countermovement jumps).

English and German publications were considered, and geographical restrictions were not applied. Studies were excluded when participants were soldiers, because of their special military training programme. Also, simulation studies were excluded.

Two independent reviewers applied predetermined eligibility criteria to screen titles and abstracts in the Rayyan-Intelligent systematic review software (Rayyan System, Cambridge, Massachusetts, USA).³⁶ An abstract was included when the two reviewers independently assessed it as satisfying the inclusion criteria. Afterwards, full texts were screened and were excluded if they did not fulfil inclusion criteria. Conflicting classifications, of both abstracts and full texts, were discussed and resolved.

Data extraction

Data extraction included: Study identifiers (authors name, year of publication), study design, study sample size and participants' characteristics (age, gender, activity level). The data were organised according to the training intervention methods, and results were added for the main outcome variables (eg, biomechanical risk factors). Training interventions were classified as (A) technique training, that is, improving the landing technique by instructions, feedback or skill training, (B) dynamic strengthening, that is, plyometrics or a combination of strengthening and plyometrics, (C) static strengthening, that is, strengthening without plyometrics or (D) balance training (static or dynamic). An included training intervention was defined as a 'prevention programme' if an apparent sequence of different forms of exercises was used and the author of the study titled this form of

training as a 'prevention programme'. In addition, the different gender responses (ie, men and women) to the training interventions were identified. Furthermore, in terms of practicability of training interventions, the required training materials were sorted according to the training techniques (A–D) and the education of the coach and training time consumption were listed.

Methodological assessment

Two independent reviewers assessed the risk of bias. The Cochrane Risk of Bias tool for randomised trials (RoB2)³⁷ was applied. The aimed target trial for the analysed studies, was a randomised clinical trial with a control and an intervention group consisting of female and male amateur athletes performing an intervention.^{33 34} In addition, the intention-to-treat effect was chosen to interpret study outcomes.³⁸ The 'Risk of bias in Non-randomised Studies-of Interventions' (ROBINS-I) tool was used to analyse other study designs.³⁸ Online supplemental table 3 summarises the adapted controlled questions for the corresponding seven bias domains.³⁸

Data analysis

For RCTs with a biomechanical risk factor as a primary outcome in results, the standardised mean differences (SMDs), known as Cohen's d effect sizes, with associated 95% CIs were calculated, and a meta-analysis was performed.³⁹

Moreover, to avoid within-group effect size correlation, the SMDs were only calculated if two experimental groups or an additional control group were analysed (ie, RCT or CT study design).⁴⁰ If a study contributed more than one effect size to the meta-analysis, a combining group approach (eg, putting means and SD of Group A and B together) was used as basis for effect size calculation³⁷ to circumvent double-counting (eg, comparing group A with group C and group B with group C again). In addition, to capture the effectiveness of the intervention, only the effect size of the measurement directly after

the intervention (ie, post-test) was considered if multiple measures were collected.

As the considerable between-study heterogeneity was anticipated, a random-effects model was used to pool effect sizes. The ‘DerSimonian-Larid’ estimator was used to calculate the heterogeneity variance τ^2 .⁴¹ Knapp-Hartung adjustment⁴² was applied to calculate the CIs around the pooled effect. Moreover, the heterogeneity index I^2 with its 95% CI and additional prediction intervals were calculated. I^2 was interpreted according to Higgins and Thompson 25%=low; 50%=moderate; 75%=substantial. In addition, if $I^2 > 50%$ an outliers (ie, CI of studies did not overlap with CI of pooled effect) and influence analyses (ie, leave-one-out method) were performed to study the robustness of the true effect.^{43–45} Finally, asymmetries of the effect size distribution due to bias distribution were assessed using Egger’s test⁴⁶ and were visually examined using a funnel plot of effect sizes relative to SE. All analyses were performed by the ‘meta’ V.5.2–0 of R language (R core team, Vienna, Austria). The thresholds for the interpretation of the effect sizes were as follows: 0.20=small; 0.50=moderate; 0.80=large.⁴⁷ Statistical significance was set at $p < 0.05$.

RESULTS

Search results

The initial database search yielded 2387 results. After removal of duplicates, 1717 records remained. Title and abstract screening resulted in 133 full-text articles that were assessed based on the inclusion criteria. The final full text screening resulted in 31 included studies, of which the flow of the PRISMA selection process is summarised in figure 1. The intervention studies included 15 RCT’s, 15 pre–post intervention studies without a control group, and one cross-over intervention study. The study characteristics are presented in table 2 and additional details are presented in online supplemental table 1.

Study characteristics

Sample sizes of the RCTs ranged from 16 to 116 participants with an overall sample size of 682. The sample size of the pre–post intervention studies ranged from 8 to 37 participants with an overall sample size of 280. Eight participants participated in the single cross-over study. Eighteen of the 31 studies included only female participants, 5 studies had only male participants and 8 studies analysed both men and women. Accordingly, more than half of the results were derived from the analysis of female athletes (64%). Three-quarter of all participants were amateur athletes (730), followed by division 1 and 2 collegiate athletes (184) and high-performance athletes (60).

Intervention characteristics

Fifteen interventions were classified as technique training (A), 12 as dynamic strengthening (B), 2 as static strengthening (C) and 2 as balance training (D). The training intervention duration varied between studies with either




Figure 1 PRISMA flow chart of study selection process. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

1 single training session in total (18 studies) or multiple training sessions performed 2–3 times per week for a total duration of three to 12 weeks (13 studies) (online supplemental table 1). The session duration of each training intervention lasted between 5 and 15 min (10 studies), 12–20 min (3 studies), 15–30 min (4 studies), 40–60 min (5 studies). In 10 studies, no exact training session duration was given, but they lasted 1 (8 studies), 2 (1 study) or a maximum of 3 (1 study) training sessions.

Different types of double limb (22 studies) and single limb (9 studies) jump landing tasks were observed to assess the outcome measurements and included variants of drop jumps (19 studies) and/or countermovement jumps (6 studies), volleyball, basketball or netball-specific jumps (3 studies), tuck jump (1 study), stop jump task (2 studies), lateral single leg jump (1 study) and maximal vertical jump (1 study).

Intervention outcomes

Fifteen studies reported ground reaction force (GRF) outcomes, 23 studies reported joint angle, 5 studies reported joint moment outcome and 3 studies reported LESS as outcome. All but two studies^{48 49} reported significant differences in outcomes before and after the intervention.

A significant reduction of biomechanical risk factors was reported for all four training types (A–D). In the following, the findings of the 15 RCTs organised by training type (A–D) and gender are reported to focus

Table 2 Characteristics and (non-)significant main results of included studies, assigned to technique training, dynamic strengthening, static strengthening or balance training

Author	Study design	Total N	Athletes	Training sessions (1/more)	GRF	Joints	LESS
A. Technique training							
Jump landing performance—instruction							
Almonroeder ⁶⁵	Pre–post	16	A	1		>knee and hip flexion angle	
Benjaminse ⁵⁴	RCT	40	S.p.	1			Improved LESS
Chijimatsu ⁷⁹	Pre–post	15	A	1		<internal rotation angle	
McNair ⁵⁶	RCT	80	A	1	<peak GRF		
Milner ⁸⁰	Cross-over	12	A	1	<peak vGRF	>peak knee flexion angle	
Mizner ⁷⁰	Pre–post	37	A	1	<peak vGRF	>peak knee flexion angle	
Tate ⁵²	RCT	26	A	1	<peak vGRF	>knee flexion angle knee abduction angle (n.s).	
Turner ⁶⁴	Pre–post	24	P	1		>knee abduction angle	
Welling ⁵¹	RCT	40	A	1	<vGRF	>knee flexion angle	
Jump landing performance—feedback							
Cronin ⁸¹	Pre–post	15	P	1	<vGRF		
Etnoyer ⁵⁷	RCT	43	A	1		>peak knee flexion angle	
Leonard ⁸²	Pre–post	23	A	1	<vGRF		
Oñate ⁵⁵	RCT	51	A	1	<peak vGRF	>knee angular displacement	
Shams ⁵³	RCT	45	A	1			Improved LESS
Jump landing performance—skill training							
Shimokochi ⁶⁹	Pre–post	20	A	1	<GRF	>knee flexion angle	
B. Dynamic strengthening							
Plyometrics							
Dello Iacano ⁸³	Pre–post	18	P	More	>peak GRF		
Herrington ⁸⁴	Pre–post	15	A	More		<knee valgus angle	
Nagano ⁸⁵	Pre–post	8	A	More		>knee flexion angle	
Makaruk ⁶²	RCT	36	A	More	<vGRF	>knee flexion angle	
Vescovi ⁴⁸	RCT	20	A	More	<vGRF		
Combi strengthening							
Dai ⁸⁶	Pre–post	28	A	1		>hip and knee flexion angle	
Peng ⁸⁷	Pre–post	12	A	1	<peak GRF	>hip and knee flexion angle >ankle plantarflex. and dorsiflex. angle	
Stearns and Powers ⁸⁸	Pre–post	21	S.p.	1		>knee and hip flexion angle <peak knee abduction angles	
Yang ⁶⁰	RCT	36	A	1	♂ < GRF ♀ < GRF (n.s.)	♂ > knee and hip flexion ♀ > knee and hip flexion (n.s.)	
Prevention programmes							
Aerts ⁵⁸	RCT	116	A	More		>hip flexion and maximal left knee flexion angle (♂)	
Fox ⁵⁹	RCT	16	A	More		>hip external rotation angle >knee angular displacement	
O'Malley ⁶¹	RCT	78	A	1			Improved LESS

Continued

Table 2 Continued

Author	Study design	Total N	Athletes	Training sessions (1/more)	GRF	Joints	LESS
C. Static strengthening							
Core workout							
Araujo ⁸⁹	Pre–post	16	A	More	<peak vGRF		
Warm-up programme							
Avedesian ⁹⁰	Pre–post	12	A	1		<peak hip adduction angle >knee abduction and internal rotation angle	
D. Balance training							
Static balance training							
Silva ⁴⁹	RCT	24	A	1	vGRF (n.s.)	Plantar flexion angle (n.s.)	
Dynamic balance training							
Letafatkar ⁶³	RCT	31	A	1		<knee flexion angle	

A, amateur; dorsiflex, dorsiflexion; GRF, ground reaction force; LESS, Landing Error Scoring System; n, sample size; n.s., non-significant; P, professional; plantarflex, plantarflexion; RCT, randomised controlled trial; S.p., semi professional; vGRF, vertical GRF.

on cause-and-effect relationships instead of reporting correlations.⁵⁰ In addition, the findings of the meta-analyses are reported.

Technique training focused on improving the athlete's landing technique through three forms of instruction (three RCTs) and feedback (four RCTs). Forms of instruction consisted of external focus (directing the athlete's attention away from the movement), internal focus (direct focus on the movement) and video instruction.

Postintervention, all three forms of instructions led to a significant improvement in knee flexion angle^{51 52} and a significant reduction in knee valgus moment in drop jump landing.⁵¹ A significant decrease in GRF was reported after all forms of technique training, except video instruction. The LESS scoring was significantly improved (fewer errors) when using external focus and video instruction or feedback.^{51–56}

The long-term effect of technique training on biomechanical risk factors is not clear: 1-week postintervention, the vertical GRF during landing can increase again^{51 52} or further improve, that is, become less during landing.⁵⁵ The positive effect of the intervention on the knee joint angle seems to reduce after 1–4 weeks^{52 57} or there is no clear trend.⁵¹ Benjaminse *et al* demonstrated an improved LESS and knee joint angle for external focus and video instruction groups 1 week after the intervention compared with baseline, however, postintervention data were not reported.⁵⁴

Dynamic strengthening training interventions included plyometric exercises (two RCTs), a combination of plyometrics and lower extremity strengthening (one RCT) and plyometric prevalent prevention programmes consisting of exercises including plyometrics but where trainers are also instructed to improve technique (three RCTs).

One prevention programme with plyometric exercises of increasing intensity significantly improved the knee flexion angle landing pattern in males.⁵⁸ A reduction in knee valgus and improvement of hip flexion and external rotation angle were reported after dynamic strengthening.^{58 59} Another programme used a combination of strengthening and plyometric exercises and was able to improve knee flexion angle at peak impact.⁶⁰ In addition, one prevention programme consisted of a warm-up programme combining strengthening, stretching and balance exercises. The application of this programme significantly decreased LESS scores directly after an 8-week intervention.⁶¹ No clear effects of plyometrics were observed for the GRF, where both significant⁶² and non-significant findings were reported.⁴⁸

Only Yang *et al* studied the long-term effect of a 4-week dynamic strengthening intervention. The positive effect on the knee flexion angle remained over time (up to 12 weeks), while the initial positive effect of the training on the vertical GRF was not consistent over time.⁶⁰

Balance training included static balance training on a wobble board (ie, no external perturbation) (one RCT) and dynamic balance training consisting of increasing perturbation-enhanced neuromuscular training, such as sport-specific technique performance on a rocker board (one RCT).

The intervention programme focusing on static balance⁴⁹ did not result in significant changes in biomechanical risk factors, whereas the dynamic balance training significantly increased the initial knee flexion angle during landing.⁶³

Regarding gender differences, both genders significantly improved risk factors in terms of knee flexion angle significantly by technique training such as teaching plantarflexed landing, feedback, external instruction,^{51 54–56} and dynamic balance training with increasing

perturbation-enhanced neuromuscular tasks⁶³ (five RCTs).

For meta-analyses, 15 RCTs were available. Due to inconsistent evaluation and reporting of outcome variables, only two variables, the knee flexion angle and vertical GRF, could be examined. The meta-analyses used 11 RCTs in total.

Four studies showed a large (>0.80),^{52 57 58 63} one study showed a moderate (>0.50)⁵⁷ and one study a small (>0.20)⁵¹ effect size for knee flexion angle. The between-study heterogeneity variance was estimated at $\tau^2 = 0.53$ (95% CI 0.25 to 5.72) with a high heterogeneity ($I^2 = 83%$; 95% CI 64% to 92%). Based on substantial heterogeneity in the main analysis ($I^2 > 50%$), an outliers and influence analysis was performed and effect sizes of Letefatkar *et al* and Makaruk *et al* were removed from the random-effect model.^{62 63} Finally, a significantly medium pooled effect size ($g = 0.77$; 95% CI 0.33 to 1.21) was found for the three technique training interventions and one prevention programme (dynamic strengthening) on knee flexion angle ($p = 0.01$).

A large effect size (>0.80) for GRF was shown in one study.⁵⁶ Four studies showed a small effect size (>0.20)^{51 52 55 62} and three studies had no effect.^{48 49 60} The

between-study heterogeneity was estimated at $\tau^2 = 0.11$ (95% CI 0.00 to 0.80) with a moderate heterogeneity ($I^2 = 48%$; 95% CI 0 to 77%) and a non-significant pooled small effect ($g = -0.24$; 95% CI -0.63 to 0.16).

Publication bias analysis of both meta-analyses showed no asymmetry in a visual inspection of funnel plots, and application of Egger's test was not possible due to the limited number of studies analysed (<10 studies each) (figure 2).

Required material and coach education characteristics

Table 3 summarises the practicability of the interventions in terms of training material and coach education required to perform the intervention. Most interventions required materials such as boxes (height ranged from 20 to 50 cm), a BOSU, ball or balance board. In addition, cones, hurdles, walls and mattresses were the second most common training materials. For feedback interventions, a video camera and monitor were needed in two out of four interventions, and in one intervention, a video of an expert performing a jump landing model must be provided. For instruction interventions, 3 out of 10 studies required an expert performing or instructing the landing manoeuvre on video.




Figure 2 Meta-analysis of knee flexion angle and ground reaction force (GRF) outcome variable.

**Table 3** Required training material and intervention description of included studies ordered by training intervention

Author	Intervention description	Trainer education	Training material
A. Technique training			
Jump landing performance—instruction			
Almonroeder ⁶⁵	Verbal internal versus external focus instruction followed by drop landings		31 cm box
Benjaminse ⁵⁴	Verbal internal, external focus and expert video instruction followed by double-legged landings		Video, monitor, box, LESS
Chijiatsu ⁷⁹	Video landing instruction of own drop vertical jumps landings		Bicycle, box, monitor, camera
McNair ⁵⁶	Technical instruction, auditory cue, imagery on double leg landing		30 cm box
Milner ⁸⁰	Three adapted verbal instructions on countermovement jump landing techniques		–
Mizner ⁷⁰	Uninstructed drop jump followed by verbal instruction on landing technique		31 cm box
Tate ⁵²	Supervised countermovement jump training with instruction/information and home-based training		Mirror, camera, instruction
Turner ⁶⁴	Bilateral barefoot drop-landings with expert video instruction		30.5 cm box, video, monitor
Welling ⁵¹	Instruction on drop vertical jump followed by feedback (LESS score)		30 cm box, LESS, camera, monitor
Jump landing performance—feedback			
Cronin ⁸¹	Bilateral jump-landings while spiking a volleyball of a toss by coach followed by verbal feedback		Volleyball, volleyball net, cone
Etnoyer ⁵⁷	Self or combination feedback (LESS Score and expert video) of box drop jump		30 cm box, camera, monitor, video, LESS
Leonard ⁸²	Dyad or expert feedback during squat jump		Feedback checklist
Oñate ⁵⁵	Basketball rebounding task on jump-ball device with self/expert/combo feedback		Camera, monitor, checklist, jump-ball (patent pending) testing device
Shams ⁵³	Plyometric exercises combined with feedback or tape		Mirror, cone, wall, scripted cueing
Jump landing performance—skill training			
Shimokochi ⁶⁹	Three different single-legged drop landing styles		Boxes (30, 45 cm)
B. Dynamic Strengthening			
Plyometrics			
Dello Iacano ⁸³	Vertical/horizontal alternate 1-leg drop jumps, landing from top of a platform		25 cm box
Herrington ⁸⁴	Progressive jump-training programme		Wall
Makaruk ⁶²	Single or repeated jumps with weekly increasing sets or jump height of four different jump forms		Slat, boxes (20, 30, 40 cm), hurdles
Nagano ⁸⁵	Balance and jump training divided in technique and performance phases		Square balance board
Vescovi ⁴⁸	Sportsmetrics: Jump exercises with increasing intensity over 6 weeks		Cones, mattress
Combi strengthening			
Dai ⁸⁶	Jump-landing-jump task with resistance band above ankles joint		30 cm box, LifeLine Medison band
Peng ⁸⁷	Drop jumps with elastic band loads (0% and 20% body weight)		Boxes (40, 50 cm), elastic band
Stearns and Power ⁸⁸	Hip-focused training programme with increasing difficulty		BOSU and ball
Yang ⁶⁰	Hip extension training and plyometrics (eg, jumps in different directions)		Box, hurdles, medicine ball, foam roller, BOSU balance board and ball
Prevention programme			
Aerts ⁵⁸	Coach-supervised jump-landing prevention programme on jump-landing technique		Poster, DVD, mattress, wall, balls
Fox ⁵⁹	Down 2 Earth		Illustrated programme, ball
C. Static strengthening			

Continued

Table 3 Continued

Author	Intervention description	Trainer education	Training material
Core workout			
Araujo ⁸⁹	Increasing core exercises (eg, plank variants, crunches, Russian twists)		–
Warm-up programme			
Avedasian ⁹⁰	Two different warm up protocols		–
O'Malley ⁶¹	GAA-15 standardised warm-up programme		Training partner, programme
D. Balance Training			
Static balance training			
Silva ⁴⁹	Wobble board training		Wobble board, ball, balloon, wall
Dynamic balance training			
Letefatkar ⁶³	Weekly increasing perturbation drills with verbal instruction		BOSU, balance boards, TheraBand, weight scale

The trainer education box was tacked if: (1) a training brochure must be studied before training intervention, (2) an instruction video must be watched, (3) LESS must be assessed, (4) expert jump-landing technique must be performed.
 BOSU, BOSU Balance trainer; Down 2 Earth, netball-specific prevention programme; DVD, Digital Video Disk; GAA-15, Gaelic Athletic Association warm up programme; LESS, Landing Error Scoring System.

Fourteen studies required coach education, with three training interventions including a training brochure or poster with detailed information on the exercises.^{58 59 61} No information was provided on how long it takes to master these programmes. One training intervention introduced instructions via DVD for coaches,⁵⁸ three interventions required knowledge about LESS^{51 54 57} and one provided a checklist of jump landing performance for coaches.⁵⁵ Five exercises required knowledge and performance of correct jump landing.^{51 54 55 57 64}

Risk of bias in studies

The risk of bias of the included studies, sorted by study design, are shown in figure 3. Twenty-six studies showed an overall moderate risk of bias. Across the RCTs, higher risk of bias was due to lack of blinding of participants and researchers (11 studies), lack of a randomisation tool (3 studies), or due to missing information regarding a prespecified analysis plan that was finalised before unblinded outcome data were available




Figure 3 Risk of bias of the included studies sorted by study design.

for the analysis (4 studies). All pre–post and cross-over studies had a moderate risk of bias mostly due to missing follow-up outcome measurement on a second date (16 studies).

DISCUSSION

This systematic review and meta-analysis evaluated 31 training interventions, aimed at reducing biomechanical risk factors for lower extremity injuries during landing movements, and their practical implications for amateur sports. Four training types were portrayed: technique, dynamic or static strengthening, and balance training. All but two studies were able to significantly improve biomechanical risk factors in terms of GRF, joint angle, joint moment and LESS outcomes. Due to the large variation in outcome measures in the RCTs, only two meta-analyses could be performed. The meta-analysis did not find a significant effect of interventions on the GRF. Only a significant medium pooled effect of 0.77 ($p=0.01$) was found on improvement of the knee flexion angle during landing.^{51 52 57 58} Moreover, all interventions were practicable in terms of required material, coach education and time consumption (tables 2 and 3, online supplemental table 1).

Relevant intervention types

Technique training and dynamic strengthening improved the knee joint flexion angle in the meta-analysis and thus can be recommended interventions in amateur setting. These intervention types were accompanied by other improvements in biomechanical risk factors and were also successfully integrated into other study programmes in this review. First, technique training as external focus instruction resulted in a significant improvement in knee flexion angle⁶⁵ and LESS score⁵⁴ compared with internal focus instruction. The positive effect of external focus is based on the enhanced motor system to self-organise the movement, which will result in an improved movement pattern (ie, optimal kinetic solution), compared with internal focus.⁶⁵ These results are in line with the general positive effects of external focus instruction on motor learning and movement technique when compared with internal focus instruction.^{66 67} Furthermore, the review showed that one to three technique training session based on feedback or instruction improved knee flexion angle,⁵⁷ LESS score^{53 54} and reduced GRF.^{52 55} This rapid performance improvement after a maximum of three sessions seems to be useful for integrating this intervention into training routine.⁶⁸

Second, changing the unilateral landing technique to landing on the toes with the body leaning forward also resulted in a reduction of risk factors in three pre–post studies.^{64 69 70} Particularly, a toe landing pattern increased the knee flexion angle and reduced GRFs.⁶⁹ Whereas heel contact (dorsiflexion) at initial ground contact can be the cause of non-contact ACL injuries.⁷¹ Thus, analysing the athletes landing technique should be considered and

landing on the toe should be preferred when performing single or double leg landings.⁶⁹

Lastly, a warm-up programme including a combination of dynamic strengthening, technique and balance training was useful as injury prevention strategy.⁶¹ These findings are in line with the study of Herman *et al*, who showed the reduction of lower extremity injuries in amateur athletes using neuromuscular warm-up strategies combining, among others, dynamic strengthening and perturbation training.⁷²

Large variation of outcome measures

General consideration of biomechanical risk factors within the included studies showed large variation in outcome measures, which affected direct comparison of risk factors (eg, maximum knee flexion angle and knee flexion angle over time). Future studies should consider analysing jump landing movements using, for example, the LESS, which identifies movement deficiencies and poor landing technique, as it serves as a validated method and simplifies assessment in the field and direct comparison between studies.⁷³ Using the LESS score enables extraction of individual biomechanical risk factors from the score (eg, knee valgus) and comparison of these partial results between studies.⁷⁴ In addition, caution should be taken when using two-dimensional motion capture analysis while focusing on the dynamic knee valgus during landing movements, as it consists of proximal (hip adduction/internal rotation) and distal (tibial abduction) movement to the knee, which take place in two different planes of motion. Therefore, three-dimensional analysis represents the gold standard for kinematic analyses.⁷⁵

Gender differences

Regarding gender differences, 65% of the results were obtained for female samples. As women have been reported to be at higher risk for knee injuries, injury prevention is crucial for them. Female athletes reduced biomechanical risk through dynamic and static strengthening, technique training and dynamic balance training, like male athletes. A study by Crossley *et al* has shown that it is beneficial to use a multifaceted intervention programme (eg, strength, plyometric exercises, dynamic balance training) to reduce the rate of lower extremity injury in women.⁷⁶ The included studies support these findings and show significant reduction of biomechanical risk factors by applying a combination of plyometrics and strengthening exercise (ie, dynamic strengthening).⁵⁹

Practicability of intervention implementation

When referring to the practicability of the training intervention the results suggested a moderate time commitment (around 30 min), minimal training material and coach education, which make the interventions integrable into training routine (table 3 and online supplemental table 1). The examined interventions mainly used materials which are often part of the existing

training equipment, for example, hurdles and boxes or no additional material, and thus facilitate the application of the training programme.^{52 67 77} A barrier to implement the intervention in amateur sports could be the incurred costs for monitors or cameras that were used in some studies to provide feedback regarding jump landing performance. However, significant improvement in LESS score were also obtained when visual feedback was provided with mirrors, which drastically reduced costs if available.⁵³

Although good coach education reduces injury risk,⁷⁷ the present review demonstrated that targeted simple training interventions can significantly reduce biomechanical risk factors, thus circumventing the time and cost hurdle of continuing education in amateur sports. However, successful training intervention depend on programme adherence. Steffen *et al* reported a larger positive effect of FIFA 11+ when participants showed high adherence compared with low adherence athletes.⁷⁸

Limitations and strength of the review

As all research, this review was subject to limitations of the included studies. First, the unaccounted-for influence on outcomes, such as sample frequency, footwear, age and activity level of participants was acknowledged as limitations. Second, more than half of the results of this review came from the analysis of female athletes (65%). Therefore, caution is advised in the general transfer of results for male amateur sport athletes. Third, direct comparison of biomechanical risk factors between studies was difficult because the collection of these parameters in the studies was not uniform. Fourth, the meta-analysis, which showed a significant result of increasing knee flexion angle, only included four RCTs (figure 2). Furthermore, baseline values differed between groups and influenced the intervention effects.^{49 55} Nevertheless, risk of bias analyses of the 31 included studies showed a mostly moderate risk. Main reason for this was the missing follow-up outcome measurement of the pre-post and cross-over studies on a second date. Therefore, future studies should aim to perform a RCT to increase the quality of the results and add a follow-up measurement to reduce the risk of bias. So far, the results suggest that the interventions have a positive effect on the biomechanical risk factors directly post intervention, but it is not clear if these effects hold long-term if training is not performed or neglected.^{51 52 57 60}

The key strength of this review is the comprehensive literature search conducted in three databases, rigorous checking of the extracted data performed by two independent reviewers, and the additional risk of bias analyses carried out with the most reliable Cochrane Risk of Bias Tools (RoB2, ROBINS-I) performed independently by the main author and a third reviewer. In addition, 70% of the included studies analysed amateur athletes, so the transfer of the results to amateur sports is given.

CONCLUSION

This systematic review emphasises that implementing technique training in form of instruction or feedback, and dynamic strengthening in form of plyometrics, can improve biomechanical risk factors for lower extremity injuries during landing movements in adult amateur athletes. The meta-analysis showed that technique training (as stand-alone or in combination with plyometric training) has a significant positive effect on the knee flexion angle and should therefore be recommended. Further, the practicability in terms of limited additional required material, coach education and training duration has been demonstrated for amateur settings. Future studies should report more consistent parameters such as the LESS and should test the long-term effects of often temporary interventions.

Acknowledgements We would like to thank Tatiana Hazenberg for the linguistic revision.

Contributors CB and RD had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis. CB, LF and RD conceptualised and designed the initial study, drafted the initial manuscript, reviewed and revised the manuscript. CB and AG systematically screened the data and carried out the initial analyses. TH, LF and RD contributed to data screening and extraction, result analyses and discussion. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of work. RD accepts full responsibility for the finished work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Chantal Bathe <http://orcid.org/0000-0002-7837-5406>

REFERENCES

- Gimigliano F, Resmini G, Moretti A, *et al*. Epidemiology of musculoskeletal injuries in adult athletes: a scoping review. *Medicina (Kaunas)* 2021;57:1118.
- Stiftung Sicherheit im Sport. Vereinssport in Der kommune-MIT sicherheit verletzungsfrei-abschlussbericht 2019-theoretische aufarbeitung Der thematik und vorbereitung Der pilotkommune. 2019. Available: <https://www.sicherheit.sport/projekte/vereinssport-in-der-kommune3/>
- Gribble PA, Bleakley CM, Caulfield BM, *et al*. Evidence review for the 2016 international ankle consortium consensus statement on

- the prevalence, impact and long-term consequences of lateral ankle sprains. *Br J Sports Med* 2016;50:1496–505.
- 4 Magnussen RA, Meschbach NT, Kaeding CC, et al. ACL graft and contralateral ACL tear risk within ten years following reconstruction: a systematic review. *JBJS Rev* 2015;3:e3.
 - 5 Barber-Westin S, Noyes FR. One in 5 athletes sustain reinjury upon return to high-risk sports after ACL reconstruction: a systematic review in 1239 athletes younger than 20 years. *Sports Health* 2020;12:587–97.
 - 6 Harringe ML, Renström P, Werner S. Injury incidence, mechanism and diagnosis in top-level teamgym: a prospective study conducted over one season. *Scand J Med Sci Sports* 2007;17:115–9.
 - 7 Bates NA, Schilaty ND, Nagelli CV, et al. Multiplanar loading of the knee and its influence on anterior cruciate ligament and medial collateral ligament strain during simulated landings and noncontact tears. *Am J Sports Med* 2019;47:1844–53.
 - 8 De Bleeker C, Vermeulen S, De Blaiser C, et al. Relationship between jump-landing kinematics and lower extremity overuse injuries in physically active populations: a systematic review and meta-analysis. *Sports Med* 2020;50:1515–32.
 - 9 Aerts I, Cumps E, Verhagen E, et al. A systematic review of different jump-landing variables in relation to injuries. *J Sports Med Phys Fitness* 2013;53:509–19.
 - 10 Ameer MA, Muaidi QI. Influence of increasing knee flexion angle on knee-ankle varus stress during single-leg jump landing. *J Taibah Univ Med Sci* 2017;12:497–503.
 - 11 Blackburn JT, Norcross MF, Cannon LN, et al. Hamstrings stiffness and landing biomechanics linked to anterior cruciate ligament loading. *J Athl Train* 2013;48:764–72.
 - 12 Louw Q, Grimmer K, Vaughan C. Knee movement patterns of injured and uninjured adolescent basketball players when landing from a jump: a case-control study. *BMC Musculoskelet Disord* 2006;7:22.
 - 13 Lian Ø, Refsnes P-E, Engebretsen L, et al. Performance characteristics of volleyball players with patellar tendinopathy. *Am J Sports Med* 2003;31:408–13.
 - 14 Leppänen M, Pasanen K, Kujala UM, et al. Stiff landings are associated with increased ACL injury risk in young female basketball and floorball players. *Am J Sports Med* 2017;45:386–93.
 - 15 Olsen O-E, Myklebust G, Engebretsen L, et al. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med* 2004;32:1002–12.
 - 16 Beynnon B, Howe JG, Pope MH, et al. The measurement of anterior cruciate ligament strain in vivo. *Int Orthop* 1992;16:1–12.
 - 17 Bisseling RW, Hof AL, Bredeweg SW, et al. Are the take-off and landing phase dynamics of the volleyball spike jump related to patellar tendinopathy? *Br J Sports Med* 2008;42:483–9.
 - 18 Caulfield BM, Garrett M. Functional instability of the ankle: differences in patterns of ankle and knee movement prior to and post landing in a single leg jump. *Int J Sports Med* 2002;23:64–8.
 - 19 Pollard CD, Sigward SM, Powers CM. Limited hip and knee flexion during landing is associated with increased frontal plane knee motion and moments. *Clin Biomech (Bristol, Avon)* 2010;25:142–6.
 - 20 Hewett TE, Ford KR, Hoogenboom BJ, et al. Understanding and preventing ACL injuries: current biomechanical and epidemiologic considerations-update 2010. *N Am J Sports Phys Ther* 2010;5:234–51.
 - 21 Yu B, Lin C-F, Garrett WE. Lower extremity biomechanics during the landing of a stop-jump task. *Clin Biomech (Bristol, Avon)* 2006;21:297–305.
 - 22 Alzahrani AM, Alzhrani M, Alshahrani SN, et al. Is hip muscle strength associated with dynamic knee valgus in a healthy adult population? A systematic review. *Int J Environ Res Public Health* 2021;18:7669.
 - 23 Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med* 2005;33:492–501.
 - 24 Aerts I, Cumps E, Verhagen E, et al. Efficacy of a 3 month training program on the jump-landing technique in jump-landing sports. design of a cluster randomized controlled trial. *BMC Musculoskelet Disord* 2010;11:281.
 - 25 Hadzic V, Sattler T, Topole E, et al. Risk factors for ankle sprain in volleyball players: a preliminary analysis. *IES* 2009;17:155–60.
 - 26 Willems TM, Witvrouw E, Delbaere K, et al. Intrinsic risk factors for inversion ankle sprains in male subjects: a prospective study. *Am J Sports Med* 2005;33:415–23.
 - 27 Parkkari J, Kujala UM, Kannus P. Is it possible to prevent sports injuries? Review of controlled clinical trials and recommendations for future work. *Sports Med* 2001;31:985–95.
 - 28 Barengo NC, Meneses-Echávez JF, Ramírez-Vélez R, et al. The impact of the FIFA 11+ training program on injury prevention in football players: a systematic review. *Int J Environ Res Public Health* 2014;11:11986–2000.
 - 29 Sadiqursky D, Braid JA, De Lira DNL, et al. The FIFA 11+ injury prevention program for soccer players: a systematic review. *BMC Sports Sci Med Rehabil* 2017;9:18.
 - 30 Aerts I, Cumps E, Verhagen E, et al. A 3-month jump-landing training program: a feasibility study using the RE-AIM framework. *J Athl Train* 2013;48:296–305.
 - 31 Gianotti S, Hume PA, Tunstall H. Efficacy of injury prevention related coach education within netball and soccer. *J Sci Med Sport* 2010;13:32–5.
 - 32 Herman DC, Oñate JA, Weinhold PS, et al. The effects of feedback with and without strength training on lower extremity biomechanics. *Am J Sports Med* 2009;37:1301–8.
 - 33 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
 - 34 Ardern CL, Büttner F, Andrade R, et al. Implementing the 27 PRISMA 2020 statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the persist (implementing PRISMA in exercise, rehabilitation, sport medicine and sports science) guidance. *Br J Sports Med* 2022;56:175–95.
 - 35 Aggarwal R, Ranganathan P. Study designs: part 4 – interventional studies. *Perspect Clin Res* 2019;10:137.
 - 36 Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5:210.
 - 37 Higgins JPT, Cochrane Collaboration. *Cochrane handbook for systematic reviews of interventions. Second edition.* Hoboken, NJ: Wiley-Blackwell, 2020.
 - 38 Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;i4919.
 - 39 Harrer M, Cuijpers P, Furukawa TA, et al. *Doing meta-analysis with R: a hands-on guide. 1st ed.* Boca Raton, FL and London: Chapman & Hall/CRC Press, 2021.
 - 40 Cuijpers P, Weitz E, Cristea IA, et al. Pre-post effect sizes should be avoided in meta-analyses. *Epidemiol Psychiatr Sci* 2017;26:364–8.
 - 41 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
 - 42 Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med* 2003;22:2693–710.
 - 43 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
 - 44 Harrer M, Adam SH, Messner E-M, et al. Prevention of eating disorders at universities: a systematic review and meta-analysis. *Int J Eat Disord* 2020;53:813–33.
 - 45 Viechtbauer W, Cheung M-L. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods* 2010;1:112–25.
 - 46 Egger M, Smith GD, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
 - 47 Cohen J. *Statistical power analysis for the behavioral sciences. 2nd ed.* Hillsdale, N J: L Erlbaum Associates, 1988.
 - 48 Vescovi JD, Canavan PK, Hasson S. Effects of a plyometric program on vertical landing force and jumping performance in college women. *Phys Ther Sport* 2008;9:185–92.
 - 49 Silva PB, Oliveira AS, Mrachacz-Kersting N, et al. Effects of wobble board training on single-leg landing neuromechanics. *Scand J Med Sci Sports* 2018;28:972–82.
 - 50 Sibbald B, Roland M. Understanding controlled trials. why are randomised controlled trials important? *BMJ* 1998;316:201.
 - 51 Welling W, Benjaminse A, Gokeler A, et al. Retention of movement technique: implications for primary prevention of ACL injuries. *Int J Sports Phys Ther* 2017;12:908–20.
 - 52 Tate JJ, Milner CE, Fairbrother JT, et al. The effects of a home-based instructional program aimed at improving frontal plane knee biomechanics during a jump-landing task. *J Orthop Sports Phys Ther* 2013;43:486–94.
 - 53 Shams F, Hadadnezhad M, Letafatkar A, et al. Valgus control feedback and taping improves the effects of plyometric exercises in women with dynamic knee valgus. *Sports Health* 2021;14:747–57.
 - 54 Benjaminse A, Welling W, Otten B, et al. Transfer of improved movement technique after receiving verbal external focus and video instruction. *Knee Surg Sports Traumatol Arthrosc* 2018;26:955–62.
 - 55 Oñate JA, Guskievicz KM, Marshall SW, et al. Instruction of jump-landing technique using videotape feedback: altering lower extremity motion patterns. *Am J Sports Med* 2005;33:831–42.
 - 56 McNair PJ, Prapavessis H, Callender K. Decreasing landing forces: effect of instruction. *Br J Sports Med* 2000;34:293–6.

- 57 Ethoyer J, Cortes N, Ringleb SI, *et al.* Instruction and jump-landing kinematics in college-aged female athletes over time. *J Athl Train* 2013;48:161–71.
- 58 Aerts I, Cumps E, Verhagen E, *et al.* The effect of a 3-month prevention program on the jump-landing technique in basketball: a randomized controlled trial. *J Sport Rehabil* 2015;24:21–30.
- 59 Fox AS, Bonacci J, McLean SG, *et al.* Exploring individual adaptations to an anterior cruciate ligament injury prevention programme. *Knee* 2018;25:83–98.
- 60 Yang C, Yao W, Garrett WE, *et al.* Effects of an intervention program on lower extremity biomechanics in stop-jump and side-cutting tasks. *Am J Sports Med* 2018;46:3014–22.
- 61 O'Malley E, Murphy JC, McCarthy Persson U, *et al.* The effects of the gaelic athletic association 15 training program on neuromuscular outcomes in gaelic football and hurling players: a randomized cluster trial. *J Strength Cond Res* 2017;31:2119–30.
- 62 Makaruk H, Czaplicki A, Saczewicz T, *et al.* The effects of single versus repeated plyometrics on landing biomechanics and jumping performance in men. *Biol Sport* 2014;31:9–14.
- 63 Letafatkar A, Rajabi R, Minoonejad H, *et al.* Efficacy of perturbation-enhanced neuromuscular training on hamstring and quadriceps onset time, activation and knee flexion during a tuck-jump task. *Int J Sports Phys Ther* 2019;14:214–27.
- 64 Turner C, Crow S, Crowther T, *et al.* Preventing non-contact ACL injuries in female athletes: what can we learn from dancers? *Phys Ther Sport* 2018;31:1–8.
- 65 Almonroeder TG, Jayawickrema J, Richardson CT, *et al.* The influence of attentional focus on landing stiffness in female athletes: a cross-sectional study. *Int J Sports Phys Ther* 2020;15:510–8.
- 66 Benjaminse A, Welling W, Otten B, *et al.* Novel methods of instruction in ACL injury prevention programs, a systematic review. *Phys Ther Sport* 2015;16:176–86.
- 67 Wulf G. Attentional focus and motor learning: a review of 15 years. *Int Rev Sport Exerc Psychol* 2013;6:77–104.
- 68 Crowell HP, Milner CE, Hamill J, *et al.* Reducing impact loading during running with the use of real-time visual feedback. *J Orthop Sports Phys Ther* 2010;40:206–13.
- 69 Shimokochi Y, Ambegaonkar JP, Meyer EG. Changing sagittal-plane landing styles to modulate impact and tibiofemoral force magnitude and directions relative to the tibia. *J Athl Train* 2016;51:669–81.
- 70 Mizner RL, Kawaguchi JK, Chmielewski TL. Muscle strength in the lower extremity does not predict postinstruction improvements in the landing patterns of female athletes. *J Orthop Sports Phys Ther* 2008;38:353–61.
- 71 Hewett TE, Torg JS, Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br J Sports Med* 2009;43:417–22.
- 72 Herman K, Barton C, Malliaras P, *et al.* The effectiveness of neuromuscular warm-up strategies, that require no additional equipment, for preventing lower limb injuries during sports participation: a systematic review. *BMC Med* 2012;10:75.
- 73 Hanzlíková I, Hébert-Losier K. Is the landing error scoring system reliable and valid? A systematic review. *Sports Health* 2020;12:181–8.
- 74 Hanzlíková I, Athens J, Hébert-Losier K. Clinical implications of landing error scoring system calculation methods. *Phys Ther Sport* 2020;44:61–6.
- 75 Holden S, Boreham C, Doherty C, *et al.* Two-dimensional knee valgus displacement as a predictor of patellofemoral pain in adolescent females. *Scand J Med Sci Sports* 2017;27:188–94.
- 76 Crossley KM, Patterson BE, Culvenor AG, *et al.* Making football safer for women: a systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. *Br J Sports Med* 2020;54:1089–98.
- 77 Bianchi FP, Labbate M, Castellana M, *et al.* Epidemiology of injuries among amateur athletes who attended fitness activities: the role of the qualification of the trainer. *J Sports Med Phys Fitness* 2020;60:422–7.
- 78 Steffen K, Emery CA, Romiti M, *et al.* High adherence to a neuromuscular injury prevention programme (FIFA 11+) improves functional balance and reduces injury risk in Canadian youth female football players: a cluster randomised trial. *Br J Sports Med* 2013;47:794–802.
- 79 Chijimatsu M, Ishida T, Yamanaka M, *et al.* Landing instructions focused on pelvic and trunk lateral tilt decrease the knee abduction moment during a single-leg drop vertical jump. *Physical Therapy in Sport* 2020;46:S1466-853X(20)30533-2:226–33..
- 80 Milner CE, Fairbrother JT, Srivatsan A, *et al.* Simple verbal instruction improves knee biomechanics during landing in female athletes. *Knee* 2012;19:399–403.
- 81 Cronin JB, Bressel E, Fkinn L. Augmented feedback reduces ground reaction forces in the landing phase of the volleyball spike jump. *Journal of Sport Rehabilitation* 2008;17:148–59.
- 82 Leonard KA, Simon JE, Yom J, *et al.* The immediate effects of expert and Dyad external focus feedback on drop landing Biomechanics in female athletes: an Instrumented field study. *International Journal of Sports Physical Therapy* 2021;16:96–105.
- 83 Dello Iacono A, Martone D, Milic M, *et al.* Vertical- vs. horizontal-oriented drop jump training: chronic effects on explosive performances of elite Handball players. *Journal of Strength and Conditioning Research* 2017;31:921–31.
- 84 Herrington L. The effects of 4 weeks of jump training on landing knee Valgus and crossover hop performance in female basketball players. *Journal of Strength and Conditioning Research* 2010;24:3427–32.
- 85 Nagano Y, Ida H, Akai M, *et al.* Effects of jump and balance training on knee Kinematics and Electromyography of female basketball athletes during a single limb drop landing: pre-post intervention study. *Sports Medicine, Arthroscopy, Rehabilitation, Therapy & Technology* 2011;3:14.
- 86 Dai B, Garrett WE, Gross MT, *et al.* The effects of 2 landing techniques on knee Kinematics, Kinetics, and performance during stop-jump and side-cutting tasks. *The American Journal of Sports Medicine* 2015;43:466–74.
- 87 Peng H-T, Song C-Y, Chen Z-R, *et al.* Effects of attaching elastic bands to the waist and heels on drop jumps. *European Journal of Sport Science* 2022;22:808–16.
- 88 Stearns KM, Powers CM. Improvements in hip muscle performance result in increased use of the hip Extensors and Abductors during a landing task. *The American Journal of Sports Medicine* 2014;42:602–9.
- 89 Araujo S, Cohen D, Hayes L. Six weeks of core stability training improves landing Kinetics among female Capoeira athletes: a pilot study. *Journal of Human Kinetics* 2015;45:27–37.
- 90 Avedesian JM, Judge LW, Wang H, *et al.* The biomechanical effect of warm-up stretching strategies on landing mechanics in female volleyball athletes. *Sports Biomech* 2020;19:587–600.