

Untangling risk factors including discipline-specific exposure for injuries in preprofessional and professional circus artists in the USA

Stephanie Greenspan ^{1,2}, Melanie I Stuckey ³

To cite: Greenspan S, Stuckey MI. Untangling risk factors including discipline-specific exposure for injuries in preprofessional and professional circus artists in the USA. *BMJ Open Sport & Exercise Medicine* 2023;**9**:e001551. doi:10.1136/bmjsem-2023-001551

Accepted 24 April 2023

ABSTRACT

Objective This prospective cohort study quantified injury patterns related to specific circus discipline exposure in preprofessional and professional circus artists.

Methods Circus artists (n=201; ages 13–69; 122 female, 29 male assigned sex at birth) were enrolled in 10 cities across the USA. Participants were followed for 1 year from enrolment, completing a weekly training log and undergoing a physical therapist evaluation for injuries. The circus-specific extension of the International Olympic Committee 2020 consensus on recording injury and illness in sports was used to analyse injury patterns.

Results The study completion rate was 77% (n=155). Data were analysed by participant subgroup (age, professional status, sex at birth). The highest injury rates in participant subgroups were for males (5.69/1000 exposures) and related to discipline subgroups, were in aerial with ground elements (5.93/1000 exposures) and aerial (4.26/1000 exposures). Adults had more injuries related to aerial, whereas adolescents had more related to ground disciplines (χ^2 (2)=10.62, p=0.005) and non-time loss injuries (χ^2 (1)=5.45, p=0.02). Females had a higher proportion of repetitive injuries (70% vs 55%) than males (χ^2 (1) = 4.43, p=0.035). Individuals with an eating disorder history had more (p<0.004) injuries (mean 2.27±2.29) than those without (mean=1.48±0.96).

Conclusions This study showed that intrinsic factors (age, sex at birth and history of eating disorder) and extrinsic factors (circus discipline exposure) affect injury risk. We need to account for the intersectionality of these factors to address risk management at an individual and group level.

INTRODUCTION

Injury prevention is a key goal for artistic or athletic recreations or occupations, including circus arts. The first step for developing an injury prevention strategy includes surveillance to understand how personal (intrinsic) and environmental (extrinsic) factors affect injury risk.¹ Circus injury research has been limited to primarily descriptive studies of professional artists in a large company² and students at professionalising schools.^{3–7} These populations represent a small segment of the

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Circus participation involves injury risk as an athletic performing art. Injury research in the circus arts is limited to professionalising students and performers in large companies, and little is understood about the influence of different circus disciplines or contexts on injury risk.

WHAT THIS STUDY ADDS

⇒ This study examines how age, professional status, sex at birth and different circus disciplines are related to injury patterns. It is the first circus study to integrate the circus-specific extension framework of the International Olympic Committee 2020 consensus into reporting injuries.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This study can inform areas of focus for research and practice for injury prevention strategies in circus arts.

overall circus context in the USA.⁸ Training or performing environments such as preprofessional programmes, professional freelance work, or small, low-resourced circus companies with different extrinsic factors may impact injury patterns. Research is needed across other settings to broaden the understanding of injury risk for circus arts.

Circus disciplines pose specific extrinsic factors regarding physical, technical and artistic demands, which likely impact injury risk, yet differences have not been explored.^{9–11} Early research compared the anatomical location of injuries for floor acrobatics versus acrobatics with equipment, including aerial (trapeze) and ground (tight-wire and slackline) disciplines with inconsistent biomechanical demands.³ A recent study found clinical injury burden (injury rate plotted against injury duration) was generally higher in ground acrobats, but the exposure was not considered.⁷ Two studies^{4,12} reported higher proportions of injuries for ground acrobatics,



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Physical Therapy, Samuel Merritt University, Oakland, CA, USA

²Artistic Science, Oakland, CA, USA

³Centre de recherche, d'innovation et de transfert en arts du cirque, École nationale de cirque, Montréal, Quebec, Canada

Correspondence to

Dr Stephanie Greenspan; sgreenspan@samuelmerritt.edu



but the exposure was also higher for these disciplines, which may have accounted for the difference. In contrast, one study showed similar 1-year injury prevalence in aerialists and non-aerialists (73% vs 81%).⁶ Still, injury rates were based on the student's primary discipline and not necessarily the discipline attributed to injury.⁶ A comprehensive understanding of injury risk in circus discipline subgroups with similar demands (eg, aerial vs ground acrobatics) is lacking.

The purpose of this study was to characterise injuries and exposure in preprofessional and professional circus artists in the USA. Secondly, the intent was to identify injury patterns by extrinsic (circus discipline subgroup) and intrinsic (age, sex at birth (SAB), medical history and prior health status) using the framework from the circus-specific extension of the International Olympic Committee 2020 consensus on recording injury and illness in sport.^{9 13}

METHODS

Participants

Ten circus training facilities across the USA were selected as host sites based on the size of the eligible target population, the presence of long-term intensive training programmes, and willingness to assist with study recruitment. Participants were recruited via flyers at area training centres, social media, American Circus Educator newsletter announcements, emails to host students and staff, and other local circus companies. Eligible participants were preprofessional (training ≥ 6 hours a week and performing in ≥ 2 shows per year) and professional (self-identified) circus artists over age 13 years, able to read/comprehend English and fulfil the requirements of the study. Exclusion criteria included a planned absence from the area ≥ 1 month over the next year. A power analysis showed that 200 was an adequate sample size for a medium-to-large (0.25–0.50) effect size with a *p* value of 0.05. Adults signed an informed consent form, and adolescents signed an assent form. Parents/guardians completed informed consent.

Design

The prospective, observational cohort design was modelled after a pilot study.^{11 12} Participants/public were not involved in the study design, conduct or dissemination.

Rolling enrolment occurred from September to December 2018 (cohort 1, four facilities) and from September 2019 to January 2020 (cohort 2, six facilities). Enrolment included online completion of informed consent/assent forms, an intake questionnaire including demographics, training experience and medical history, and a physical examination^{11 12} of intrinsic factors by a physical therapist (PT). PTs had at least 5 years of orthopaedic or sports-related experience and underwent standardised training for study procedures.

Participants were tracked for 1 year following their enrolment date. Participants were emailed a Qualtrics

(V.9/2017–2012/2018, Provo, Utah, USA) survey link to complete a weekly training log where they reported weekly exposure as the total number of sessions (classes, rehearsals, performances or self-directed training by circus discipline and strength or flexibility conditioning). New or ongoing injuries and missed training sessions due to injury were recorded. For weeks without exposure, participants indicated the reason (eg, vacation, illness, COVID-19 closure). For new or recurrent injuries, the PTs conducted an interview and physical examination to determine the injury's mechanism, body region, tissue and nature. Injuries at study enrolment were not included in the injury frequency calculation unless an exacerbation occurred.

Injury classification

The injury was defined as an anatomical tissue-level impairment.^{11 12 14} Time loss injury (TL) resulted in the full loss of participation in at least one circus discipline for one or more days after injury onset.^{11 12 14} Injuries not meeting the TL criteria were defined as non-time loss (NTL). The type of injury and mode of onset followed the circus-specific extension.^{9 13} New injury represented the first time a specific injury occurred within the study period.^{9 13 15 16} Subsequent recurrent injury was the same location and nature, and subsequent local injury was the same location but different nature after returning to 100% participation from TL injury.^{9 13 15 16} Acute sudden onset related to a single traumatic event (eg, fall, awkward landing).^{9 13} Repetitive sudden or gradual onset was associated with repetitive use.^{9 13 17} Injuries were defined as directly associated with participation in circus training, rehearsal or performance, or indirectly associated if related to the circus context outside of those activities (eg, spotting), or not related if outside of the circus context.^{9 13} Disciplines associated with injuries were grouped using the circus arts discipline subgroup classification system, which classified disciplines as aerial acrobatics (aerial, aerial with ground elements), ground acrobatics (human propulsion, apparatus propulsion, balance/control), manipulation and character.^{9 11}

Data analysis

Descriptive statistics were conducted using Microsoft Excel 365 (vV.2211, Redmond, Washington, USA) for baseline intake and examination data, injury rate (frequency per 1000 sessions of circus exposure) and frequency for participant subgroups by age, SAB and professional status. Brief descriptive statistics were also included for the non-binary participants to provide some information on this under-reported group. The remaining analyses used IBM SPSS Statistics (V.26) with significance set at *p*<0.05. A regression analysis used total training sessions, age and years of circus experience in years as predictor variables for injury incidence. Multiple *t*-tests compared injury types by discipline subgroups, age, professional status and SAB. Pearson's correlational analysis tested the relationship between the duration

of training experience and injuries. The influence of medical history and prior health status was tested using multiple regression analyses.

RESULTS

Two hundred and one participants (ages 13–69 years) enrolled, and 155 (77.11%) completed the study. Drop-outs (1 adolescent; 45 adults) resulted from non-adherence to weekly training log completion ($n=32$), personal issues, illness or moving away. Drop-outs' data were included for the duration of their participation. Participants were identified by SAB (172 assigned female at birth, AFAB; 29 assigned male at birth, AMAB) for the primary analysis due to a low number identifying as non-binary ($n=8$). Non-binary participants included six preprofessionals and two professionals (ages 24–36 years), with seven completing the study.

Participant characteristics

Select data from study enrolment are in [table 1](#). Average years of circus experience was higher ($t(195)=-3.96$, $p<0.001$), accounting for different variances ($F=7.26$, $p<0.008$), in professionals (8.63 years; $SD=6.01$) compared with preprofessionals (5.58 years; $SD=3.23$). Amenorrhoea was more prevalent in adolescents than adults and preprofessionals than professionals. No adolescents reported an eating disorder, although 36% of adolescent AFAB had amenorrhoea. Fewer participants reported a connective tissue disorder than met the criteria for generalised joint hypermobility (GJH) per their Beighton score.¹⁸

Exposure

Circus exposure by discipline subgroup and training context, excluding drop-outs, are in [tables 2 and 3](#). Weekly circus exposure was higher ($t(199)=10.76$, $p<0.001$) in professionals (9.27; $SD=1.44$ sessions) than preprofessionals (6.87; $SD=1.65$ sessions). Across participant subgroups, exposure was highest for aerial and ground (balance/control), except for adolescents with similar exposure to ground (human propulsion).

Injury rates

Injury rates, excluding drop-outs, are reported per 1000 exposures in [table 4](#). By participant subgroup, AMAB had the highest injury rates (5.69/1000 sessions), and by discipline subgroup, aerial with ground elements, aerial then ground (balance/control) were highest (5.93, 4.26, 3.66/1000 discipline-specific sessions).

Injury types or patterns

[Tables 5 and 6](#) show injury distribution by subgroup. The proportion of injuries was highest for the shoulder (22%), followed by the lumbosacral region (13%) and then the elbow (10%). Muscle/tendon injuries were most common (52%) across tissue types.

Age, sex, professional status and discipline

Adults had more aerial-related injuries compared with adolescents. In contrast, adolescents had more ground-related injuries ($\chi^2(2)=10.62$, $p=0.005$) and non-time loss injuries ($\chi^2(1)=5.45$, $p=0.02$). There were no differences between participant groups for 1-year injury incidence but a trend towards a correlation between increased age and decreased injury incidence ($r=-0.122$, $p=0.085$) was found. No significant differences existed by age or SAB for new versus subsequent injuries, body region or tissue type affected. There was a difference by SAB such that AFAB had a higher proportion of repetitive injuries than AMAB (70% vs 55%) and a lower proportion of acute (30% vs 43%) injuries ($\chi^2(1)=4.43$, $p=0.035$). There were no differences between injuries related to aerial versus ground disciplines or by professional status for body region, tissue type affected or mode of onset. TL in days across all participants excluding dropouts was higher for AFAB (21.79; $SD=54.96$) than AMAB (12.57; $SD=23.45$), professionals (29.88; $SD=70.66$) compared with preprofessionals (15.04; $SD=33.61$) and adults (20.86; $SD=51.79$) vs adolescents (17.53; $SD=50.62$). Across participants with TL injuries (excluding dropouts), adolescents had greater average TL over one year compared to adults (65.8; $SD=50.62$; 30.8; $SD=59.93$). Among the 8 non-binary participants, 14 injuries occurred in 6 participants. These included four shoulder, three thigh injuries and one each neck, abdominal, hip, ankle, foot, wrist and elbow injury.

Medical history and physical status

Individuals with an eating disorder history averaged more injuries (2.27; $SD=2.29$) than those without (1.48; $SD=0.96$; $p<0.004$). There were no significant differences in injury incidence between AFAB with amenorrhoea or participants with a history of connective tissue, musculoskeletal, neurological disorder or head injury/concussion compared with those without, or based on baseline physical characteristics per the enrolment physical examination that included selected measures of strength, flexibility, joint mobility and balance.

COVID-19 effects

The COVID-19 pandemic impacted cohort 2. Training facility and show closures started 1–7 months into their study period. Overall training and performance exposure were lower by 21.83% and 34.53% compared with cohort 1. There were changes in the disciplines trained due to facility access. The environment for training often changed from facility to home or outdoors, and performances became recorded or virtual. Injury assessments were completed virtually with a modified physical examination, sometimes limiting our ability to specify the affected body tissue.

DISCUSSION

This was the first multicentred, prospective longitudinal study to examine injury patterns for preprofessional and

Table 1 Participant demographics (entire cohort)

	Entire cohort			Subgroups by age			Subgroups by level			Subgroups by sex at birth	
	All	Adolescent	Adult	Preprofessional	Professional	Female	Male*				
No of participants	201	16	185	130	71	172	29				
Age	31.40 (8.91)	14.50 (1.51)	32.87 (7.68)	30.45 (9.97)	33.13 (6.26)	31.22 (8.60)	32.50 (10.72)				
Years of circus experience	6.98 (4.17)	5.88 (2.73)	7.07 (4.27)	5.71 (3.50)	9.27 (4.34)	7.01 (3.96)	6.75 (5.38)				
Height (cm)	164.37 (7.96)	164.30 (7.22)	164.38 (8.04)	164.22 (7.68)	164.65 (8.50)	162.85 (6.98)	173.69 (7.34)				
Mass (kg)	62.48 (10.90)	57.64 (9.90)	62.90 (10.91)	62.57 (11.49)	62.32 (9.82)	60.62 (9.12)	73.95 (13.81)				
History of eating disorder	18.00%	0.00%	19.57%	17.83%	18.31%	20.35%	3.57%				
History of amenorrhoea†	NA	35.71%	16.46%	21.05%	12.07%	18.02%	NA				
History of musculoskeletal injury	89.50%	75.00%	90.76%	87.60%	92.96%	90.12%	85.71%				
History of concussion/head injury	30.50%	12.50%	32.07%	21.71%	46.48%	29.07%	39.29%				
History of connective tissue disorder	1.50%	6.25%	1.09%	1.55%	1.41%	1.16%	3.57%				
Positive Beighton score‡	34.50%	31.25%	34.78%	31.78%	39.44%	37.21%	17.86%				

Measures are reported as mean (SD) or percentage (%).

*One male preprofessional participant did not complete a medical history questionnaire, so percentages were adjusted.

†History of amenorrhoea excludes males and accounts for the onset of puberty and the use of birth control.

‡Positive Beighton score as criteria for generalised joint hypermobility was $\geq 6/9$ in prepubertal children, $\geq 5/9$ for postpubertal up to age 50 and $\geq 4/9$ for over age 50.¹⁸

NA, not available;

Table 2 Average total and weekly session exposure to disciplines subgroups or conditioning (excludes drop-outs)

	All	Adolescent	Adult	Preprofessional	Professional	Female	Male
Total sessions/participant							
All	403.13	411.73	402.21	357.25	482.00	401.15	415.07
Aerial	130.16	71.87	136.41	103.45	176.09	122.34	177.43
Aerial with ground elements	29.36	5.13	31.96	31.37	25.91	31.99	13.45
Ground (human propulsion)	47.02	81.93	43.28	43.64	52.82	46.94	47.50
Ground (apparatus propulsion)	3.10	3.47	3.06	2.00	4.98	3.15	2.77
Ground (balance/control)	100.42	121.40	98.18	84.20	128.32	101.92	91.36
Manipulation	19.08	43.67	16.45	20.01	17.49	18.00	25.64
Character	0.38	0.00	0.42	0.47	0.23	0.44	0.00
Conditioning—flexibility	0.35	0.00	0.39	0.40	0.26	0.41	0.00
Conditioning—strength	73.25	84.27	72.07	71.71	75.89	75.95	56.91
Weekly sessions/participant							
All	7.75	7.92	7.73	6.87	9.27	7.71	7.98
Aerial	2.50	1.38	2.62	1.99	3.39	2.35	3.41
Aerial with ground elements	0.56	0.10	0.61	0.60	0.50	0.62	0.26
Ground (human propulsion)	0.90	1.58	0.83	0.84	1.02	0.90	0.91
Ground (apparatus propulsion)	0.06	0.07	0.06	0.04	0.10	0.06	0.05
Ground (balance/control)	1.93	2.33	1.89	1.62	2.47	1.96	1.76
Manipulation	0.37	0.84	0.32	0.38	0.34	0.35	0.49
Character	0.01	0.00	0.01	0.01	0.00	0.01	0.00
Conditioning—flexibility	0.01	0.00	0.01	0.01	0.01	0.01	0.00
Conditioning—strength	1.41	1.62	1.39	1.38	1.46	1.46	1.09

professional circus artists in the USA by intrinsic (age, SAB, medical history, physical characteristics) and extrinsic (circus discipline) risk factors. Injury risk is multifactorial and environmental factors are an important consideration.¹⁹ Many contextual differences were apparent

in our sample compared with previous studies. Circus students at professionalising schools trained 19 or more hours per week,^{3–5,7} whereas preprofessionals in this study trained 6.68 sessions per week on average. Sessions had close to a 1:1 relationship to hours in the pilot study.¹² In

Table 3 Average total and weekly session exposure by context (excludes drop-outs)

	All	Adolescent	Adult	Preprofessional	Professional	Female	Male
Total sessions/participant							
All	403.13	411.73	402.21	357.25	482	401.15	415.07
Classes	125.07	216.4	115.29	151.53	79.58	127.35	111.27
Additional training	130.59	115.87	132.17	120.96	147.16	126.14	157.52
Rehearsal	31.08	33.13	30.86	19.33	51.3	32.01	25.5
Performance	15.03	22.67	14.21	9.89	23.86	15.87	9.91
Teach	101.35	23.67	109.68	55.55	180.11	99.78	110.86
Weekly sessions/participant							
All	7.75	7.92	7.73	6.87	9.27	7.71	7.98
Classes	2.41	4.16	2.22	2.91	1.53	2.45	2.14
Additional training	2.51	2.23	2.54	2.33	2.83	2.43	3.03
Rehearsal	0.6	0.64	0.59	0.37	0.99	0.62	0.49
Performance	0.29	0.44	0.27	0.19	0.46	0.31	0.19
Teach	1.95	0.46	2.11	1.07	3.46	1.92	2.13

**Table 4** Injury rates by subgroup (excludes drop-outs)

Subgroup	Injury rate
All	3.82
Adolescent	2.42
Adult	3.98
Preprofessional	4.08
Professional	3.49
Female	3.5
Male	5.69
Aerial	4.26
Aerial with ground elements	5.93
Ground (human propulsion)	2.2
Ground (apparatus propulsion)	2.08
Ground (balance/control)	3.66
Manipulation	0.33
Character	0

For participant subgroups, the injury rate is per 1000 exposures (sessions of all circus training and performances). For circus discipline, the subgroup injury rate is the number of discipline subgroup-related injuries per 1000 exposures only in the same circus discipline subgroup.

large circus companies with resident shows, professionals may work 300 or more shows annually,¹⁰ compared with our professionals, who averaged 23.86 performances annually by discipline (cohort 1=29.94). Further, professionalising schools and large companies,^{2-5 7} had dedicated medical staff, and our training facilities did not, which may impact injury management. Finally, our sex distribution differed from previous studies, with 80% compared with 29%–59%²⁻⁷ AFAB. This study demonstrates the importance of describing and accounting for contextual differences in circus injury research to ensure injury risk is understood across the breadth of settings.

Discipline

Determining extrinsic risk for circus disciplines, including accounting for exposure and artists training multiple disciplines, is challenging⁹⁻¹² but important to inform injury prevention practices. We found higher injury rates attributed to aerial versus ground acrobatics, but rates varied across discipline subgroups. Aerial with ground elements, including Chinese pole and pole dance, had the highest injury rate (5.93/1000 session exposures). In a professional training programme, injuries from Chinese pole were most prevalent among aerial disciplines but less frequent than ground acrobatics.⁴ Injury rates for recreational pole dance were higher (7.65/1000 hours training)²⁰ than in circus arts (0.3–3.3/1000 hours).^{3 5 7} Although study methods and population differences limit comparison between studies, aerial with ground elements may have an elevated injury risk.

Ground acrobatics-related injury rates were highest for the balance/control subgroup (3.66/1000 sessions),

which includes contortion, hand-balancing and partner acrobatics. In contrast, Munro⁴ found ground acrobatics was the leading cause of injury with a similar proportion in balance/control and human propulsion disciplines. Exposure was not quantified, but the tumbling practice was a sizeable part of the school curriculum. Similarly, floor exercise is associated with the highest proportion of gymnastics injuries.^{21 22} Future studies should account for discipline exposure in injury rates to improve our understanding of injury risk across disciplines and to avoid misrepresenting injury risk by just reporting frequency.

Age

Age is an intrinsic risk factor. Like the pilot study,¹² adults had a higher injury rate than adolescents (3.98 vs 2.42/1000 session exposures), although incidence decreased with age. Adolescents had more ground than aerial-related injuries (60%;14%) but also higher exposure to the ground than aerial (3.98; 1.48 sessions/week). Adults showed the opposite pattern with lower injury frequency (37%;44%) and exposure (2.78; 3.23) in ground than aerial disciplines, highlighting the importance of interpreting injury risk relative to exposure. Adolescents had significantly more NTL injuries (77%) than adults (47%), suggesting that adults have more severe injuries. However, across participants with TL injuries (excluding dropouts), adolescents had greater average TL over one year (65.8; SD=50.62; 30.8; SD=59.93). Accounting for injury management might be useful to understand this contrast as high school circus arts students were less likely than basketball players to report the injury to the athletic trainer.²³ Adults may also be better at implementing rest or accessing healthcare resources.

Preprofessional versus professional

A high absolute workload can positively or negatively impact injury risk depending on ramp-up and internal factors such as stress, anxiety and fatigue.²⁴ Fatigue is a known injury risk factor in sport²⁵ and dance,^{26 27} and is affected by training load in circus students, despite sufficient sleep.²⁸ Professional circus artists reported less fatigue than students at a circus college,²⁹ which could explain why despite higher exposure (9.27 vs 6.87 sessions/week), professional circus artists experienced a lower injury rate than preprofessionals (3.49 vs 4.08/1000 sessions).

Injury rate differences may also relate to greater professional experience (9.27 years) than preprofessionals (5.71 years). Third-year students at a circus college experienced lower injury rates than first-year students suggesting that experience may be protective.⁷ This could be attributed to training adaptations, mental performance or focus on artistry and consistency for a marketable act rather than technical development and building a skill repertoire. Professional circus artists have also reported lower social isolation, better social skills, fewer problems in shows/evaluations and greater performance satisfaction

Table 5 Injury frequency (percentage) of injury types (entire cohort)

	All	Adolescent	Adult	Preprofessional	Professional	Female	Male
Total injuries	276	22 (8%)	254 (92%)	169 (61%)	107 (39%)	222 (80%)	54 (20%)
Time loss	115 (42%)	5 (23%)	110 (43%)	71 (42%)	44 (41%)	96 (43%)	19 (35%)
Non-time loss	161 (58%)	17 (77%)	144 (57%)	98 (58%)	63 (59%)	126 (57%)	35 (65%)
New	257 (93%)	20 (91%)	57 (22%)	156 (92%)	101 (94%)	205 (92%)	52 (96%)
Subsequent recurrent	9 (3%)	2 (9%)	7 (3%)	6 (4%)	3 (3%)	8 (4%)	2 (4%)
Subsequent local	10 (4%)	0	10 (4%)	7 (4%)	3 (3%)	9 (4%)	0
Acute—sudden	89 (32%)	7 (32%)	82 (32%)	57 (34%)	32 (30%)	66 (30%)	23 (43%)
Repetitive—sudden	34 (12%)	4 (18%)	30 (12%)	23 (14%)	11 (10%)	28 (13%)	6 (11%)
Repetitive—gradual	150 (54%)	14 (64%)	136 (54%)	89 (53%)	61 (57%)	126 (57%)	24 (44%)
Unknown	3 (1%)	1 (5%)	2 (1%)	1 (1%)	2 (2%)	3 (1%)	0
Head/face	1 (0%)	0	1 (0%)	1 (1%)	0	1 (0%)	0
Neck/cervical spine	15 (5%)	0	15 (6%)	7 (4%)	8 (7%)	11 (5%)	4 (7%)
Thoracic spine/upper back	10 (4%)	0	10 (4%)	5 (3%)	5 (5%)	8 (4%)	2 (4%)
Chest	10 (4%)	0	10 (4%)	4 (2%)	6 (6%)	8 (4%)	2 (4%)
Lumbosacral	35 (13%)	5 (23%)	30 (12%)	21 (12%)	14 (13%)	30 (14%)	5 (9%)
Abdomen	10 (4%)	1 (5%)	9 (4%)	8 (5%)	2 (2%)	8 (4%)	2 (4%)
Shoulder	61 (22%)	5 (23%)	56 (22%)	36 (21%)	25 (23%)	47 (21%)	14 (26%)
Upper arm	1 (0%)	1 (5%)	0	1 (1%)	0	1 (0%)	0
Elbow	27 (10%)	0	27 (11%)	17 (10%)	10 (9%)	21 (9%)	6 (11%)
Forearm	8 (3%)	0	8 (3%)	6 (4%)	2 (2%)	5 (2%)	3 (6%)
Wrist	21 (8%)	4 (18%)	17 (7%)	13 (8%)	8 (7%)	17 (8%)	4 (7%)
Hand	11 (4%)	1 (5%)	10 (4%)	9 (5%)	2 (2%)	9 (4%)	2 (4%)
Hip/groin	20 (7%)	1 (5%)	19 (7%)	12 (7%)	8 (7%)	19 (9%)	1 (2%)
Thigh	15 (5%)	1 (5%)	14 (6%)	8 (5%)	7 (7%)	11 (5%)	4 (7%)
Knee	15 (5%)	1 (5%)	14 (6%)	8 (5%)	7 (7%)	13 (6%)	2 (4%)
Lower leg	3 (1%)	2 (9%)	1 (0%)	3 (2%)	0	3 (1%)	0
Ankle	7 (3%)	0	7 (3%)	6 (4%)	1 (1%)	5 (2%)	2 (4%)
Foot	6 (2%)	0	6 (2%)	4 (2%)	2 (2%)	5 (2%)	1 (2%)
Muscle/tendon	144 (52%)	12 (55%)	132 (52%)	95 (56%)	49 (46%)	114 (51%)	30 (56%)
Cartilage/synovium/bursa	57 (21%)	6 (27%)	51 (20%)	35 (21%)	22 (21%)	49 (22%)	8 (15%)
Ligament/joint capsule	32 (12%)	1 (5%)	31 (12%)	15 (9%)	17 (16%)	26 (12%)	6 (11%)
Bone	2 (1%)	1 (5%)	1 (0%)	1 (1%)	1 (1%)	2 (1%)	0
Nervous	16 (6%)	1 (5%)	15 (6%)	7 (4%)	9 (8%)	11 (5%)	5 (9%)
Superficial tissues/skin	6 (2%)	0	6 (2%)	5 (3%)	1 (1%)	5 (2%)	1 (2%)
Non-specific	19 (7%)	1 (5%)	18 (7%)	11 (7%)	8 (7%)	15 (7%)	4 (7%)

compared with circus students,²⁹ reinforcing the need for a more holistic consideration of factors related to an injury.

Sex

Sex differences included a higher injury rate (5.69 vs 3.5/1000 exposures) and a higher proportion of acute than repetitive injuries for AMAB than AFAB. A study in collegiate athletes also found AMAB had more acute injuries.³⁰ However, differences were no longer apparent when analyses were limited to 10 sports with similar movement skills for both sexes.³⁰ In gymnastics, AFAB had more overuse injuries than AMAB,³⁰ possibly due to

apparatus differences. These studies suggest that gender biases in training are more likely responsible for this difference than anatomical or physiological sex differences.

In circus arts, sex differences in overall injury rates are uncommon,^{2 4 6 7} but become apparent for injury location. In our study, lumbosacral (14%; 9%) and hip/groin (9%; 2%) injuries were higher in AFAB than in AMAB. In other studies, AFAB had more hip/groin injuries^{2 4} and AMAB more wrist, forearm and ankle injuries.⁴ These differences may arise from gender biases in circus education, where women are often encouraged to train

Table 6 Injury frequency in relationship to activity (entire cohort)

	All	Adolescent	Adult	Preprofessional	Professional	Female	Male
Warm-up	4 (1%)	0	4 (2%)	2 (1%)	2 (2%)	3 (1%)	1 (2%)
Training	187 (66%)	13 (57%)	174 (67%)	115 (66%)	72 (65%)	151 (66%)	36 (55%)
Rehearsal	16 (6%)	2 (9%)	14 (5%)	12 (7%)	4 (4%)	11 (5%)	5 (9%)
Performance	16 (6%)	0	16 (6%)	6 (3%)	10 (9%)	11 (5%)	5 (9%)
Conditioning flexibility	22 (8%)	2 (9%)	20 (8%)	15 (9%)	7 (6%)	19 (8%)	3 (5%)
Conditioning strength	13 (5%)	2 (9%)	11 (4%)	10 (6%)	3 (3%)	12 (5%)	1 (2%)
Indirectly circus related	11 (4%)	2 (9%)	9 (3%)	6 (3%)	5 (5%)	9 (4%)	2 (4%)
Not circus related	7 (2%)	1 (4%)	6 (2%)	5 (3%)	2 (2%)	6 (3%)	1 (2%)
Unknown	7 (2%)	1 (4%)	6 (2%)	2 (1%)	5 (5%)	6 (3%)	1 (2%)
Aerial	95 (34%)	2 (9%)	93 (36%)	50 (29%)	45 (41%)	76 (33%)	19 (35%)
Aerial with ground elements	27 (10%)	0	27 (10%)	22 (13%)	5 (5%)	27 (12%)	0
Ground (human propulsion)	20 (7%)	3 (13%)	17 (7%)	14 (8%)	6 (5%)	15 (7%)	5 (9%)
Ground (apparatus propulsion)	1 (0%)	1 (4%)	0	1 (1%)	0	1 (0%)	0
Ground (balance/control)	69 (24%)	9 (39%)	60 (23%)	42 (24%)	27 (25%)	48 (21%)	21 (38%)
Manipulation	0	0	0	0	0	0	0
Character	0	0	0	0	0	0	0
Multidisciplinary	7 (2%)	0	7 (3%)	4 (2%)	3 (3%)	6 (3%)	1 (2%)
Warm-up	4 (1%)	0	4 (2%)	2 (1%)	2 (2%)	3 (1%)	1 (2%)
Conditioning flexibility	22 (8%)	2 (9%)	20 (8%)	15 (9%)	7 (6%)	19 (8%)	3 (5%)
Conditioning strength	13 (5%)	2 (9%)	11 (4%)	10 (6%)	3 (3%)	12 (5%)	1 (2%)
Indirectly circus related	11 (4%)	2 (9%)	9 (3%)	6 (3%)	5 (5%)	9 (4%)	2 (4%)
Not circus related	7 (2%)	1 (4%)	6 (2%)	5 (3%)	2 (2%)	6 (3%)	1 (2%)
Unknown	7 (2%)	1 (4%)	6 (2%)	2 (1%)	5 (5%)	6 (3%)	1 (2%)

Indirectly circus related, if related to circus but did not occur from participation in training or performance (eg, walking across the gym, spotting); not circus related, known cause outside of circus context (eg, riding a bike); unknown, insidious onset or participant unable to identify a cause; multidisciplinary, multiple disciplines involved in more than one discipline subgroup.

in disciplines exhibiting grace and flexibility and men in more dynamic disciplines.³¹ Understanding the impact of gender identity is also important, including the implications for non-binary (3.9% of our study population) and transgender/gender diverse artists (4.6% in another study).³²

Medical history

Overall self-reported eating disorder prevalence (18%) was lower than in another study of circus artists (36%),³³ however, 35.71% of AFAB adolescents in our study had a history of amenorrhoea despite none reporting an eating disorder. Amenorrhoea could indicate the physiological effects of underfueling and an undetected eating disorder. Similarly, only 1.5% of participants reported having a connective tissue disorder despite 34.5% meeting the GJH criteria. Lack of healthcare access may have contributed to the underdiagnosis of eating and connective tissue disorders and impacted injury burden and should be further examined in future research. Similar to other studies,^{34 35} eating disorders were associated with higher injury risk,

but unlike other sports research,^{36 37} amenorrhoea and history of connective tissue disorders were not. Baseline physical characteristics such as strength, flexibility and balance did not influence injury incidence. However, these factors are often the focus of injury prevention, suggesting that prevention strategies must shift to address our emerging understanding of risk factors.

Strengths

Despite the required weekly training log, the study had a high completion rate (77.11%). Including NTL injuries versus only TL or medical attention injuries better represents the breadth of circus injuries. Injury assessment by a PT enhances diagnostic accuracy for the injury cause and type compared with self-report.¹⁴ Using the circus-specific extension framework, including the circus discipline subgroups to report our data, will allow for valid comparisons to future research.

Limitations

Based on challenges during the pilot study,¹² exposure by hours was not tracked, limiting the comparison to some

research. Future studies should include total circus exposure in hours.⁹ Recruitment challenges limited adolescent participation (n=16), which could be increased with greater parent and coach engagement. There were also limited AMAB participants (n=29/201), which seems to reflect the circus demographics in the USA.³⁸

CONCLUSION

This study characterised injuries in preprofessional and professional circus artists in the USA. It showed that intrinsic (age, SAB and eating disorder history) and extrinsic factors (specific circus discipline exposure) affected injury risk in circus artists. When considering injury risk management, these factors must be applied holistically and contextually to the artist and their environment.

Correction notice The article has been corrected since it was published online. The acknowledgement section was missing which has been reinstated.

Acknowledgements We would like to thank the 201 circus artists who gave time to this study, Sarah Tiffin for her role as study coordinator, Mary McCall and Robin Greenspan for assistance with data analysis. We are grateful to our host facilities AcroSports, Circadium, The Circus Project, Esh Circus Arts, Kinetic Arts Center, Kinetic Theory, MOTH, New England Center for Circus Arts, Sky Candy, and Versatile Arts. And finally, we would like to acknowledge the study physical therapists Carol Bailey, Melissa Buffer-Trenouth, Michele Frances Benedisuk, Marisa Hentis, Traci Hurley, Greg Kowal, Emily Scherb, Tracey Wagner, and Pan Zhan.

Contributors SG conceived, designed, conducted and directed the analysis for this study, and is the guarantor. SG and MIS both significantly contributed to the data interpretation, writing and review of this manuscript.

Funding This work was supported by the Samuel Merritt University FSGP grant (grant number 215111820).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and the Samuel Merritt University Internal Review Board approved this study (SMUIRB#18-021). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Stephanie Greenspan <http://orcid.org/0000-0002-6570-5680>
Melanie I Stuckey <http://orcid.org/0000-0003-3478-1469>

REFERENCES

- Finch C. A new framework for research leading to sports injury prevention. *J Sci Med Sport* 2006;9:3–9.
- Shrier I, Meeuwisse WH, Matheson GO, et al. Injury patterns and injury rates in the Circus arts: an analysis of 5 years of data from cirque Du SOLEIL. *Am J Sports Med* 2009;37:1143–9.
- Wanke EM, McCormack M, Koch F, et al. Acute injuries in student Circus artists with regard to gender specific differences. *Asian J Sports Med* 2012;3:153–60.
- Munro D. Injury patterns and rates amongst students at the National Institute of Circus arts: an observational study. *Med Probl Perform Art* 2014;29:235–40.
- Stubbe JH, Richardson A, van Rijn RM. Prospective cohort study on injuries and health problems among Circus arts students. *BMJ Open Sport Exerc Med* 2018;4:e000327.
- Hakim H, Puel F, Bertucci W. Injury assessment in Circus student-artists population; preliminary study. *Sci Sports* 2020;35:154–60.
- Stuckey MI, Bruinooge B, Aubertin P, et al. Clinical burden of injuries in students at a professional Circus College: a 7.5-year longitudinal study. *Med Probl Perform Art* 2022;37:98–105.
- American Circus Educators, American Youth Circus Organization. Circus census 2022 final report. Available: <https://www.americancircuseducators.org/wp-content/uploads/2022/10/ACE-AYCO-Survey-Final-Report.pdf> [Accessed 05 Jan 2023].
- Greenspan S, Munro D, Nicholas J, et al. Circus-specific extension of the International Olympic Committee 2020 consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sport. *BMJ Open Sport Exerc Med* 2022;8:e001394.
- Faltus J, Richard V. Considerations for the medical management of the Circus performance artist and acrobat. *Int J Sports Phys Ther* 2022;17:307–16.
- Greenspan S. Injury patterns in subgroups of Circus artists by Circus discipline: a pilot study. *Orthop Phys Ther Pract* 2022;34:162–70.
- Greenspan S. Injury frequency and characteristics in adolescent and adult Circus artists: a pilot prospective cohort study. *Med Probl Perform Art* 2021;36:103–7.
- Bahr R, Clarsen B, Derman W, et al. International Olympic Committee consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sport 2020 (including STROBE extension for sport injury and illness surveillance (STROBE-SIIS)). *Br J Sports Med* 2020;54:372–89.
- Liederbach M, Hagins M, Gamboa JM, et al. Assessing and reporting dancer capacities, risk factors, and injuries: recommendations from the IADMS standard measures consensus initiative. *J Dance Med Sci* 2012;16:139–53.
- Fuller CW, Bahr R, Dick RW, et al. A framework for recording recurrences, reinjuries, and exacerbations in injury surveillance. *Clin J Sport Med* 2007;17:197–200.
- Lee L, Reid D, Cadwell J, et al. Injury incidence, dance exposure and the use of the movement competency screen (MCS) to identify variables associated with injury in full-time pre-professional dancers. *Int J Sports Phys Ther* 2017;12:352–70.
- Bronner S, McBride C, Gill A. Musculoskeletal injuries in professional modern dancers: a prospective cohort study of 15 years. *J Sports Sci* 2018;36:1880–8.
- Malfait F, Francomano C, Byers P, et al. The 2017 International classification of the Ehlers-Danlos syndromes. *Am J Med Genet* 2017;175:8–26.
- Bolling C, Mellette J, Pasman HR, et al. From the safety net to the injury prevention web: applying systems thinking to unravel injury prevention challenges and opportunities in cirque Du SOLEIL. *BMJ Open Sport Exerc Med* 2019;5:e000492.
- Nicholas J, Weir G, Alderson JA, et al. Incidence, mechanisms, and characteristics of injuries in pole dancers: a prospective cohort study. *Med Probl Perform Art* 2022;37:151–64.
- Campbell RA, Bradshaw EJ, Ball NB, et al. Injury epidemiology and risk factors in competitive artistic gymnasts: a systematic review. *Br J Sports Med* 2019;53:1056–69.
- Kerr ZY, Kroshus E, Grant J, et al. Epidemiology of national collegiate athletic association men's and women's cross-country injuries, 2009–2010 through 2013–2014. *J Athl Train* 2016;51:57–64.
- Long AS, Ambegaonkar JP, Fahringer PM. Injury reporting rates and injury concealment patterns differ between high-school cirque performers and basketball players. *Med Probl Perform Art* 2011;26:200–5.
- Soligard T, Schwellnus M, Alonso J-M, et al. How much is too much? (Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *Br J Sports Med* 2016;50:1030–41.
- Schwellnus M, Soligard T, Alonso J-M, et al. How much is too much? (Part 2) International Olympic Committee consensus statement on load in sport and risk of illness. *Br J Sports Med* 2016;50:1043–52.
- Liederbach M, Schanfein L, Kremenik IJ. What is known about the effect of fatigue on injury occurrence among dancers? *J Dance Med Sci* 2013;17:101–8.
- Campbell RS, Lehr ME, Livingston A, et al. Intrinsic modifiable risk factors in ballet dancers: applying evidence based practice principles to enhance clinical applications. *Phys Ther Sport* 2019;38:106–14.



- 28 Decker A, Aubertin P, Kriellaars D. Sleep and fatigue of elite Circus student-artists during one year of training. *Med Probl Perform Art* 2019;34:125–31.
- 29 Donohue B, Gavrilova Y, Galante M. Empirical development of a screening method for mental, social, and physical wellness in amateur and professional Circus artists. *Psychol Aesthet Creat Arts* 2020;14:313–24.
- 30 Yang J, Tibbetts AS, Covassin T, *et al.* Epidemiology of overuse and acute injuries among competitive collegiate athletes. *J Athl Train* 2012;47:198–204.
- 31 Funk A. Gender asymmetry and Circus education. *Performance Matters* 2018;4:19–35.
- 32 van Rens F, Heritage B. Mental health of Circus artists: psychological resilience, Circus factors, and demographics predict depression, anxiety, stress, and flourishing. *Psychol Sport Exerc* 2021;53:101850.
- 33 van Rens F, Metse AP, Heritage B. Exploring the mental health of Circus artists: Circus factors, psychological resilience, and demographics predict disordered eating and exercise addictions. *Psychol Sport Exerc* 2022;59:102107.
- 34 Meng K, Qiu J, Benardot D, *et al.* The risk of low energy availability in Chinese elite and recreational female aesthetic sports athletes. *J Int Soc Sports Nutr* 2020;17:13.
- 35 Rauh MJ, Nichols JF, Barrack MT. Relationships among injury and disordered eating, menstrual dysfunction, and low bone mineral density in high school athletes: a prospective study. *J Athl Train* 2010;45:243–52.
- 36 Pacey V, Nicholson LL, Adams RD, *et al.* Generalized joint hypermobility and risk of lower limb joint injury during sport: a systematic review with meta-analysis. *Am J Sports Med* 2010;38:1487–97.
- 37 Liaghat B, Pedersen JR, Young JJ, *et al.* Joint hypermobility in athletes is associated with shoulder injuries: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2021;22.
- 38 Huberman C, Scales M, Vallabhajosula S. Shoulder range of motion and strength characteristics in Circus acrobats. *Med Probl Perform Art* 2020;35:145–52.