


Perspectives of professional experts in relation to the development of community-based exercise for young adults with schizophrenia: a qualitative study

Martin Færch Andersen ^{1,2} Kickan Roed,¹ Allan Riis,^{2,3} Bolette Skjødt Rafn,^{1,4} Bjørn Hylsebeck Ebdrup,^{5,6} Julie Midtgaard^{1,6}

To cite: Andersen MF, Roed K, Riis A, *et al.* Perspectives of professional experts in relation to the development of community-based exercise for young adults with schizophrenia: a qualitative study. *BMJ Open Sport & Exercise Medicine* 2023;**9**:e001658. doi:10.1136/bmjsem-2023-001658

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjsem-2023-001658>).

Accepted 8 September 2023

ABSTRACT

Physical activity is a key modifiable factor in protecting physical and mental health in people with severe mental illness including schizophrenia. Therefore, early promotion of physical activity is recommended and programmes supporting long-term maintenance of physically active behaviour are warranted. This study aimed to explore the perspectives of professional experts in relation to the development of a sustainable community-based exercise programme tailored to young adults with schizophrenia and intended to promote change and adoption of physical activity. We conducted 9 semistructured interviews with 11 clinical and professional experts. Qualitative content analysis, as described by Graneheim and Lundman, was applied to analyse data. We identified four categories: (1) living a physical active life with schizophrenia, (2) exercise as promotor of personal recovery, (3) prescribing safe and relevant exercise and (4) instructors' qualifications and formation. When developing sustainable community-based exercise programmes tailored to young adults with schizophrenia, developers should ensure instructors' qualifications and provide an exercise protocol. In addition, developers should consider providing an inclusive and recovery-oriented exercise environment.

BACKGROUND

Severe mental illness, including schizophrenia, is among the most burdensome and costly illnesses worldwide.¹ The clinical symptoms usually manifest in early adult life,² and a substantial proportion of patients with schizophrenia experience persistent physical, functional and cognitive impairments.³ Psychotic symptoms are often effectively treated with antipsychotic medication, however, treatment is commonly accompanied by metabolic side effects. Patients treated with antipsychotic medication have a 4-fold higher prevalence of metabolic syndrome,⁴ and a 2–3 fold higher risk of cardiovascular disease compared with

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Physical activity is a key modifiable factor in protecting physical and mental health in people with severe mental illness, including schizophrenia.
- ⇒ Sustainable community-based programmes to support adoption of physical activity for young adults with schizophrenia are warranted.

WHAT THIS STUDY ADDS

- ⇒ Developers of community-based exercise programmes for young adults with schizophrenia must ensure a strategy for identifying and training exercise instructors as well as providing a protocol for delivering safe and clinically relevant exercise.
- ⇒ Community-based exercise can promote personal recovery and considerations of how to balance overcoming potential barriers to participation with the goal of creating a non-clinical exercise environment are important.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ These findings could support and inform the development of community-based programmes promoting physical activity for people with schizophrenia and may be transferable or adaptable to other psychiatric populations in other countries.

the general population,⁴ which contributes to a premature mortality rate of 15–20 years.⁵ Lifestyle factors, such as poor diet, smoking⁶ and physical inactivity,⁷ may contribute to the increased morbidity and mortality.

The Lancet Psychiatry Commission has pointed to physical activity as a key modifiable factor in protecting physical health in people with mental illness.⁸ Specifically, introducing exercise in the early stages of schizophrenia may be a sustainable solution in preventing or attenuating metabolic dysfunction associated with antipsychotic medication.^{8 9} Furthermore, exercise has been found efficacious in



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Martin Færch Andersen;
mfan@ucn.dk

improving clinical symptoms, quality of life, global functioning and cognitive deficits,^{10–12} and early improvements in these outcomes may reduce the likelihood of enduring symptoms and functional disability.¹³ Moreover, patients in the early stages of schizophrenia are more physically active than patients with long-term schizophrenia,¹⁴ and thus potentially easier to engage in exercise. However, long-term maintenance of physical active behaviour is challenging.¹⁵ Replicable and scalable methods of delivering physical activity to patients with schizophrenia in the early stages of their disease in a format that is accessible, engaging and effective for a large number are warranted.¹⁶ Indeed, the WHO's Mental Health Action Plan 2013–2030 calls for the provision of mental health services integrated in communities for service users and families.¹⁷ Accordingly, community-based group exercise may hold promise to support sustainable physical activity.¹⁸ Still, the initiation of community-based exercise in people with serious mental illness is not a simple step from intention to participation, but a non-linear slow phased transition with various challenges and setbacks at every phase.¹⁹

This study was conducted as a part of a project aiming to develop, evaluate and implement community-based exercise for young adults in antipsychotic treatment (ClinicalTrials.gov identifier: NCT05461885). Previously, we evaluated the impact and feasibility of a gym-based group exercise programme supervised by non-health professional exercise instructors for young adults in antipsychotic treatment (the COPUS trial).^{20–21} The results indicated that gym-based group exercise has the potential to support and promote personal recovery, which can be defined as 'changing values, feelings, goals, abilities and roles in order to achieve a satisfactory, hopeful and productive way of life, with the possible limitation of the illness'.²² However, feasibility was challenged due to a lack of incentives and infrastructure to ensure consecutive recruitment from outpatient clinics to gym-based exercise in communities. Consequently, it was concluded that refinements in both the intervention itself and in its the delivery were warranted before initiating a large-scale, definitive trial.²⁰ Also, community-based exercise programmes may be considered a complex intervention due to the properties of the intervention itself, that is, how the intervention produces change, and the interaction between the intervention and its context, that is, how the context affects implementation and outcomes. Hence, early considerations and continually revisiting core elements, that is, considering context, identifying key uncertainties, developing a programme theory and refining the intervention, are recommended throughout the research process when evaluating complex interventions.^{23–24} The current study aimed to explore the perspectives of various professional experts in relation to the development of community-based exercise for young adults with schizophrenia intended to promote long-term maintenance of physical activity.

METHODS

Design

We applied a descriptive qualitative design²⁵ using researcher triangulation and semi-structured interviews. This study follows the Consolidated Criteria for Reporting Qualitative Research checklist²⁶ (online supplemental material 1).

Sampling

We used a purposeful sampling strategy to support information richness.²⁷ Specifically, we used snowball sampling to reach unique key informants with expert knowledge, experience or interest concerning the study's aims.²⁸ Key informants were thus recruited among professional experts within exercise as health promotion and/or treatment of psychosis, for example, psychiatry, physiology, exercise psychology. All key informants who were contacted agreed to participate. The final sample was defined by theoretical saturation, that is, different expertise and perspectives represented, and by inductive thematic saturation during the analysis focusing on emerging new perspectives.²⁹

Data collection

KR (female investigator, registered nurse, full-time PhD student) or MFA (male investigator, certified physiotherapist, full-time PhD student) carried out the semistructured individual interviews, which lasted 45–60 min, using video calls on Microsoft Teams or physical face-to-face meetings. Only the interviewer and the key informants were present during interviews. Both KR and MFA have previous experience in conducting qualitative interviews and had no personal knowledge of the key informants beforehand. All key informants were provided with written or oral information regarding the study aim prior to the interviews. Hence, it was explained that community-based exercise was defined as group exercise delivered by non-health professional exercise instructors outside a hospital setting (eg, commercial gyms or sporting clubs). The interview guide comprised a standard open-ended question.

According to you, as an expert, what should be considered when developing a community-based exercise programme for young adults with schizophrenia?

In addition, each key informant was then asked specified questions thematically related to their area of expertise (online supplemental material 2). The face-to-face interviews were audiorecorded, and the online video calls were video recorded. The interviewer (KR or MFA) who had conducted a specific interview wrote selective transcriptions, that is, focusing exclusively on content relevant to the research question,³⁰ which was then validated by the other researcher (KR or MFA) while focusing on and discussing the emergence of new perspectives to determine data saturation.²⁹ To enhance trustworthiness, the key informants received the transcript within 1 week after the interview for member checking.³¹ Selective




Figure 1 Example of the analytical process from meaning unit to category: living a physical active life with schizophrenia.

transcripts were adjusted according to comments from member checking when needed. Additional methodological considerations are available in online supplemental material 2.

Data analysis

The interviews were analysed using inductive qualitative content analysis, as described by Graneheim and Lundman.³² NVivo V.12 (QRS International, Melbourne, Australia) and Microsoft Excel were used to assist in data management.

MFA and KR initially read and reread the transcripts separately before jointly discussing them to obtain a sense of the complete data material. Subsequently, MFA identified and extracted meaning units and labelled them with descriptive codes. Next, MFA compared, abstracted and sorted codes into subcategories and categories, which were discussed with KR and JM (female psychologist, full-time researcher and principal investigator). Codes, subcategories and categories were continuously compared with original condensed summaries in an iterative process. Figure 1 provides an example of the condensation-abstraction process.

RESULTS

Characteristics of informants

Nine semistructured interviews were conducted with 11 key informants (3 key informants were interviewed simultaneously) representing different professional backgrounds, that is, clinical psychiatry (n=1), psychology (n=4), human physiology (n=1), consultants with expertise in exercise for people with mental illness (n=2) and physiotherapy (n=2) and occupational therapy (n=1) within mental healthcare. All key informants had more than 10 years of experience working with people diagnosed with schizophrenia, exercise physiology and/or exercise psychology.

Analysis of findings

Four main categories, 11 subcategories and 34 codes were identified (table 1). In the following presentation of the results young adults with schizophrenia will be addressed only as participants and non-health professional exercise instructors only as instructors as these were the terms used most frequently by the key informants.

Category 1: living a physical active life with schizophrenia

Schizophrenia is often associated with debilitating symptoms which have a major impact on daily life and therefore also on the ability to engage in physical activities. The key informants emphasise that taking these potential barriers into consideration when developing a community-based exercise programme is vital along with

Table 1 Results of the analysis

Categories	Subcategories
Living a physical active life with schizophrenia	Schizophrenia and living life
	How schizophrenia affects physical activity
	Exercise as a promotor of personal recovery
Exercise as a promotor of personal recovery	Being part of an exercise culture
	Meeting at eye level
	Striving for commonness
	Feeling empowered
Prescribing safe and relevant exercise	Flexible exercise protocol
	Contact to trusted others
	Do no harm
Instructors' qualifications and formation	Experienced and passionate
	Learning through acting and reflection

ensuring that the exercise is supervised by instructors, who recognise these barriers.

Schizophrenia and living life

Suffering from schizophrenia may be perceived as a life crisis that may dampen hopes and dreams of living a 'normal' life for the affected person. Community-based exercise may serve as a break from the disorder but may also constitute with yet another stressor:

Being struck by psychosis for a period in life can be exceedingly difficult... For some, physical exercise may be a ray of light that distracts them from the disease... But for others it may add to the list of things that have become unmanageable. (Expert in clinical psychiatry)

According to key informants with knowledge about mental illness, symptoms of schizophrenia may include a psychophysical split, implying that participants experience detachment from their own body. This may challenge participation in physical activity; however, exercise also has the potential to improve this phenomenon:

You can feel bodily disorganised and do not experience the body as one entity. There may also be the sensation that the body is levitating, and here physical exercising may be positive. (Expert in physiotherapy within mental healthcare)

How schizophrenia affects physical activity

The key informants acknowledge that community-based exercise should be delivered by instructors, who have received basic education about schizophrenia, so they understand the participants' potential reactions. Nonetheless, the education should not encourage an excessive focus on the disease:

Must NOT dominate! Knowledge about this [psychotic disorders] should enable instructors to have realistic and informed expectations... But they should also know that people with mental illness are not significantly different from other people. (Expert in psychology and recovery)

Especially, knowledge about psychotic and negative symptoms and how they may affect the participants' ability to engage in physical activity with others is important, as the general population often misinterprets these symptoms.

Negative symptoms play an important role in the disease and are experienced as the most debilitating aspect associated with a schizophrenia diagnosis, in contrast to the general perception that it must be the psychotic symptoms... The intentions are present, but the initiative is absent. (Expert in clinical psychiatry)

Furthermore, severe weight gain is a typical side effect of antipsychotic medication which is often accompanied by considerable psychosocial distress. Several key informants caution, however, against weight loss becoming

the most important goal as losing weight is difficult to achieve with physical activity alone. Instead, the experience of well-being and having a strong body should be promoted.

Category 2: exercise as a promotor of personal recovery

The key informants promoted the view that a community-based group exercise programme may have potential to facilitate participants' personal recovery process. However, to achieve this potential, the programme should create an inclusive exercise environment.

Being part of an exercise culture

Of importance, the key informants stated that the participants should ideally achieve a feeling of being a part of something bigger than just an exercise programme; thus, participants should enter an exercise culture where being physically active is part of a regular everyday routine. To succeed, the setting must feel like a safe space where participants know the instructors and vice versa and atmosphere should reflect that all levels of participation are acceptable.

It must be a cool experience. If that succeeds... then the potential for physical activity has a chance to unfold. (Expert in psychology and physical activity for recovery)

A safe atmosphere is created not only by the exercise instructor's facilitation but more importantly within the group of participants. As a result, group interaction should be equally, if not more highly, prioritised compared with the content of the exercise. Nurturing social relationships between the participants may be challenged because of, for example, negative symptoms or anxiety; hence the instructor should take charge of facilitating a feeling of being a part of a group before, during and after the exercise session. This may include having team exercise activities or encourage socialising after exercise sessions.

Meeting at eye level

The key informants expressed that it is important to have a caring communication without talking down to or being overprotective towards the participants. Participants may have experienced various disappointments and defeats regarding physical activity, potentially resulting in negative expectations and low self-confidence. Thus, self-stigma is considerable, and it must be acknowledged that even showing up may be a large accomplishment:

They [participants] will be tough enough on themselves if they only show up two out of three times. It's important to say, "So nice to see you" rather than "Where were you last time?" Small things make the difference. (Expert in psychology and schizophrenia)

Striving for commonness

One key informant underpins that the community-based exercise must not be a sanctuary for people with mental disorders disconnected from the rest of the world. Others state that being physically active where other young people are present and physically active might potentially be of great importance to the participants' self-worth and in the (re)building of an exercise identity:

For a young patient to tell other young people that: "in my spare time I go to the gym three times a week" can be of great importance. (Expert in physiotherapy within mental healthcare)

The participants should be met with the same demands and commitments as everyone else without violating their autonomy; it should be okay to withdraw for a moment if needed. There may be a general conception that people with schizophrenia are fragile and vulnerable, which may constitute a pitfall, and the exercise instructor may accidentally take on a therapeutic role:

Instructors should listen but abstain from being therapeutic and giving advice [on mental health] and be aware of their task—which is exercise. (Expert in physiotherapy within mental healthcare)

Feeling empowered

According to the key informants, a community-based exercise programme should seek to facilitate empowerment and draw attention to a positive sensation of being in control, being strong, having more energy and sleeping better; in other words, of connecting exercise with a feeling of well-being. One key informant, however, emphasised the importance of recognising that exercise alone may not function as the main driver for all participants, but that they may find the context surrounding the exercise, such as the people and the place, more important to their well-being.

It may not be exercise as such making the difference, but the meaningfulness placed within the exercise. (Expert in exercise for people with mental illness)

Category 3: prescribing safe and relevant exercise

Key informants agree that when developing a community-based exercise programme for young adults with schizophrenia, it is pivotal to ensure safety procedures in the case of physical and mental adverse events. Furthermore, the exercise protocol should have the flexibility to allow individual adjustments.

Flexible exercise protocol

Several key informants acknowledge that the exercise protocol should have a clear and recognisable structure. When the structure is known, varying the content and complexity of the exercise is easier for the instructors and more acceptable to the participants. It is important that

the exercise content can be adjusted to fit the individual's physical, mental and social capacity.

It's damn difficult because it's not only about making the exercise easier or harder, but also about what the patient is like in the room and how they interact with the others. Instructors must acknowledge the complexity of an exercise situation. (Expert in physiotherapy within mental healthcare)

The protocol must be flexible enough to accommodate individual goals of participants. However, even though individual goals may be mental or social, it is important that the instructors pursue clinically relevant exercise intensities given the considerably increased risk of metabolic diseases.

It's important to pay attention to every single patient's level of activity throughout the sessions, so each patient reaches a high level of activity over a longer period of time. (Expert in human physiology)

Contact to trusted others

The key informants highlight that the support of both relatives and healthcare professionals is essential since they serve as external motivators and improve the ability of participants to attend the exercise sessions. Furthermore, inviting a friend or next of kin may ease the initial process of joining the group exercise. Because participants may express psychotic, suicidal or aggressive behaviour during an exercise session, it is important that exercise instructors are provided safety procedures regarding what to do and who to contact. However, the key informants emphasise that such situations are very rare.

Do no harm

Some key informants express that elements of or being physically touched by others may be distressing for some participants. Furthermore, bodily discomfort such as delayed onset muscle soreness or an elevated heartbeat may be misinterpreted as dangerous and the normality of this needs to be addressed before, during and after exercise sessions.

What can be expected after exercise, such as muscle soreness, and what can be expected during exercise, such as sweating, elevated heartbeat, and shortness of breath. Some of it may be confused with symptoms of anxiety. (Expert in physiotherapy in mental healthcare)

Category 4: instructors' qualification and formation

When developing a community-based exercise programme for young adults with schizophrenia, identifying the right exercise instructors is crucial. The key informants highlight that they should be fully engaged in their role as exercise instructors and receive formal



training containing theory and the exchange of practical experience.

Experienced and passionate

According to the key informants, exercise instructors should be unpretentious and reliable to best support participants, who might be entering an unknown activity and culture. To do so, experience as an exercise instructor along with being passionate about this role is needed to focus on resonating with the participants. Furthermore, instructors should have relationship-building skills and be able to act as external motivation.

A motivational instructor is crucial for success.
Without that, a good programme is worth nothing.
(Expert in exercise psychology)

Learning through acting and reflection

The key informants recommend that exercise instructors must receive formal training combining theoretical sessions as well as reflection on their own experiences to facilitate exercise for people with mental illness. Since instructors may have sparse experience and knowledge about mental illness, facilitating an ongoing exchange of experience is essential.

They [instructors] should receive more than just one day of training, actually a whole course, with the possibility to try something out, exchange experiences, and receive qualified feedback. (Expert in physiotherapy within mental healthcare)

DISCUSSION

This qualitative study of the perspectives of clinical and professional experts offers a unique insight into core considerations when developing community-based exercise tailored to young adults with schizophrenia. It was widely acknowledged that mental health and social care initiatives in community-based settings are needed for people with severe mental illness to provide an opportunity to attain the highest level of health and participate in society free of stigmatisation and discrimination.³³ Thus, key informants acknowledge that community-based exercise programmes have the potential to promote long-term maintenance of physical and mental health for young adults with schizophrenia, and thereby achieve the proposed potential of exercise.^{8 10–12}

Notably, our findings show a belief among experts in the potential of community-based exercise as an important contributor to personal recovery. They highlight that exercise could facilitate the experience of empowerment, that is, of being strong and in control, which is an important factor in the recovery process according to one of the most accepted theoretical frameworks present to understand personal recovery, CHIME (Connectedness, Hope and optimism about the future, Identity, Meaning of life and Empowerment).³⁴ Furthermore, qualitative findings indicate that participating in sports contributes

to experiencing a sense of achievement³⁵ and may represent an untapped resource in personal recovery as it can serve as an arena that promotes physically active behaviour while providing an opportunity to build life skills and social connectivity.³⁶ Indeed, the key informants assert that group interaction should be prioritised at least as high as the content of the exercise. In addition, the community and the location should be a place that supports the participants in transitioning beyond their role as patients to help them (re)build an exercise identity. Qualitative findings from our recent COPUS trial found that group exercise delivered in a conventional fitness centre by non-health professional exercise instructors supported a feeling of being a normal young adult.²¹

The COPUS trial found that community-based exercise in a local gym was meaningful and safe for young adults with schizophrenia; however, it also reported that its feasibility was challenged due to limited recruitment and retention rates as the study setup did not provide the mental health staff with sufficient incitements and infrastructure to secure consecutive screening and promotion of the intervention.²⁰ The key informants in this study highlight that support from trusted individuals, such as the primary healthcare provider, has an immense influence on the ability of people with schizophrenia to participate. This aligns with a qualitative review suggesting that emotional and practical support from either health professionals and/or relatives plays an important role.¹⁹ Moreover, while the exercise protocol should allow all levels of participation, clinically relevant exercise doses and intensities should be striven for to improve metabolic conditions. Current evidence suggests that higher doses of aerobic exercise are not only recommended to improve cardiovascular health but also clinical and functional outcomes.³⁷

Another important finding is the need for carefully recruiting dedicated exercise instructors as their personal engagement and ability to resonate with participants is crucial for success. This is in line with other developmental research on community-based physical activity programmes for people with mental illness and other clinical populations. Here, relationship-building skills, including empathy and the ability to generate team spirit are described as important instructor qualifications.^{19 38 39} The key informants in our study stressed the importance of instructors having formal education and knowledge on how symptomatology and the potential side effects of medical treatment may affect participation to allow them to have realistic and informed expectations as also supported by an international consensus statement.⁴⁰ Especially negative symptoms, such as lack of energy, low self-esteem, depressive symptoms and apprehensive attitude to socialising, are the most frequently reported barriers towards exercise participation among people with schizophrenia.^{19 41} Furthermore, cognitive symptoms related to thinking about, planning and getting to an activity is also reported as a significant barrier.¹⁹ Knowledge about these symptoms may help instructors

understand the importance of creating a welcoming and safe environment.

Methodological considerations

This study has some strengths and limitations which should be taken into consideration. Despite our confidence in using purposeful sampling and snowball sampling aiming for variation in relation to the key informants' areas of expertise, we cannot rule out an element of recruitment bias implying that key informants may have been positive about incorporating physical activity in community mental healthcare. Due to the geographical distance between key informants and the interviewers, most interviews were conducted using video calls. We found that this format was flexible, served the intended objective and provided sufficient information power. For pragmatic reasons three informants who were colleagues were interviewed simultaneously. To avoid potential conformity and authority bias the interviewer continuously asked the respective expert to justify their answers and posed critical follow-up questions, for example, related to disadvantages or issues associated with a specific viewpoint. Furthermore, each expert received the transcript for additional comments which were not shared with the other informants. Selective transcripts were used as empirical data, a process in which some relevant content may have been lost. Thus, to enhance trustworthiness, the transcripts were validated by a second researcher and sent to the experts for member checking.

Clinical implications

Based on the perspectives of clinical and professional experts in the field, we propose the following recommendations encompassing structural and cultural factors in relation to promotion of long-term maintenance of physical activity in young adults with schizophrenia (figure 2).

- Consider how schizophrenia may produce various barriers toward exercise participation and how

paying special attention to them can be balanced with promoting a normal exercise environment.

- Consider how exercise may promote the personal and social recovery process, for example, the CHIME framework, in the development of gym-based exercise programmes to support the sensation of empowerment, social connectivity and exercise identity in participants.
- Ensure that safe and relevant exercise is prescribed, specifically by formulating a flexible exercise protocol and including other trusted individuals, for example, relatives, friends or mental health staff, to support participation and to agree on safety procedures.
- Ensure instructors' qualifications and training, specifically by drawing up a strategy to identify and recruit experienced and passionate exercise instructors and by establishing an educational course and professional network that allows instructors to gain insight into schizophrenia and share reflection on practical experiences.

The current study was carried out in Denmark where the mental healthcare system and the community sector are closely interconnected to provide comprehensive care to people with mental illness. Despite this context, we believe that the recommendations delivered from this study may be applied or adjusted for the development of programmes promoting physical activity among diverse psychiatric populations and in other countries.

CONCLUSION

This study provides key considerations when developing sustainable community-based exercise programmes tailored to young adults with schizophrenia intended to promote long-term maintenance of physical activity. Developers should focus on structural factors by ensuring instructors' qualifications and providing a flexible exercise protocol, and cultural factors such as facilitating an inclusive and recovery-oriented exercise environment. These findings may be transferable to the development of programmes promoting physical activity in other psychiatric populations.

Author affiliations

¹Centre for Applied Research in Mental Health Care, Psychiatric Center Glostrup, Glostrup, Denmark

²Department of Physiotherapy, University College of Northern Denmark, Aalborg, Denmark

³Center for General Practice, Aalborg University, Aalborg, Denmark

⁴Department of Oncology, Rigshospitalet Department of Oncology, Copenhagen, Denmark

⁵Center for Neuropsychiatric Schizophrenia Research, Psychiatric Center Glostrup, Glostrup, Denmark

⁶Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Acknowledgements The authors are thankful to the key informants who participated in this study, for giving their time and providing detailed accounts of their expertise to make this work possible.

Contributors All authors participated in conceptualising the study. MFA, KR and JM planned the study design and methodology. MFA and KR were responsible for



Figure 2 Structural and cultural factors of importance when developing community-based exercise targeting young adults with schizophrenia.

the data collection and MFA was responsible for the initial data analysis. MFA, KR and JM contributed to the data analysis process. MFA generated the first draft, and all authors critically revisited the draft for important intellectual content. Lastly, the final version was sent to all authors for approval. All authors read and approved the final manuscript. MFA is the guarantor of the overall content of the study.

Funding The study was funded by TrygFonden (Grant No: 151603).

Competing interests BHE is part of the Advisory Board of Eli Lilly Denmark A/S, Janssen-Cilag, Lundbeck Pharma A/S and Takeda Pharmaceutical Company; and has received lecture fees from Bristol-Myers Squibb, Boehringer Ingelheim, Otsuka Pharma Scandinavia AB, Eli Lilly Company, and Lundbeck Pharma A/S.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval In addition to receiving written information about the nature of the study prior to participation, all key informants provided oral informed consent. All key informants were guaranteed anonymity and confidentiality. Thus, quotes presented in the results section are not linked with the informant ID. The study was conducted in accordance with the Declaration of Helsinki. The Regional Ethics Committee of Northern Denmark has confirmed that no formal ethical approval was required (2023000206) for the current study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Martin Færch Andersen <http://orcid.org/0000-0003-0469-7557>

REFERENCES

- Charlson FJ, Ferrari AJ, Santomauro DF, *et al*. Global epidemiology and burden of schizophrenia: findings from the global burden of disease study 2016. *Schizophr Bull* 2018;44:1195–203.
- Immonen J, Jääskeläinen E, Korpela H, *et al*. Age at onset and the outcomes of schizophrenia: a systematic review and meta-analysis. *Early Interv Psychiatry* 2017;11:453–60.
- Secher RG, Hjorthøj CR, Austin SF, *et al*. Ten-year follow-up of the OPUS specialized early intervention trial for patients with a first episode of psychosis. *Schizophr Bull* 2015;41:617–26.
- Vancampfort D, Stubbs B, Mitchell AJ, *et al*. Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. *World Psychiatry* 2015;14:339–47.
- Hjorthøj C, Stürup AE, McGrath JJ, *et al*. Years of potential life lost and life expectancy in schizophrenia: a systematic review and meta-analysis. *Lancet Psychiatry* 2017;4:295–301.
- Teasdale SB, Ward PB, Samaras K, *et al*. Dietary intake of people with severe mental illness: systematic review and meta-analysis. *Br J Psychiatry* 2019;214:251–9.
- Vancampfort D, Firth J, Schuch FB, *et al*. Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. *World Psychiatry* 2017;16:308–15.
- Firth J, Siddiqi N, Koyanagi A, *et al*. The lancet psychiatry commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry* 2019;6:675–712.
- Gates J, Killackey E, Phillips L, *et al*. Mental health starts with physical health: current status and future directions of non-pharmacological interventions to improve physical health in first-episode psychosis. *Lancet Psychiatry* 2015;2:726–42.
- Bredin SSD, Kaufman KL, Chow MI, *et al*. Effects of aerobic, resistance, and combined exercise training on psychiatric symptom severity and related health measures in adults living with schizophrenia: a systematic review and meta-analysis. *Front Cardiovasc Med* 2021;8:753117.
- Dauwan M, Begemann MJH, Heringa SM, *et al*. Exercise improves clinical symptoms, quality of life, global functioning, and depression in schizophrenia: a systematic review and meta-analysis. *Schizophr Bull* 2016;42:588–99.
- Firth J, Stubbs B, Rosenbaum S, *et al*. Aerobic exercise improves cognitive functioning in people with schizophrenia: a systematic review and meta-analysis. *Schizophr Bull* 2017;43:546–56.
- Álvarez-Jiménez M, Gleeson JF, Henry LP, *et al*. Road to full recovery: longitudinal relationship between symptomatic remission and psychosocial recovery in first-episode psychosis over 7.5 years. *Psychol Med* 2012;42:595–606.
- Walther S, Stegmayer K, Horn H, *et al*. Physical activity in schizophrenia is higher in the first episode than in subsequent ones. *Front Psychiatry* 2014;5:191.
- Firth J, Carney R, French P, *et al*. Long-term maintenance and effects of exercise in early psychosis. *Early Interv Psychiatry* 2018;12:578–85.
- Stubbs B, Vancampfort D, Hallgren M, *et al*. EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and position statement from the European Psychiatric Association (EPA), supported by the International organization of physical therapists in mental health (IOPTMH). *Eur Psychiatry* 2018;54:124–44.
- WHO. *Comprehensive mental health action plan 2013–2030*. Geneva: World Health Organization, 2021.
- Soundy A, Freeman P, Stubbs B, *et al*. The transcending benefits of physical activity for individuals with schizophrenia: a systematic review and meta-ethnography. *Psychiatry Res* 2014;220:11–9.
- Quirk H, Hock E, Harrop D, *et al*. Understanding the experience of initiating community-based group physical activity by people with serious mental illness: a systematic review using a meta-ethnographic approach. *Eur Psychiatry* 2020;63:e95.
- Midtgaard J, Schnor H, Bjerre ED, *et al*. Exercise training complementary to specialised early intervention in patients with first-episode psychosis: a feasibility randomised trial. *Pilot Feasibility Stud* 2021;7:162.
- Larsen LQ, Schnor H, Tersbøl BP, *et al*. The impact of exercise training complementary to early intervention in patients with first-episode psychosis: a qualitative sub-study from a randomized controlled feasibility trial. *BMC Psychiatry* 2019;19:192.
- Anthony WA. Recovery from mental illness: the guiding vision of the mental health service system in the 1990s. *Psychosocial Rehabilitation Journal* 1993;16:11–23.
- Skivington K, Matthews L, Simpson SA, *et al*. A new framework for developing and evaluating complex interventions: update of medical research Council guidance. *BMJ* 2021;374:n2061.
- Moore GF, Audrey S, Barker M, *et al*. Process evaluation of complex interventions: medical research council guidance. *BMJ* 2015;350:h1258.
- Sandelowski M. Whatever happened to qualitative description? *Res Nurs Health* 2000;23:334–40.
- Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- Patton MQ. Enhancing the quality and credibility of qualitative analysis. *Health Serv Res* 1999;34:1189–208.
- Penrod J, Preston DB, Cain RE, *et al*. A discussion of chain referral as a method of sampling hard-to-reach populations. *J Transcult Nurs* 2003;14:100–7.
- Saunders B, Sim J, Kingstone T, *et al*. Saturation in qualitative research: exploring its conceptualization and operationalization. *Qual Quant* 2018;52:1893–907.
- McMullin C. Transcription and qualitative methods: implications for third sector research. *Voluntas* 2023;34:140–53.
- Birt L, Scott S, Cavers D, *et al*. Member checking. *Qual Health Res* 2016;26:1802–11.
- Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse Educ Today* 2004;24:105–12.
- Killaspay H, Harvey C, Brasier C, *et al*. Community-based social interventions for people with severe mental illness: a systematic review and narrative synthesis of recent evidence. *World Psychiatry* 2022;21:96–123.
- Leamy M, Bird V, Le Boutillier C, *et al*. Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *Br J Psychiatry* 2011;199:445–52.

- 35 Carless D, Douglas K. Narrative, identity and mental health: how men with serious mental illness re-story their lives through sport and exercise. *Psychology of Sport and Exercise* 2008;9:576–94.
- 36 Brooke LE, Lin A, Ntoumanis N, *et al.* Is sport an untapped resource for recovery from first episode psychosis? A narrative review and call to action. *Early Interv Psychiatry* 2019;13:358–68.
- 37 Stubbs B, Vancampfort D, Hallgren M, *et al.* EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and position statement from the European Psychiatric Association (EPA) supported by the International organization of physical therapists in mental. *Eur Psychiatry* 2018;54:124–44.
- 38 Bjerre ED, Leth M, Hammer NM, *et al.* Development of an educational program for non-professional soccer coaches in charge of community-based soccer in men with prostate cancer: a qualitative study. *Sports Med Open* 2018;4:31.
- 39 Heston A-H, Schwartz AL, Justice-Gardiner H, *et al.* Addressing physical activity needs of survivors by developing a community-based exercise program: LIVESTRONG® at the YMCA. *Clin J Oncol Nurs* 2015;19:213–7.
- 40 Rosenbaum S, Davison K, Stanton R. The role of sport, exercise, and physical activity in closing the life expectancy gap for people with mental illness: an international consensus statement by exercise and sports SC... pragmatic evaluation course 2017 Vancouver network view project surf therapy view project. 2018. Available: <http://www.acsm-tj.org>
- 41 Firth J, Rosenbaum S, Stubbs B, *et al.* Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychol Med* 2016;46:2869–81.

Perspectives of professional experts in relation to the development of community-based exercise for young adults with schizophrenia – A qualitative study

MF Andersen^{1,2*}, K Roed¹, A Riis^{2,3} BS Rafn^{1,4}, BH Ebdrup^{5,6} J Midtgaard^{1,6}

*Correspondence: MF Andersen, mfan@ucn.dk

¹ Centre for Applied Research in Mental Health Care (CARMEN), Mental Health Centre Glostrup, University of Copenhagen, Copenhagen, Denmark

² Department of Physiotherapy, University College of Northern Denmark, Aalborg, Denmark

Supplemental Material 1 Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

Please note that the COREQ checklist originally operates with reporting items according to which page number they can be found on. The page numbers in this supplementary file refer to the original manuscript. An added row to the checklist provides information as to which section and heading of the article specific items are reported in.

Developed from:

Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Heal Care*. 2007;19(6):349-357. doi:10.1093/INTQHC/MZM042

No. Item	Guide questions/description	Reported on Page #	Section and heading
Domain 1: Research team and reflexivity			
<i>Personal Characteristics</i>			
1. Inter viewer/facilitator	Which author/s conducted the interview or focus group?	6	Methods under 'Data collection'
2. Credentials	What were the researcher's credentials? E.g. PhD, MD	6	Methods under 'Data collection'
3. Occupation	What was their occupation at the time of the study?	6	Methods under 'Data collection'
4. Gender	Was the researcher male or female?	6	Methods under 'Data collection'
5. Experience and training	What experience or training did the researcher have?	6	Methods under 'Data collection'
<i>Relationship with participants</i>			
6. Relationship established	Was a relationship established prior to study commencement?	6	Methods under 'Data collection'
7. Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	6	Methods under 'Data collection'
8. Interviewer characteristics	What characteristics were reported about the inter viewer/facilitator? e.g. Bias, assumptions, reasons and	16	Discussion under 'Methodological considerations'

	interests in the research topic		
Domain 2: study design			
<i>Theoretical framework</i>			
9. Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	7	Methods under 'Data analysis'
<i>Participant selection</i>			
10. Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	6	Methods under 'Sampling'
11. Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	6	Methods under 'Sampling'
12. Sample size	How many participants were in the study?	8	Results under 'Characteristics of informants'
13. Non-participation	How many people refused to participate or dropped out? Reasons?	6	Methods under 'sampling'
<i>Setting</i>			
14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	6	Methods under 'Data collection'
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	6	Methods under 'Data collection'
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	8, Table 1	Results under 'Characteristics of informants'
<i>Data collection</i>			
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	6 Appendix 2	Methods under 'Data collection'
18. Repeat interviews	Were repeat interviews carried out? If yes, how many?	7	Methods under 'Data collection'
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	6	Methods under 'Data collection'
20. Field notes	Were field notes made during and/or after the interview or focus group?	7	Methods under 'Data collection'
21. Duration	What was the duration of the interviews or focus group?	6	Methods under 'Data collection'
22. Data saturation	Was data saturation discussed?	6, 7	Methods under 'Sampling' and 'Data collection'
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	6, 7	Methods under 'Data collection'
Domain 3: analysis and findings			
<i>Data analysis</i>			
24. Number of data coders	How many data coders coded the data?	7	Methods under 'Data analysis'
25. Description of the coding tree	Did authors provide a description of the coding tree?	8 Table 2	Results under 'Analysis of findings'
26. Derivation of themes	Were themes identified in advance or derived from the data?	7	Methods under 'Data analysis'
27. Software	What software, if applicable, was used to manage the data?	7	Methods under 'Data analysis'

28. Participant checking	Did participants provide feedback on the findings?	7	Methods under 'data collection'
<i>Reporting</i>			
29. Quotations presented	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	8-14	Results under 'Analysis of findings'
30. Data and findings consistent	Was there consistency between the data presented and the findings?	8-14	Results under 'Analysis of findings'
31. Clarity of major themes	Were major themes clearly presented in the findings?	8-14	Results under 'Analysis of findings'
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	8-14	Results under 'Analysis of findings'

Perspectives of professional experts in relation to the development of community-based exercise for young adults with schizophrenia – A qualitative study

MF Andersen^{1,2*}, K Roed¹, A Riis^{2,3}, BS Rafn^{1,4}, BH Ebdrup^{5,6}, J Midtgaard^{1,6}

*Correspondence: MF Andersen, mfan@ucn.dk

¹ Centre for Applied Research in Mental Health Care (CARMEN), Mental Health Centre Glostrup, University of Copenhagen, Copenhagen, Denmark

² Department of Physiotherapy, University College of Northern Denmark, Aalborg, Denmark

Supplemental Material 2 Interview guide and additional methodological considerations

Standard opening and open-ended question:

According to you, as an expert, what should be considered when developing a community-based exercise programme for young adults with schizophrenia?

Expert specific question:

A:

- *With your practical experience in supervising exercise for people with schizophrenia, is there something you think non-health professional exercise instructors should pay close attention to? Motivation? Exercise content and structure?*

B:

- *With your knowledge of and experience in working with people with mental illness, what knowledge regarding pathophysiology should be shared with exercise instructors? Are there any contraindications that would inhibit people with psychotic disorders from participating in gym-based exercise? How should physical and mental complaints in relation to the exercise be handled?*

C:

- *With your knowledge of health psychology in relation to physical activity how would you create a motivating exercise environment outside the mental health system for people with mental illness? How do we avoid barriers towards change in physical active behavior?*

D:

- *With your practical experience in supervising exercise to people in antipsychotic treatment, is there something you think non-health professional exercise instructors should pay close attention to? Symptoms? Metabolic health?*

E:

- *With your experience in exercise programmes for people with mental illness, how would you create an inclusive and non-stigmatizing exercise environment? What should be considered in regard to the participants' social and personal recovery process?*

F:

- *With your knowledge of rehabilitation and recovery, what should be considered when delivering gym-based exercise to people in antipsychotic treatment in order to promote and support social and personal recovery? Are there any pitfalls that will work against recovery?*

G:

- *With your knowledge of and experience with physical activity in recovery processes, what would you consider when promoting exercise to people in antipsychotic treatment? Are there unique possibilities in using exercise instead of other social settings?*

H:

- *With your knowledge of exercise physiology, what should be considered to obtain clinically relevant exercise intensity and/or doses? How should the exercise content be organized? How should physical and mental complaints be handled during exercise?*

I, J, K:

- *With your knowledge of and clinical experience in working within early intervention outpatient clinics for people with schizophrenia spectrum disorders, how could an exercise community complement outpatient treatment?*

Additional methodological considerations

We chose online video interviews (Microsoft Teams) due to the geographical distance between experts and researchers, as well as due to the assumption that sharing professional expertise would not evoke strong emotions. Furthermore, data elicited via video calls have demonstrated substantial richness akin to that gained in a face-to-face interview (1).

We chose selective transcripts due to the fact that the analysis of the interviews was purely descriptive (not subject to in-depth analysis and interpretation). Moreover, the key informants were professional experts conveying professional knowledge and not personal experience, and we wished to concentrate on the specific aspects of the data that were most relevant to the research question rather than more latent content requiring more detailed transcripts.

No fieldnotes were made during or after the interview and interviews were not repeated.

No patients took part in planning, conduct or interpretation of the results of this study.

1. *Keen S, Lomeli-Rodriguez M, Joffe H. From Challenge to Opportunity: Virtual Qualitative Research During COVID-19 and Beyond. Int J Qual Methods [Internet]. 2022 May 1 [cited 2023 Jan 12];21:1–11. Available from: /pmc/articles/PMC9167989/*